

Introduction to Partial Differential Equations in  
Physics

Francesco Zanlungo



# Contents

<b>I</b>	<b>Introduction</b>	<b>9</b>
<b>1</b>	<b>Geometrical meaning of vector calculus</b>	<b>11</b>
1.1	A very short introduction to diffusion (and dimensional analysis) . . . . .	11
1.2	Curvilinear coordinates . . . . .	14
1.2.1	Cylindrical coordinates . . . . .	16
1.2.2	Spherical coordinates . . . . .	16
1.3	The gradient . . . . .	17
1.3.1	Gradient for cylindrical coordinates . . . . .	18
1.3.2	Gradient for spherical coordinates . . . . .	18
1.3.3	Dimensional analysis . . . . .	18
1.4	The divergence . . . . .	18
1.4.1	Definition . . . . .	18
1.4.2	Cartesian coordinates . . . . .	19
1.4.3	Gauss theorem . . . . .	21
1.4.4	Curvilinear coordinates . . . . .	22
1.4.5	Dimensional analysis . . . . .	24
1.5	The Laplacian . . . . .	25
1.5.1	Cartesian coordinates . . . . .	25
1.5.2	Cylindrical coordinates . . . . .	25
1.5.3	Spherical coordinates . . . . .	26
1.5.4	Dimensional analysis . . . . .	26
1.5.5	Geometrical meaning . . . . .	26
1.6	Another approach to diffusion . . . . .	27
<b>II</b>	<b>Separation of variables</b>	<b>31</b>
<b>2</b>	<b>Separation of variables in the heat equation</b>	<b>33</b>
2.1	Boundary conditions . . . . .	33

## CONTENTS

---

2.2	1D heat equation with fixed temperature ends: dimensional analysis . . . . .	37
2.3	Separation of variables for the 1D heat equation . . . . .	39
2.3.1	Separation of variables and linearity . . . . .	42
2.4	Fourier series and general solution to the 1D heat equation with homogeneous boundary conditions . . . . .	43
2.4.1	General solution . . . . .	43
2.5	Computation of coefficients . . . . .	44
2.6	Time dependence . . . . .	46
2.7	Flux boundary conditions . . . . .	47
2.7.1	Stationary solution . . . . .	47
2.7.2	General solution . . . . .	48
2.7.3	Computation of coefficients . . . . .	50
2.8	Diffusion on the circle . . . . .	51
2.8.1	Periodic boundary conditions . . . . .	51
2.8.2	Separation of variables . . . . .	52
2.8.3	Complex “orthogonal” solutions . . . . .	54
2.8.4	General solution . . . . .	55
<b>3</b>	<b>The Schrödinger equation</b>	<b>57</b>
3.1	Historical remarks . . . . .	57
3.1.1	Plank . . . . .	57
3.1.2	Einstein . . . . .	58
3.1.3	Relativity . . . . .	59
3.1.4	De Broglie . . . . .	61
3.1.5	Heisenberg . . . . .	62
3.1.6	Schrödinger . . . . .	63
3.2	A simple 1D problem in Quantum mechanics: a trapped particle	65
3.2.1	Boundary conditions . . . . .	65
3.2.2	Separation of variables: the Time-independent Schrödinger equation . . . . .	65
3.2.3	Solution of the time-independent Schrödinger equation	66
3.2.4	General solution . . . . .	68
3.3	The Philosophy of Quantum Mechanics . . . . .	68
3.4	Getting rid of $\hbar$ . . . . .	71
3.5	The particle in a 3D box . . . . .	73
<b>4</b>	<b>Vibrating membrane</b>	<b>77</b>
4.1	The wave equation . . . . .	77
4.1.1	A “deduction” . . . . .	77

4.1.2	From string to membrane . . . . .	79
4.1.3	Getting rid of $c$ . . . . .	80
4.2	Separation of variables in the vibrating membrane problem . . . . .	81
<b>5</b>	<b>Laplace equation</b>	<b>85</b>
5.1	Introduction . . . . .	85
5.1.1	Electrostatics . . . . .	85
5.1.2	Unicity of solution for the Laplace equation . . . . .	88
5.2	2D Laplace equation in a rectangular area . . . . .	89
5.2.1	A simpler problem . . . . .	90
5.2.2	Solution to the full problem . . . . .	91
5.3	Laplace equation in polar coordinates . . . . .	92
<b>6</b>	<b>A short introduction to function spaces and Dirac formalism</b>	<b>97</b>
6.1	Introduction . . . . .	97
6.2	Vector spaces . . . . .	98
6.2.1	Basic geometry and physics . . . . .	98
6.2.2	Generalisation . . . . .	99
6.2.3	The scalar or inner product . . . . .	101
6.2.4	Linear independence, bases, dimension and the like . . . . .	102
6.2.5	Orthonormalisation . . . . .	103
6.2.6	Scalar product and orthonormal bases . . . . .	104
6.3	Linear operators . . . . .	105
6.3.1	Definition . . . . .	105
6.3.2	Linear operators on a finite dimensional space . . . . .	106
6.3.3	Eigenvalues and eigenvectors . . . . .	107
6.3.4	Inverse operator . . . . .	108
6.3.5	Adjoint operator . . . . .	108
6.3.6	Commuting operators . . . . .	111
6.3.7	Normal operators . . . . .	113
6.3.8	Unitary operator . . . . .	114
6.4	Dirac notation . . . . .	115
6.4.1	Bras, kets, and scalar products . . . . .	115
6.4.2	Ketbras, projectors and representations of unity . . . . .	116
6.4.3	Operators . . . . .	117
6.5	Function spaces . . . . .	120
6.5.1	Introduction . . . . .	120
6.5.2	Continuum limit . . . . .	121
6.5.3	Adjoint operator . . . . .	122
6.5.4	Enumerable basis . . . . .	123

## CONTENTS

---

<b>7</b>	<b>Laplace equation in spherical coordinates</b>	<b>125</b>
7.1	Introduction . . . . .	125
7.2	Axial symmetry . . . . .	125
7.2.1	Radial equation . . . . .	126
7.2.2	Angular equation . . . . .	126
7.2.3	Legendre polynomials . . . . .	128
7.2.4	General solution . . . . .	130
7.3	Introduction to the general case . . . . .	131
<b>8</b>	<b>Orthogonality (an invitation to Sturm-Liouville theory)</b>	<b>135</b>
8.1	An other approach to orthogonality . . . . .	135
8.2	The Sturm-Liouville problem . . . . .	137
8.2.1	Definition . . . . .	137
8.2.2	Scalar product . . . . .	137
8.2.3	Adjoint operator . . . . .	138
8.2.4	Defining $w$ so that $\hat{L}$ is self-adjoint . . . . .	138
8.2.5	A different formulation of the problem . . . . .	141
8.2.6	Orthogonality . . . . .	142
8.3	Application to Legendre polynomials . . . . .	143
<b>9</b>	<b>The hydrogen atom</b>	<b>145</b>
9.1	Introduction . . . . .	145
9.2	The equation . . . . .	146
9.3	Time independent Schrödinger equation . . . . .	147
9.4	The angular solution . . . . .	148
9.5	The radial equation . . . . .	149
9.5.1	First change of variable . . . . .	149
9.5.2	Dimensional analysis (characteristic length) . . . . .	150
9.5.3	Second change of variables . . . . .	151
9.5.4	A further dimensional analysis (characteristic energy) . . . . .	151
9.5.5	Asymptotic behaviour . . . . .	152
9.5.6	The equation for the function $f(u)$ . . . . .	153
9.5.7	A third change of variables . . . . .	154
9.5.8	Power series . . . . .	155
9.5.9	Conditions imposed by the power series solution . . . . .	156
9.6	The solution . . . . .	157
	<b>Appendices</b>	<b>159</b>
9.A	A very shorth introduction to analytical mechanics . . . . .	159
9.A.1	Cartesian coordinates . . . . .	159

---

## CONTENTS

---

9.A.2	Arbitrary coordinates . . . . .	161
9.B	Two body problem . . . . .	163
9.B.1	Introduction . . . . .	163
9.B.2	Newtonian treatment . . . . .	163
9.B.3	Lagrangian treatment . . . . .	165
9.C	Angular momentum in Classical and Quantum Mechanics . . . . .	167
9.C.1	Definitions . . . . .	167
9.C.2	Lagrangian for a central potential . . . . .	168
9.C.3	Angular momentum in Quantum Mechanics . . . . .	171
9.C.4	Commutators in Quantum Mechanics . . . . .	173
9.C.5	The operators $\hat{L}_+$ , $\hat{L}_-$ and their properties . . . . .	180
9.C.6	Eigenvectors and eigenvalues of $\hat{L}$ . . . . .	181
9.D	Explicit form of $\hat{L}^2$ in spherical coordinates . . . . .	187
9.D.1	Change of coordinate . . . . .	187
9.D.2	Partial derivatives . . . . .	187
9.D.3	Components of $\hat{L}$ . . . . .	189
9.E	Mathematical formulation of the uncertainty principle . . . . .	191
9.E.1	Expectation values . . . . .	191
9.E.2	Schwartz inequality . . . . .	195
9.E.3	Physical meaning . . . . .	196

## CONTENTS

---



## Part I

# Introduction



# Chapter 1

## Geometrical meaning of vector calculus

### 1.1 A very short introduction to diffusion (and dimensional analysis)

Let us consider the function

$$\rho(t, \mathbf{x}) \tag{1.1}$$

giving, for example, the density of a substance in still air as a function of time  $t$  and position  $\mathbf{x}$  (a 3D vector). We may assume that at time  $t = 0$  such density assumes a given distribution

$$\rho(0, \mathbf{x}) = \rho_0(\mathbf{x}) \tag{1.2}$$

and wonder how  $\rho$  will change with time. Let us make the hypothesis that at fixed  $\mathbf{x} = \mathbf{x}_0$  the variation in time of  $\rho$  will depend on how much material is in  $\mathbf{x}_0$  with respect to the neighbouring space. We may define a small volume  $V_{\mathbf{x}_0}$  centred in  $\mathbf{x}_0$  and measure the average of  $\rho$  in the volume

$$\bar{\rho}(t, \mathbf{x}_0) = \frac{\int_{V_{\mathbf{x}_0}} \rho(t, \mathbf{x}) d\mathbf{x}}{\int_{V_{\mathbf{x}_0}} d\mathbf{x}} \tag{1.3}$$

This quantity depends obviously on the details of the choice of  $V_{\mathbf{x}_0}$ , but we assume that it is uniquely defined if we let the size of  $V_{\mathbf{x}_0}$  go to zero,

$$\bar{\rho}(t, \mathbf{x}_0) = \lim_{V_{\mathbf{x}_0} \rightarrow 0} \frac{\int_{V_{\mathbf{x}_0}} \rho(t, \mathbf{x}) d\mathbf{x}}{\int_{V_{\mathbf{x}_0}} d\mathbf{x}} \tag{1.4}$$

In a intuitive way, we may expect the amount of  $\rho$  to grow in time if  $\bar{\rho} > \rho$ , and to decrease if  $\bar{\rho} < \rho$ . Namely, the substance will diffuse in  $\mathbf{x}_0$  if a larger

## 1.1 A very short introduction to diffusion (and dimensional analysis)

---

amount of it is in the surrounding small volume, and diffuse out of  $\mathbf{x}_0$  if a smaller amount of it is in the volume.

Mathematically, we may express the change of  $\rho$  in time while keeping  $\mathbf{x}$  fixed as the partial derivative in time

$$\frac{\partial \rho(t, \mathbf{x})}{\partial t} \equiv \partial_t \rho(t, \mathbf{x}) \equiv \partial_t \rho \quad (1.5)$$

In these notes, we will often use  $\partial_t$  as a short hand for the partial derivative with respect to variable  $t$ , and we will not write the list of variables when there is no risk of confusion.

We now assume that this variation in time is proportional to the difference between  $\rho$  and  $\bar{\rho}$

$$\partial_t \rho(t, \mathbf{x}) = \kappa(\bar{\rho}(t, \mathbf{x}) - \rho(t, \mathbf{x})) \quad (1.6)$$

In this equation, we used  $\mathbf{x}$  in place of  $\mathbf{x}_0$  to stress that the equation is valid everywhere<sup>1</sup>.

We note that the constant has “the dimension of inverse time”, i.e., if time is measured in seconds,  $\kappa$  is measured in  $\text{s}^{-1}$ . This is due to the fact that on the left  $\rho$  is divided by time<sup>2</sup>, while on the right  $\rho$  is multiplied by  $\kappa$ . As a result, if we want the equation to make sense if we change units,  $\kappa$  has to be the equivalent to a division by time.

Such a “dimensional analysis” is extremely useful in all physics problems, so that it may be useful to explain it better with a simple example. Let us assume we have a law<sup>3</sup> that relates the average velocity of walking  $v$  to the walked distance  $L$ ,

$$v = \alpha/\sqrt{L} \quad (1.7)$$

Let us also assume that we know that if we measure distances in kilometres, and times in hours, we have  $\alpha = 7.2$ , meaning that we walk at 7.2 km/h over a 1 km distance. The law will then tell us that we walk at a 3.6 km/h velocity over a 4 km distance. Let us now use meters and seconds. 7.2 km equal 7200 m, an 1 hour is 3600 s, so that a velocity of 7.2 km/h corresponds to 2 m/s. We have now  $2 = \alpha/\sqrt{1000}$  or  $\alpha \approx 63.25$ . It is clear that  $\alpha$  is not a pure number, i.e. a number that does not change when we change a unit system. This is obvious, since on the left we have a velocity, and on the right  $\alpha$  divided by the square of a length, so that  $\alpha$  must have its own “dimension”, i.e. it must be expressed as a function of length, time, mass,

---

<sup>1</sup>Or at least inside a proper domain in which we study the evolution of  $\rho$ .

<sup>2</sup>Since the (partial) derivative is given by the difference between two values of  $\rho$  at small time separation  $\delta t$ , divided by  $\delta t$ , or, more precisely given by  $\lim_{\delta t \rightarrow 0} \frac{\rho(t+\delta t, \mathbf{x}) - \rho(t, \mathbf{x})}{\delta t}$ .

<sup>3</sup>The actual law will be more complex, and dependent on individual fitness.

etc, so that also the right side will be a velocity. A velocity has to be equal to a velocity!<sup>4</sup>

We may write an equation as

$$[V] = \frac{[L]}{[T]} = [L][T]^{-1} \quad (1.8)$$

that we read as “velocity has the dimensionality of length over time”. This means that the physical quantity velocity is obtained as the ratio between a length and a time. Then, the dimensional equation corresponding to eq. (1.7) is

$$[L][T]^{-1} = [\alpha][L]^{-1/2} \quad (1.9)$$

that we solve as

$$[\alpha] = [L]^{3/2}[T]^{-1} \quad (1.10)$$

This means that  $\alpha$  is measured in square roots of volumes divided by time. This looks quite complex, but as we know that a quantity measured in kilometres becomes  $10^3$  times bigger when we measure it in meters (i.e., when we pass to a unit that is  $10^{-3}$  smaller)  $\alpha$  will be scaled of a factor

$$(10^3)^{3/2}(3.6 \cdot 10^3)^{-1} = 10^{3/2}/3.6 \approx 8.784$$

passing from kilometres and hours to meters and seconds. Indeed,

$$7.2 \cdot 8.748 \approx 63.25$$

Let us go back to our original problem, eq. (1.6). In this form, the equation is very intuitive, and may already tell us something interesting. If, at time  $t = 0$ , we have  $\rho(0, \mathbf{x}) \geq 0 \forall \mathbf{x}$ , as we expect for the density of our substance, we will have  $\rho(t, \mathbf{x}) \geq 0 \forall \mathbf{x}, t \geq 0$ . Why? To have a negative  $\rho$ , we need (by eq. 1.6 and assuming the function and its derivatives to be continuous) a point with  $\rho = 0$  and with  $\bar{\rho} < 0$ . This means that starting from  $\rho \geq 0$  everywhere, we will never get a negative  $\rho$ .

Nevertheless, eq. (1.6) is written in a very peculiar way. It relates a partial derivative to the limit of an integral. It is possible to re-write it as an equation involving only partial derivative of  $\rho$ , i.e. as a *partial differential equation*. In our path of obtaining such a *diffusion equation* (or *heat equation*) we will also learn the geometrical and physical meaning of vector calculus, and how to write differential equations in different coordinate systems.

---

<sup>4</sup>So that the above expression  $2 = \alpha/\sqrt{1000}$  is actually not correct, or better valid only after a specific set of units has been chosen.

## 1.2 Curvilinear coordinates

---

### 1.2 Curvilinear coordinates

Let us consider two points on the plane,  $A$  and  $B$ . We assume their coordinates in a given Cartesian frame to be  $A(x, y)$  and  $B(x + \Delta x, y + \Delta y)$ . By Pythagoras' theorem, we know that the distance between the two points is

$$\Delta s = \sqrt{\Delta x^2 + \Delta y^2} \quad (1.11)$$

We will often write this as

$$ds^2 = dx^2 + dy^2 \quad (1.12)$$

meaning that  $dx$  and  $dy$  are small and we are authorised to retain only leading terms in Taylor polynomials and the like.

We can also use other coordinates in the plane, e.g. polar coordinates  $(r, \theta)$ . The relation giving the Cartesian coordinates as functions of the polar ones is

$$x = r \cos \theta, \quad y = r \sin \theta \quad (1.13)$$

Using a Taylor polynomial, we may write the change in  $x$  as a function of changes in  $r$  and  $\theta$ , namely

$$\Delta x = \frac{\partial x}{\partial \theta} \Delta \theta + \frac{\partial x}{\partial r} \Delta r + \text{higher powers of } \Delta r, \Delta \theta \quad (1.14)$$

As stated above, assuming variations to be small we may also write this expression as

$$dx = \frac{\partial x}{\partial \theta} d\theta + \frac{\partial x}{\partial r} dr \quad (1.15)$$

We also have

$$dy = \frac{\partial y}{\partial \theta} d\theta + \frac{\partial y}{\partial r} dr \quad (1.16)$$

From eq. (1.13) we have

$$\frac{\partial x}{\partial r} = \cos \theta, \quad \frac{\partial x}{\partial \theta} = -r \sin \theta, \quad \frac{\partial y}{\partial r} = \sin \theta, \quad \frac{\partial y}{\partial \theta} = r \cos \theta \quad (1.17)$$

and substitution of eqs. (1.13,1.17) in eq. (1.12) gives

$$\begin{aligned} ds^2 &= \cos^2 \theta dr^2 + r^2 \cos^2 \theta d\theta^2 - 2r \cos \theta \sin \theta + \sin^2 \theta dr^2 + r^2 \sin^2 \theta d\theta^2 + 2r \sin \theta \cos \theta \\ &= r^2 d\theta^2 + dr^2 \end{aligned} \quad (1.18)$$

This equation tells us two important things

1.  $ds^2$  may be written as a sum of quadratic terms in  $dr$  and  $d\theta$ , i.e. the polar coordinates are orthogonal and no mixed terms appear

2. the expression is anyway different from the one we obtained for Cartesian coordinates, since the contribution of  $d\theta$  is weighted by a  $r^2$  term. The reason of this should be evident from Fig. 1.1. For larger values of  $r$ , to same variation of  $\theta$  corresponds a larger distance




Figure 1.1: A variation in  $\theta$  causes a larger displacement for larger  $r$  values.

In general, a point in 3D space may be identified by Cartesian coordinates  $(x, y, z)$  or more general *curvilinear coordinates*  $(q_1, q_2, q_3)$ . We will have a relation expressing the former as a function for the latter

$$x(q_1, q_2, q_3), \quad y(q_1, q_2, q_3), \quad z(q_1, q_2, q_3) \quad (1.19)$$

We may also use the notation  $x = x_1, y = x_2, z = x_3$ , and write sometimes the relations 1.19 with the short hand  $x_i(q_j)$ .

We have

$$ds^2 = \sum_i dx_i^2 = \sum_{i,j,k} \left( \frac{\partial x_i}{\partial q_j} dq_j \right) \left( \frac{\partial x_i}{\partial q_k} dq_k \right) = \sum_{j,k} h_{j,k} dq_j dq_k \quad (1.20)$$

where we defined

$$h_{j,k} \equiv \sum_i \frac{\partial x_i}{\partial q_j} \frac{\partial x_i}{\partial q_k} \quad (1.21)$$

When, as in the case of polar coordinates, the variations in the  $q_i$  are orthogonal between them, the expression above simplifies to

$$ds^2 = \sum_i h_i^2 dq_i^2 \quad (1.22)$$

The  $h_i$  are clearly functions of the  $q_i$ , so it is useful to write explicitly

$$ds^2 = h_1^2(q_1, q_2, q_3) dq_1^2 + h_2^2(q_1, q_2, q_3) dq_2^2 + h_3^2(q_1, q_2, q_3) dq_3^2 \quad (1.23)$$

## 1.2 Curvilinear coordinates

---

In these notes we will focus on coordinates that allow us to use an expression in the form 1.23, and in particular on two useful coordinate systems;

### 1.2.1 Cylindrical coordinates

This coordinate system corresponds to choosing a plane, identified by Cartesian coordinates  $(x, y)$ , and express it in polar coordinates  $(r, \theta)$ , while the  $z$  coordinate is left unchanged. The Cartesian coordinates are expressed as function of the cylindrical ones through

$$x = r \cos \theta, \quad y = r \sin \theta, \quad z = z \quad (1.24)$$

and it is straightforward to show that

$$ds^2 = dr^2 + r^2 d\theta^2 + dz^2 \quad (1.25)$$

so that

$$h_r = 1(= h_1), \quad h_\theta = r(= h_2), \quad h_z = 1(= h_3) \quad (1.26)$$

We will use this system of coordinates when we are facing a problem with a particular symmetry, namely rotation around a given axis (identified by the  $z$  coordinate).

### 1.2.2 Spherical coordinates

This coordinate system is used when we have a spherical symmetry in our problem, and somehow correspond to use a combination of two polar system. First we divide the component on the  $z$  axis from the component on the  $(x, y)$  plane by using  $r$  and the angle  $\theta$ , and then we divide the  $x$  and  $y$  components introducing another angle,  $\varphi$ . In this “second application” of the polar transformation, the role of the distance is played by the component of  $r$  in the plane,  $r \sin \theta$ . The Cartesian coordinates are expressed as functions of the spherical ones through

$$x = r \sin \theta \cos \varphi, \quad y = r \sin \theta \sin \varphi, \quad z = r \cos \theta \quad (1.27)$$

Both from explicit computation or geometrical considerations (the two polar systems described above, one using  $r$  and the other one using  $r \sin \theta$  as distance coordinates) we get

$$ds^2 = dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 \quad (1.28)$$

so that

$$h_r = 1(= h_1), \quad h_\theta = r(= h_2), \quad h_\varphi = r \sin \theta(= h_3) \quad (1.29)$$



### 1.3 The gradient

The geometrical definition of the gradient operator  $\nabla$  is the following. Let us consider a function  $\phi$  and its values in points  $A$  and  $B$ . Let us assume the separation between  $A$  and  $B$  being given by the vector

$$\Delta \mathbf{s} \equiv B - A \quad (1.30)$$

and define

$$\Delta \phi = \phi(B) - \phi(A) \quad (1.31)$$

Then we ask the gradient to satisfy

$$\Delta \phi = \nabla \phi \cdot \Delta \mathbf{s} + \text{higher powers of the components of } \Delta \mathbf{s} \quad (1.32)$$

In Cartesian coordinates (2D) we have

$$\nabla \phi \cdot \Delta \mathbf{s} = (\nabla \phi)_x \Delta x + (\nabla \phi)_y \Delta y \quad (1.33)$$

At the same time, in any coordinate system we have the Taylor expansion

$$\Delta \phi = \sum_i \frac{\partial \phi}{\partial q_i} \Delta q_i + \text{higher powers of } \Delta q_i \quad (1.34)$$

so that by comparison of eqs. (1.32,1.34) we obtain, for Cartesian coordinates,

$$(\nabla \phi)_i = \frac{\partial \phi}{\partial x_i} \quad (1.35)$$

But this relation will not hold in curvilinear coordinates, if we want eq. (1.32) to hold when the displacement will have a less trivial dependence on coordinates (as in eq. 1.23).

Let us consider first the 2D polar case. The same  $B - A = \Delta \mathbf{s}$  will be now given by (again, keeping only first order terms)

$$(\Delta \mathbf{s})_r = \Delta r, \quad (\Delta \mathbf{s})_\theta = r \Delta \theta \quad (1.36)$$

where by  $(\Delta \mathbf{s})_q$  we mean component of the vector along the variation of coordinate  $q$ . As a result, comparing eqs. (1.32,1.34) we get

$$(\nabla \phi)_r \Delta r + (\nabla \phi)_\theta r \Delta \theta \approx \Delta \phi \approx \frac{\partial \phi}{\partial r} \Delta r + \frac{\partial \phi}{\partial \theta} \Delta \theta \quad (1.37)$$

and

$$(\nabla \phi)_r = \frac{\partial \phi}{\partial r}, \quad (\nabla \phi)_\theta = \frac{1}{r} \frac{\partial \phi}{\partial \theta} \quad (1.38)$$

The  $\theta$  component of the gradient becomes smaller for large  $r$  to compensate for the growth of the displacement  $(\Delta \mathbf{s})_\theta$ .

## 1.4 The divergence

---

In general, we will have for orthogonal coordinates

$$(\Delta \mathbf{s})_i = h_i(q_j) \Delta q_i \quad (1.39)$$

so that by comparing eqs. (1.32,1.34) we get

$$(\nabla \phi)_i = \frac{1}{h_i(q_j)} \frac{\partial \phi}{\partial q_i} \quad (1.40)$$

### 1.3.1 Gradient for cylindrical coordinates

$$(\nabla \phi)_r = \frac{\partial \phi}{\partial r}, \quad (\nabla \phi)_\theta = \frac{1}{r} \frac{\partial \phi}{\partial \theta}, \quad (\nabla \phi)_z = \frac{\partial \phi}{\partial z} \quad (1.41)$$

### 1.3.2 Gradient for spherical coordinates

$$(\nabla \phi)_r = \frac{\partial \phi}{\partial r}, \quad (\nabla \phi)_\theta = \frac{1}{r} \frac{\partial \phi}{\partial \theta}, \quad (\nabla \phi)_\varphi = \frac{1}{r \sin \theta} \frac{\partial \phi}{\partial z} \quad (1.42)$$

### 1.3.3 Dimensional analysis

A gradient is physically a spatial derivative, so that its dimension should be

$$[\phi][L]^{-1} \quad (1.43)$$

Angles such as  $\theta$  and  $\varphi$  are pure numbers. The correct dimensionality in eqs. (1.41,1.42) is given anyway by the division by  $r$  in front of derivations that involve angles. The dimensional analysis suggests thus the correctness of our results.

## 1.4 The divergence

### 1.4.1 Definition

We consider now the vector field (i.e., a function from  $\mathbb{R}^3$  to  $\mathbb{R}^3$ , assigning a vector to each point in space)  $\mathbf{A}(\mathbf{x})$ . We geometrically define its divergence as

$$\operatorname{div} \mathbf{A}(\mathbf{x}_0) \equiv \nabla \cdot \mathbf{A}(\mathbf{x}_0) = \lim_{V_{\mathbf{x}_0} \rightarrow 0} \frac{\int_{S_{V_{\mathbf{x}_0}}} \mathbf{A}(\mathbf{x}) \cdot \mathbf{n} dS}{\int_{V_{\mathbf{x}_0}} d\mathbf{x}} \quad (1.44)$$

A few explanations are due. First of all, the notation  $\nabla \cdot \mathbf{A}$  for the divergence recalls us of the usual Cartesian expression  $\sum_i \partial_i A_i$ . Nevertheless, in these note we prefer to write  $\operatorname{div}$  to recall the geometrical definition and the different forms assumed in different coordinate systems. As we did above for eq. (1.4), we define an integral in a small volume  $V_{\mathbf{x}_0}$  centred around

$\mathbf{x}_0$ , and let this volume go to zero (see below for a practical example of how this is done).  $S_{V_{\mathbf{x}_0}}$  is the surface delimiting the volume,  $\mathbf{n}$  is the unit vector that is orthogonal in each point to the surface, and  $\mathbf{A}(\mathbf{x}) \cdot \mathbf{n} dS$  stands for the integration of the scalar product of the vectors  $\mathbf{A}$  and  $\mathbf{n}$ . We are thus measuring on the whole surface the outgoing component of  $\mathbf{A}$ , and then dividing by the volume. This definition originates from fluid dynamics (how much fluid is flowing out of a volume).

### 1.4.2 Cartesian coordinates

Let us do this for the Cartesian coordinates. We operate on a small cube centred in  $(x_0, y_0, z_0)$  with sides  $\Delta x = \Delta y = \Delta z = a$  (Fig 1.2). We have




Figure 1.2: Small cube for divergence integral.

obviously

$$\int_{V_{\mathbf{x}_0}} d\mathbf{x} = \Delta x \Delta y \Delta z = a^3 \quad (1.45)$$

The limit  $V_{\mathbf{x}_0} \rightarrow 0$  corresponds to sending  $a \rightarrow 0$ <sup>5</sup>.

We now need to integrate  $\mathbf{A}(\mathbf{x}) \cdot \mathbf{n}$  on all faces of the cube. Let us name this faces  $1, \dots, 6$  and the corresponding integrals  $I_1, \dots, I_6$ . If, as in Fig. 1.2 we name the face in the direction of growing  $x$  as 1, we have (due to

---

<sup>5</sup>If we use different values of  $\Delta x, \Delta y, \Delta z$ , the derivation would be unchanged provided that we send each of these to zero. Our proof is actually performed keeping explicit values of these variations, so that the need to send all of them to zero results evident. One way of getting the limit is to multiply the possibly different  $\Delta x_i$  by  $a$ , and then send  $a$  to 0.

## 1.4 The divergence

---

$$\mathbf{A} \cdot \mathbf{n}_1 = A_x)$$

$$I_1 = \int_{-\frac{\Delta z}{2}}^{\frac{\Delta z}{2}} dz \int_{-\frac{\Delta y}{2}}^{\frac{\Delta y}{2}} A_x \left( x_0 + \frac{\Delta x}{2}, y_0 + y, z_0 + z \right) dy \quad (1.46)$$

Now we use the Taylor expansion

$$\begin{aligned} & A \left( x_0 + \frac{\Delta x}{2}, y_0 + y, z_0 + z \right) = \\ & = A \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) + \partial_y A_x \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) y + \\ & \quad \partial_z A_x \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) z + O(y^2, z^2) \end{aligned} \quad (1.47)$$

where  $O(y^2, z^2)$  stands for terms like  $x^2$ ,  $y^2$ ,  $zy$  or higher powers. If we compute an integral such as

$$\bar{I} = \int_{-\frac{\Delta z}{2}}^{\frac{\Delta z}{2}} dz \int_{-\frac{\Delta y}{2}}^{\frac{\Delta y}{2}} C y^2 dy \quad (1.48)$$

we obtain (using for simplicity  $a$  for the value of all sides)

$$\bar{I} = C \frac{a^4}{12} \quad (1.49)$$

Once we divide this contribution by the volume  $a^3$ , we get  $Ca/12 \rightarrow 0$  as  $a \rightarrow 0$ . This is true of all quadratic terms (and higher). A quadratic term, integrated on the surface, will give a quartic term that will go to zero when divided by the volume.

We may now consider

$$\begin{aligned} I^* &= \int_{-\frac{\Delta z}{2}}^{\frac{\Delta z}{2}} dz \int_{-\frac{\Delta y}{2}}^{\frac{\Delta y}{2}} \partial_y A_x \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) y dy = \\ &= \partial_y A_x \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) \int_{-\frac{\Delta z}{2}}^{\frac{\Delta z}{2}} dz \int_{-\frac{\Delta y}{2}}^{\frac{\Delta y}{2}} y dy = 0 \end{aligned} \quad (1.50)$$

since

$$\int_{-\frac{\Delta x}{2}}^{\frac{\Delta x}{2}} y dy = \frac{a^2}{8} - \frac{a^2}{8} \quad (1.51)$$

Namely we are integrating an odd function on a symmetrical interval, so we get 0. The same will happen for the integral involving  $\partial_z A_x(x_0, y_0, z_0)z$ . Taking in account the only remaining term we have

$$I_1 = A_x \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) \Delta y \Delta z + O(a^4) \quad (1.52)$$

In a similar way

$$I_2 = -A_x \left( x_0 - \frac{\Delta x}{2}, y_0, z_0 \right) \Delta y \Delta z + O(a^4) \quad (1.53)$$

the different sign being due to  $\mathbf{A} \cdot \mathbf{n}_2 = -A_x$ .

Now we have

$$\begin{aligned} I_1 + I_2 &= \\ &= \left[ A_x \left( x_0 + \frac{\Delta x}{2}, y_0, z_0 \right) - A_x \left( x_0 - \frac{\Delta x}{2}, y_0, z_0 \right) \right] \Delta y \Delta z + O(a^4) = \\ &= \left[ A_x(x_0, y_0, z_0) + \partial_x A_x(x_0, y_0, z_0) \frac{\Delta x}{2} \right. \\ &\quad \left. - \left( A_x(x_0, y_0, z_0) - \partial_x A_x(x_0, y_0, z_0) \frac{\Delta x}{2} \right) + O(a^2) \right] \Delta y \Delta z + O(a^4) = \\ &= \partial_x A_x(x_0, y_0, z_0) \Delta x \Delta y \Delta z + O(a^4) \end{aligned} \quad (1.54)$$

We may repeat this for the other faces of the cube just changing names of the variables, and we get

$$\operatorname{div} \mathbf{A}(\mathbf{x}) = \lim_{a \rightarrow 0} \frac{[\partial_x A_x(\mathbf{x}) + \partial_y A_y(\mathbf{x}) + \partial_z A_z(\mathbf{x})] \Delta x \Delta y \Delta z + O(a^4)}{\Delta x \Delta y \Delta z} \quad (1.55)$$

and thus, in *Cartesian coordinates*

$$\operatorname{div} \mathbf{A} = \sum_i \partial_i A_i \quad (1.56)$$

### 1.4.3 Gauss theorem

From the (coordinate independent) definition eq. (1.44) we have, for an infinitesimal cubic volume  $\delta V$

$$\operatorname{div} \mathbf{A}(\mathbf{x}) \delta V = \int_{S_{\delta V}} \mathbf{A}(\mathbf{x}) \cdot \mathbf{n} dS \quad (1.57)$$

The integral over a finite volume may be defined as the integral over a partition of very small cubes, in the limit of the volume of all cubes going to zero. Namely, for a finite volume  $V$  we have

$$\int_V \operatorname{div} \mathbf{A}(\mathbf{x}) d\mathbf{x} = \sum_i \operatorname{div} \mathbf{A}(\mathbf{x}) \delta V_i = \sum_i \int_{S_{\delta V_i}} \mathbf{A}(\mathbf{x}) \cdot \mathbf{n} dS \quad (1.58)$$

In general, the sum of the integrals over the surfaces  $S_{\delta V_i}$  of the cubes that give the total volume  $V$  would be different from an integral over the

---

## 1.4 The divergence

---

surface  $S_V$ , due to the contribution of the internal surfaces of the cubes. But due to the nature of the integrand, all internal terms will be present in two neighbouring cubes that have opposite  $\mathbf{n}$ , and thus all these terms will cancel, leaving only the integral over the external surface. This result is called Gauss theorem

$$\int_V \operatorname{div} \mathbf{A}(\mathbf{x}) d\mathbf{x} = \int_{S_V} \mathbf{A}(\mathbf{x}) \cdot \mathbf{n} dS \quad (1.59)$$

### 1.4.4 Curvilinear coordinates

To understand how to generalise in the curvilinear coordinate case, let us check the cylindrical case described in Fig. 1.3. Our small volume will be




Figure 1.3: Small volume for divergence integral in the cylindrical coordinate case.

now characterised by having faces that are surfaces with constant  $r$ ,  $\theta$  or  $z$ . In each point of 3D space, vectors orthogonal to such surfaces define a set of orthonormal vectors, that we may call  $\mathbf{e}_r$ ,  $\mathbf{e}_\theta$ ,  $\mathbf{e}_z$ . These vectors (with the exception of  $\mathbf{e}_z$ ) are different in different point of space, but at any given point they define a basis for the vector space at the location. Thus any vector  $\mathbf{A}$  may be written as

$$\mathbf{A} = A_r \mathbf{e}_r + A_\theta \mathbf{e}_\theta + A_z \mathbf{e}_z \quad (1.60)$$

We may name again the faces of our small volume from 1 to 6, and for example, as shown in Fig. 1.3, name 1 and 2 the faces at constant  $r$ . We

will clearly have  $\mathbf{n}_1 = \mathbf{e}_r$  and  $\mathbf{n}_2 = -\mathbf{e}_r$ . We may repeat the operations performed above in the Cartesian case remembering that all considerations involving Taylor expansions will be unchanged since for a general function  $g(q)$

$$\Delta g = \partial_q g \Delta q \quad (1.61)$$

applies also to curvilinear coordinates. On the other hand, integration in a curvilinear coordinate has to be multiplied by  $h_q$  to take in account the fact that the displacement in the curvilinear direction is  $h_q \Delta q$ <sup>6</sup>. For example, an integral in  $\theta$  will involve a multiplication by the locally constant value of  $r$ .

If our volume is centred in  $(r_0, \theta_0, z_0)$  we have

$$I_1 = A_r \left( r_0 + \frac{\Delta r}{2}, \theta_0, z_0 \right) \left[ r_0 + \frac{\Delta r}{2} \right] \Delta \theta \Delta z + O(a^4) \quad (1.62)$$

and

$$I_2 = A_r \left( r_0 - \frac{\Delta r}{2}, \theta_0, z_0 \right) \left[ r_0 - \frac{\Delta r}{2} \right] \Delta \theta \Delta z + O(a^4) \quad (1.63)$$

If we now consider the auxiliary function

$$f_r(r, \theta, z) = A_r(r, \theta, z)r \quad (1.64)$$

we may write

$$\begin{aligned} I_1 + I_2 &= \left[ f_r \left( r_0 + \frac{\Delta r}{2}, \theta_0, z_0 \right) - f_r \left( r_0 - \frac{\Delta r}{2}, \theta_0, z_0 \right) \right] \Delta \theta \Delta z + O(a^4) = \\ &= \partial_r f(r_0, \theta_0, z_0) \Delta r \Delta \theta \Delta z + O(a^4) \end{aligned} \quad (1.65)$$

The contribution of these two faces to the divergence is thus

$$\lim_{a \rightarrow 0} \frac{I_1 + I_2}{\int_{V_{\mathbf{x}_0}} d\mathbf{x}} = \lim_{a \rightarrow 0} \frac{I_1 + I_2}{r_0 \Delta r \Delta \theta \Delta z} = \frac{1}{r_0} \partial_r (r A_r(r_0, \theta_0, z)) \quad (1.66)$$

How can we generalise to other curvilinear systems? The volume will be given in general by

$$\int_{V_{\mathbf{x}_0}} d\mathbf{x} = h_1(q_1, q_2, q_3) h_2(q_1, q_2, q_3) h_3(q_1, q_2, q_3) \quad (1.67)$$

---

<sup>6</sup>Obviously, this result agrees with the more formal and rigorous one that performs the integration taking in account the Jacobian determinant of the coordinate transformation. Such a determinant is indeed trivially  $h_1 h_2 h_3$  for orthogonal coordinates.

## 1.4 The divergence

---

while the auxiliary function  $f_i$  generated by the integral on the two faces at constant  $q_i$  has contribution  $h_j$  with  $j \neq i$  to the integral, so that

$$f_i(q_j) = A_i(q_j) \prod_{j \neq i} h_j(q_k) \quad (1.68)$$

and finally

$$\operatorname{div} \mathbf{A} = \frac{\sum_i \partial_i f_i}{\prod_i h_i} \quad (1.69)$$

or, in an explicit way

$$\begin{aligned} \operatorname{div} \mathbf{A}(q_1, q_2, q_3) &= \frac{1}{h_1(q_1, q_2, q_3) h_2(q_1, q_2, q_3) h_3(q_1, q_2, q_3)} \\ &\quad \left[ \partial_1 [A_1(q_1, q_2, q_3) h_2(q_1, q_2, q_3) h_3(q_1, q_2, q_3)] + \right. \\ &\quad + \partial_2 [A_2(q_1, q_2, q_3) h_1(q_1, q_2, q_3) h_3(q_1, q_2, q_3)] + \\ &\quad \left. + \partial_3 [A_3(q_1, q_2, q_3) h_1(q_1, q_2, q_3) h_2(q_1, q_2, q_3)] \right] \end{aligned} \quad (1.70)$$

### Cylindrical coordinates

For cylindrical coordinates eq. (1.70) gives

$$\operatorname{div} \mathbf{A} = \frac{1}{r} \left[ \partial_r [r A_r] + \partial_\theta [A_\theta] + \partial_z [r A_z] \right] \quad (1.71)$$

where  $\mathbf{A}$  and its components are functions of  $(r, \theta, z)$ . Taking out some variables from the partial derivative we get

$$\operatorname{div} \mathbf{A}(r, \theta, z) = \frac{1}{r} \partial_r [r A_r(r, \theta, z)] + \frac{1}{r} \partial_\theta [A_\theta(r, \theta, z)] + \partial_z A_z(r, \theta, z) \quad (1.72)$$

### Spherical coordinates

For spherical coordinates eq. (1.70) gives

$$\operatorname{div} \mathbf{A} = \frac{1}{r^2 \sin \theta} \left[ \partial_r [r^2 \sin \theta A_r] + \partial_\theta [r \sin \theta A_\theta] + \partial_\varphi [r A_\varphi] \right] \quad (1.73)$$

where  $\mathbf{A}$  and its components are functions of  $(r, \theta, \varphi)$ . Taking out some variables from the partial derivative we get

$$\operatorname{div} \mathbf{A}(r, \theta, \varphi) = \frac{1}{r^2} \partial_r [r^2 A_r(r, \theta, \varphi)] + \frac{1}{r \sin \theta} \partial_\theta [\sin \theta A_\theta(r, \theta, \varphi)] + \frac{1}{r \sin \theta} \partial_\varphi A_\varphi(r, \theta, \varphi) \quad (1.74)$$

#### 1.4.5 Dimensional analysis

You can easily check that the divergence has the expected dimensionality of

$$[A][L]^{-1} \quad (1.75)$$



## 1.5 The Laplacian

Since the gradient, when applied on a function, generates a vector, we may apply the divergence operator on the gradient. In such a way, we will define a new operator, the Laplacian

$$\operatorname{div}(\nabla\phi) \equiv \nabla^2\phi \quad (1.76)$$

Since both the gradient and the divergence have been defined in a geometrical, coordinate independent way, also the Laplacian will have a geometrical, coordinate independent definition. Before investigating it, let us obtain its expression in an arbitrary (orthogonal) coordinate system From (eq. 1.40)

$$(\nabla\phi) = \sum_i \frac{1}{h_i} \partial_i \phi \quad (1.77)$$

and (eq. 1.69)

$$\operatorname{div}\mathbf{A} = \frac{\sum_i \partial_i f_i}{\prod_i h_i} \quad (1.78)$$

where the  $f_i$  were defined in eq. (1.68), we obtain

$$\nabla^2\phi = \frac{\sum_i \partial_i \left[ \frac{\prod_{j \neq i} h_j}{h_i} \partial_i \phi \right]}{\prod_i h_i} \quad (1.79)$$

The above is a short hand for

$$\nabla^2\phi = \frac{1}{h_1 h_2 h_3} \left[ \partial_1 \left[ \frac{h_2 h_3}{h_1} \partial_1 \phi \right] + \partial_2 \left[ \frac{h_1 h_3}{h_2} \partial_2 \phi \right] + \partial_3 \left[ \frac{h_1 h_2}{h_3} \partial_3 \phi \right] \right] \quad (1.80)$$

where again we should remember that the  $h$  and  $\phi$  are functions of the  $q_i$ .

### 1.5.1 Cartesian coordinates

In Cartesian coordinates all the  $h_i$  equal to 1 and we have

$$\nabla^2\phi(x, y, z) = \partial_x^2\phi(x, y, z) + \partial_y^2\phi(x, y, z) + \partial_z^2\phi(x, y, z) \quad (1.81)$$

### 1.5.2 Cylindrical coordinates

In cylindrical coordinates, substituting the values of the  $h_i$  and bringing out of derivatives variables when possible, we get

$$\nabla^2\phi(r, \theta, z) = \frac{1}{r} \partial_r (r \partial_r \phi(r, \theta, z)) + \frac{1}{r^2} \partial_\theta^2 \phi(r, \theta, z) + \partial_z^2 \phi(r, \theta, z) \quad (1.82)$$

## 1.5 The Laplacian

---

### 1.5.3 Spherical coordinates

In spherical coordinates, substituting again the values of the  $h_i$  and bringing out of derivatives variables when possible, we get

$$\nabla^2\phi(r, \theta, \varphi) = \frac{1}{r^2}\partial_r(r^2\partial_r\phi(r, \theta, \varphi)) + \frac{1}{r^2\sin\theta}\partial_\theta(\sin\theta\partial_\theta\phi(r, \theta, \varphi)) + \frac{1}{r^2\sin^2\theta}\partial_\varphi^2\phi(r, \theta, \varphi) \quad (1.83)$$

### 1.5.4 Dimensional analysis

You can easily check that the divergence has the expected dimensionality of

$$[\phi][L]^{-2} \quad (1.84)$$

### 1.5.5 Geometrical meaning

In section 1.1 we suggested the form

$$\partial_t\rho(t, \mathbf{x}) = \kappa(\bar{\rho}(t, \mathbf{x}) - \rho(t, \mathbf{x})) \quad (1.85)$$

for a diffusion equation, where  $\bar{\rho}$  stands for an average over a small volume centred in  $\mathbf{x}$ . Since then we got a good experience on small volume integrations and Taylor expansions. Let us try to compute  $\bar{\phi}$  for a cube of size  $a$  centred in  $\mathbf{x}_0 = (x_0, y_0, z_0)$ . Using Cartesian coordinates we have

$$\begin{aligned} \phi(x, y, z) &= \phi(\mathbf{x}_0) + \partial_x\phi(\mathbf{x}_0)(x - x_0) + \partial_y\phi(\mathbf{x}_0)(y - y_0) + \partial_z\phi(\mathbf{x}_0)(z - z_0) + \\ &\partial_x\partial_y\phi(\mathbf{x}_0)(x - x_0)(y - y_0) + \partial_x\partial_z\phi(\mathbf{x}_0)(x - x_0)(z - z_0) + \partial_y\partial_z\phi(\mathbf{x}_0)(y - y_0)(z - z_0) + \\ &\frac{1}{2}\partial_x^2\phi(\mathbf{x}_0)(x - x_0)^2 + \frac{1}{2}\partial_y^2\phi(\mathbf{x}_0)(y - y_0)^2 + \frac{1}{2}\partial_z^2\phi(\mathbf{x}_0)(z - z_0)^2 \\ &+ \text{cubic terms} \end{aligned} \quad (1.86)$$

Now we write, remembering that the volume of the cube is  $a^3$ , and calling  $\Delta x \equiv x - x_0$ ,  $\Delta y \equiv y - y_0$ ,  $\Delta z \equiv z - z_0$ ,

$$\begin{aligned} \bar{\phi} &= \frac{1}{a^3} \int_{-\frac{a}{2}}^{\frac{a}{2}} dx \int_{-\frac{a}{2}}^{\frac{a}{2}} dy \int_{-\frac{a}{2}}^{\frac{a}{2}} \phi(x, y, z) dz = \\ &= \phi(x_0, y_0, z_0) + \frac{1}{a^3} \left( \int \int \int \text{linear terms in } \Delta x, \Delta y, \Delta z d(\Delta x)d(\Delta y)d(\Delta z) + \right. \\ &\int \int \int \text{terms of the form } (\Delta x\Delta y), (\Delta y\Delta z), (\Delta x\Delta z) d(\Delta x)d(\Delta y)d(\Delta z) + \\ &\left. \frac{1}{2} \int \int \int (\partial_x^2\phi(\mathbf{x}_0)\Delta x^2 + \partial_y^2\phi(\mathbf{x}_0)\Delta y^2 + \partial_z^2\phi(\mathbf{x}_0)\Delta z^2) d(\Delta x)d(\Delta y)d(\Delta z) \right) \end{aligned} \quad (1.87)$$

From the trivial result

$$\int_{-\frac{a}{2}}^{\frac{a}{2}} q dq = 0 \quad (1.88)$$

(and Fubini's theorem, namely the possibility of integrating first in one variable, etc.) we obtain that integrals involving the linear terms and terms such as  $xy$  give zero. We are left with integrals in the form (thanks again to Mr. Fubini, since the integral in the two variables unrelated to the integrand gives the face area  $a^2$ )

$$\frac{a^2}{2} \int_{-\frac{a}{2}}^{\frac{a}{2}} \partial_q^2 \phi(\mathbf{x}_0) q^2 dq = \frac{a^2}{2} \partial_q^2 \phi(\mathbf{x}_0) \left[ \frac{q^3}{3} \right]_{-\frac{a}{2}}^{\frac{a}{2}} = \frac{a^5}{24} \quad (1.89)$$

Substituting in eq. 1.87 we get, at the lowest order in  $a$ ,

$$\bar{\phi}(\mathbf{x}_0) - \phi(\mathbf{x}_0) = \frac{a^2}{24} \sum_i \partial_i^2 \phi(\mathbf{x}) = \frac{a^2}{24} \nabla^2 \phi(\mathbf{x}) \quad (1.90)$$

Thanks to our geometrical definitions, this will be true in any coordinate system, and the diffusion equation may be written as

$$\partial_t \rho = \gamma \nabla^2 \rho \quad (1.91)$$

A simple dimensional analysis shows now that

$$[\gamma] = [L]^2 [T] \quad (1.92)$$

## 1.6 Another approach to diffusion

Let us consider a fixed volume  $V$ . The amount of our “substance” in  $V$  is given by the integral of  $\rho$  over the volume<sup>7</sup>.

$$Q_V(t) = \int_V \rho(t, \mathbf{x}) d\mathbf{x} \quad (1.93)$$

$Q_V$  is a function of the only time. Its time derivative is

$$\frac{dQ_V(t)}{dt} = \int_V \partial_t \rho(t, \mathbf{x}) d\mathbf{x} \quad (1.94)$$

If we assume that the substance is “locally conserved” meaning that it cannot disappear but only move to a different place, then the change in time of

---

<sup>7</sup>For example,  $\rho$  may be the electric charge density. The dimensional analysis tells us then that if the dimension of charge is  $[Q]$ , the dimension of  $\rho$  has to be  $[Q][L]^{-3}$ , so that the integral in eq. (1.93) has the proper dimension.  $\rho$  gives a charge only after integration.

## 1.6 Another approach to diffusion

---

$Q_V$  has to be given by the flux of it through the surface of  $V$ . Imagine now that  $\rho$  gives the density of a liquid. This liquid moves in a cylinder (pipe) of section  $A$  with velocity  $v$ . This means that in time  $T$  an amount  $\rho v T A$  of fluid will exit the pipe (Fig. 1.4). Once we divide by time, the “change of amount of water in time” is  $\rho v A$ . If now the “mouth of the pipe” is not orthogonal to the flow of water, although the surface increases, the flow of water does not change. We can thus see that the flow is not given by the product of the velocity by the area, but by  $A \rho \mathbf{v} \cdot \mathbf{n}$ , where  $\mathbf{n}$  is the outgoing unit vector orthogonal to the surface  $A$ . The product  $A \mathbf{v} \cdot \mathbf{n}$  gives us, indeed, the projection of the area in the direction of the water flow and leads us back to the same flow obtained for the orthogonal mouth (refer also to Fig. 1.4). In the discussion above we considered uniform velocities and flat surfaces, but we can generalise by taking an integral over “many infinitesimal flat surfaces and uniform velocities” (the usual Riemann definition of integral).




Figure 1.4: The amount of water that will exit the pipe with the oblique red mouth in time  $T$  is given by the blue parallelogram. The height of the parallelogram is  $h = L \cos \theta$ , where  $\cos \theta = (\mathbf{v} \cdot \mathbf{n})/v$ .

We define a flux vector

$$\Phi(t, \mathbf{x}) = \rho(t, \mathbf{x}) \mathbf{v}(t, \mathbf{x}) \quad (1.95)$$

and write

$$\frac{dQ_V(t)}{dt} = - \int_{S_V} \Phi(t, \mathbf{x}) \cdot \mathbf{n} dS \quad (1.96)$$

The sign is negative because a positive flux in the direction of  $\mathbf{n}$  causes  $Q$  to decrease. Now we use Gauss theorem (eq. 1.59) to obtain

$$\frac{dQ_V(t)}{dt} = - \int_V \operatorname{div} \Phi(t, \mathbf{x}) d\mathbf{x} \quad (1.97)$$

so that, using eq. (1.93)

$$\int_V \partial_t \rho(t, \mathbf{x}) d\mathbf{x} = - \int_V \operatorname{div} \Phi(t, \mathbf{x}) d\mathbf{x} \Rightarrow \int_V (\partial_t \rho(t, \mathbf{x}) + \operatorname{div} \Phi(t, \mathbf{x})) d\mathbf{x} = 0 \quad (1.98)$$

This has to be true regardless of the volume  $V$ , so that we need to have<sup>8</sup>.

$$\partial_t \rho(t, \mathbf{x}) + \operatorname{div} \Phi(t, \mathbf{x}) = 0 \quad (1.99)$$

From eq. (1.32) we know that the gradient gives us the direction in which we have the maximum growth of the function<sup>9</sup>. We previously suggested that  $\rho$  locally grows if its integral in a small surrounding volume is higher than the local value. We may also think that “ $\rho$  flows where  $\rho$  is minimum, or

$$\Phi = -\gamma \nabla \rho \quad (1.100)$$

Here the dimensional analysis tell us that

$$[\rho][L][T]^{-1} = [\gamma][L]^{-1}[\rho] \Rightarrow [\gamma] = [L]^2[T]^{-1} \quad (1.101)$$

Substituting eq. (1.100) in eq.(1.99) we obtain the same form of the diffusion equation

$$\partial_t \rho = \gamma \nabla^2 \rho \quad (1.102)$$

This equation is often called heat equation since  $\rho$  may stand for termic energy (temperature).

---

<sup>8</sup>Otherwise, we could integrate in a small volume around the area where the integrand is not zero, and obtain a non zero integral (these kind of arguments are not completely rigorous due to the possibility of having the integrand different from zero an a set of zero measure

<sup>9</sup>Imagine to use vectors of fixed length  $\Delta s = 1$  in eq. (1.32). The maximum growth will be given for a  $\Delta \mathbf{s}$  in the same direction of the gradient.

## 1.6 Another approach to diffusion

---

## Part II

# Separation of variables





## Chapter 2

# Separation of variables in the heat equation

### 2.1 Boundary conditions

Let us consider the diffusion equation in Cartesian coordinates

$$\partial_t \rho(t, x, y, z) = \gamma \left[ \partial_x^2 \rho(t, x, y, z) + \partial_y^2 \rho(t, x, y, z) + \partial_z^2 \rho(t, x, y, z) \right] \quad (2.1)$$

This equation express a relation between partial derivatives. A function  $\rho$  that represents a solution of the (modelled) physical problem needs to satisfy this relation in every point of a given area of space and time<sup>1</sup>.

We may for example consider the time evolution of the temperature on a metal bar. The bar has a finite length  $L$ . The initial temperature ( $t = 0$ ) is known for each  $x \in [0, L]$ <sup>2</sup>. We also need to provide, in order to obtain a unique solution for the temperature at  $t > 0$ , information about “what

---

<sup>1</sup>Modern physics speaks about space-time. This is due to the fact that in Relativistic physics, time plays a role similar (although not exactly equal) to the other coordinates. As we may “mix” the spatial coordinates  $x, y, z$  by performing a rotation, according to Relativity we may also “mix” time and space, just by moving at constant velocity. We do not realise that this mixing happens because the effect is related to the ratio  $v/c$  where  $v$  is the velocity of movement and  $c$  the velocity of light, and  $v \ll c$  for common life velocities. The physics of Newton is obtained indeed from the physics of Einstein in the  $c \rightarrow \infty$  limit. Eq. (2.1) deals with time and space derivatives in a different way, and thus it is not a relativistic equation. Indeed, according to eq. (2.1), diffusion may happen at infinite velocity. The wave equation of electromagnetism (light propagation),  $[\partial_t^2 - c(\partial_x^2 + \partial_y^2 + \partial_z^2)]\mathbf{A} = 0$ , is a relativistic equation (the minus generates the difference between, time and space in Relativity, while a dimensional analysis easily shows that  $c$  is a velocity). It was indeed reasoning about light that Lorentz, Poincaré and Einstein developed the (Special, i.e. not including Gravitation) Relativity theory.

<sup>2</sup>We approximate the problem as a 1D one, i.e. we consider the bar “infinitely thin”.

## 2.1 Boundary conditions

---

happens at the bars ends",  $x = 0, x = L$ . This information will be needed and will be enough to solve the problem. Why?

Let us remember the situation that we face when we want to solve an *ordinary* differential equation. For a first order differential equation, such as the equation for radioactive decay

$$\frac{dN(t)}{dt} = -\gamma N(t) \quad (2.2)$$

we have a general solution

$$N(t) = Ae^{-\gamma t} \quad (2.3)$$

The solution for a specific problem is obtained by noticing that  $N(0) = A$ . So if initially we have  $N_0$  atoms of the radioactive material, the solution is

$$N(t) = N_0 e^{-\gamma t} \quad (2.4)$$

If we have a second order equation such as the Harmonic oscillator one

$$\frac{d^2 X(t)}{dt^2} = -\omega^2 X(t) \quad (2.5)$$

the general solution is

$$X(t) = A \sin \omega t + B \cos \omega t \quad (2.6)$$

We have now two constants, so we need two conditions to specify our solution. The usual way of doing it is to notice that<sup>3</sup>

$$X(0) = B, \quad \frac{dX(0)}{dt} = \omega A \quad (2.7)$$

so that, calling  $X(0) = X_0$  and  $\frac{dX(0)}{dt} = V_0$ , the solution for a specific problem is

$$X(t) = \frac{V_0}{\omega} \sin \omega t + X_0 \cos \omega t \quad (2.8)$$

But we can solve the oscillator problem in another way, for example by asking to have  $X(0) = X_0$  and  $X(T) = X_T$  so that

$$X_T = A \sin \omega T + X_0 \cos \omega T \Rightarrow A = \frac{X_T - X_0 \cos \omega T}{\sin \omega T} \quad (2.9)$$

provided that  $\omega T \neq n\pi$ . In the latter case, we have to check the equation *before* dividing by the sine, and we get

$$X_T = X_0 \quad (2.10)$$

---

<sup>3</sup>You should perform a dimensional analysis for each equation you see. They are always useful in checking the validity of your results.

so that no other condition may be satisfied.

We have thus seen that for a second order ordinary differential equation we can get a solution either by specifying the initial value of the function and of its derivative<sup>4</sup>, or by specifying its value at two different times. Nevertheless, while the first method always leads to a solution for our problem<sup>5</sup>, as we have seen this is not true for the second approach. The second approach may nevertheless be useful to restrict the possible values that *the model's parameter may assume*. For example, let us ask  $X_0 = 0$  and  $X_{T=1} = 0$ . If we want our solution to be different from zero everywhere we will need (left of eq. 2.9)  $\omega = n\pi$ . We will use this method very often in the following.

Before going back to partial differential equations, let us notice that from the general solution eq. (2.6) we may obtain the expression for the derivative

$$\frac{dX(t)}{dt} = \omega(A \cos \omega t - B \sin \omega t) \quad (2.11)$$

We may thus specify the state of our system also by asking  $\frac{dX(0)}{dt} = V_0$ ,  $\frac{dX(T)}{dt} = V_T$ , i.e. by fixing the derivative (velocity) at two different points.

Let us analyse Fig. 2.1. We want to obtain the solution in the interior of the red area, where  $\rho$  will satisfy eq. (2.1). It looks reasonable that we may need to specify what happens on the *boundary* of the region (*boundary conditions* or BC). We need to know the initial state of the system

$$\rho(0, x) \quad (2.12)$$

but also what happens, for each  $t$ , at the borders. By studying the temperature of the bar, we are assuming that the bar itself follows the heat equation, but what happens around it may be changed in different ways. For example, we could put the bar in contact with one or two different external bodies at  $x = 0$  and  $x = L$ . The temperature of these bodies could be fixed (for example, the bodies being much larger than the bar) or change in time (because connected to some thermic engine, or just because environmental temperature changes with time). Or we could insulate the ends in such a way that heat does not flow (or, in case the equation describes the diffusion of a gas: at the ends we could have walls that do not let the particles pass).

---

<sup>4</sup>For a physical particle, initial position and velocity. Since Newton's equation are second order, this means that the state of a system in classical physics is given by specifying the initial value of all particles and velocities (or, in a more general coordinate system, momenta).

<sup>5</sup>This is a very informal discussion, in which we assume all functions to be as well-behaved as possible. Refer to any book on differential equations for a more formal and rigorous treatment.

## 2.1 Boundary conditions

---




Figure 2.1: Space and time diagram for the problem of diffusion of heat on a 1D bar.

It is clear that, *for the same behaviour given by eq. (2.1) in the prescribed area*, the solution would be different given different boundary conditions, so that the importance of such conditions should be intuitive from a physical view point. Regarding which kind of boundary conditions are necessary we may, in a completely informal and not rigorous way, use the following argument based on a comparison with ordinary differential equations.

Let us wonder what happens at fixed  $x = \bar{x}$ , i.e. on the green line of Fig. 2.1. Since  $x$  is fixed, we may expect our equation to behave as

$$\frac{d\rho_{\bar{x}}(t)}{dt} = f_{\bar{x}}(t) \quad (2.13)$$

so that we need just an initial condition for it,  $\rho(t = 0, \bar{x})$ . But since we want to solve the problem *for all*  $x \in [0, L]$ , we actually need to provide the function

$$\rho(0, x) = f_{t=0}(x) \quad x \in [0, L] \quad (2.14)$$

But obviously we are not dealing with a first order equation in  $t$ , since also what happens on  $x$  is “dynamical”. If we try to think about what happens at fixed  $t = \bar{t}$ , i.e. on the blue line of Fig. 2.1, we expect an equation like

$$\frac{d^2\rho_{\bar{t}}(x)}{dx^2} = f_{\bar{t}}(x) \quad (2.15)$$

---

## Separation of variables in the heat equation

---

We thus need two conditions at fixed  $t$  to specify the solution, since we are dealing with a second order equation. For example, we can fix the temperature at  $x = 0$  and  $x = L$  for all values of  $t > 0$

$$\rho(0, t) = f_{x=0}(t), \quad \rho(L, t) = f_{x=L}(t), \quad t > 0 \quad (2.16)$$

Or we can fix the flow of temperature on the ends. We have seen (eq. 1.100) that the flow is related to the gradient, i.e., for our 1D problem to the derivative. Fixing the flow amounts thus to provide the derivative

$$\frac{d\rho(0, t)}{dt} = g_{x=0}(t), \quad \frac{d\rho(L, t)}{dt} = g_{x=L}(t), \quad t > 0 \quad (2.17)$$

a choice motivated both by our understanding of the physical problem, and by our discussion on the conditions needed to specify second order problems. Insulating the bar, for example, would correspond to fix

$$g_{x=0}(t) = g_{x=L}(t) = 0 \quad \forall t > 0 \quad (2.18)$$

## 2.2 1D heat equation with fixed temperature ends: dimensional analysis

We are almost ready to solve our first PDE problem, but before doing it let us define it properly. We want to study the diffusion of temperature on a 1D bar<sup>6</sup>, while keeping the ends at a fixed (zero) temperature<sup>7</sup>.

The problem is specified by the equation

$$\partial_{t'}\rho(t', x') = \gamma\partial_{x'}^2\rho(t', x') \quad (2.19)$$

initial condition

$$\rho(0, x') = f_{t=0}(x') \quad x' \in [0, L] \quad (2.20)$$

and boundary conditions

$$\rho(0, t') = \rho(L, t') = 0, \quad t' > 0 \quad (2.21)$$

As a first step, we start by simplifying our problem through an appropriate change of variables (this is the reason we called the above variables  $x'$  and  $t'$ ... since we want to get rid of them as soon as possible!). As it will be

---

<sup>6</sup>These 1D problems are obviously idealisations, but they are very useful in understanding mathematical techniques in simple settings.

<sup>7</sup>Or at a given temperature  $\bar{T}$ . In such a case, we can re-define our temperature scale so that  $\rho = T - \bar{T}$ .

## 2.2 1D heat equation with fixed temperature ends: dimensional analysis

---

clear soon, in order to fix the the value of the function to zero at the ends, it will be handy to work on the interval  $[0, \pi]$ . Let us thus define

$$x = \frac{\pi}{L}x' \quad \Rightarrow \quad x' = \frac{L}{\pi} \quad (2.22)$$

so that  $x$  will assume values between 0 and  $L$ . Since we changed variables, also the form of the equation will change. Defining  $\alpha = \pi/L$  we have

$$\frac{\partial}{\partial x'} = \frac{\partial}{\partial x} \frac{\partial x'}{\partial x} = \alpha \frac{\partial}{\partial x} \quad \Rightarrow \quad \partial_{x'}^2 = \alpha \partial_x (\alpha \partial_x) = \alpha^2 \partial_x^2 \quad (2.23)$$

and the equation becomes

$$\partial_{t'} \rho(t', x) = \gamma \alpha^2 \partial_x^2 \rho(t, x) \quad (2.24)$$

Now we notice that if we if we define

$$t = \beta t' \quad \Rightarrow \quad \partial_{t'} = \beta \partial_t \quad (2.25)$$

the equation becomes

$$\partial_t \rho(t, x) = \frac{\gamma \alpha^2}{\beta} \partial_x^2 \rho(t, x) \quad (2.26)$$

The trick is now to define  $\beta = \gamma \alpha^2$  so that the change in the time variable is

$$t = \gamma \frac{\pi^2}{L^2} t' \quad \Rightarrow \quad t' = \frac{L^2}{\pi^2 \gamma} \quad (2.27)$$

and the equation becomes

$$\partial_t \rho(t, x) = \partial_x^2 \rho(t, x) \quad (2.28)$$

We may call this the a-dimensional heat equation since, from eqs. (1.101,2.22,2.27),

$$[x] = \frac{[L]}{[L]} = [*]^0 \quad [t] = [\gamma] \frac{[*]^0}{[L]^2} [T] = [*]^0 \quad (2.29)$$

where with the the notation  $[*]^0$  we just mean a pure number, independent on unit systems.

The form of the equation got simplified. We will now proceed to find a solution for  $\rho(t, x)$  from which we will be able to obtain the solution in the original dimensional variable through the transformation rules eqs. (2.22,2.27). Obviously, the same rules may also be used to express the boundary conditions eqs. (2.20,2.21) as functions of the a-dimensional variables  $(t, x)$ .

## 2.3 Separation of variables for the 1D heat equation

The *separation of variables* method is based on assuming that the solution may be written as the product of a function of  $t$  and  $x$

$$\rho(t, x) = T(t)X(x) \quad (2.30)$$

The method is basically based on trusting that the method will work. We will follow the consequence of the *ansatz*<sup>8</sup> eq. (2.30), perform a few doubtful steps like dividing by some function without worrying if it is zero or not, and finally arrive at a general solution for our equation. Only after we will reach the solution we will discuss about when this method may work or not. Let us substitute<sup>9</sup> eq. (2.30) in eq. (2.28)

$$\partial_t [T(t)X(x)] = \partial_x^2 [T(t)X(x)] \quad \Rightarrow \quad X(x)\partial_t [T(t)] = T(t)\partial_x^2 [X(x)] \quad (2.31)$$

Let us now divide both sides by  $T(t)X(x)$  to obtain

$$\frac{\partial_t T(t)}{T(t)} = \frac{\partial_x^2 X(x)}{X(x)} \quad (2.32)$$

On the left we have a function of  $t$ , and on the right a function of  $x$ , and they are equal everywhere, i.e. *for each arbitrary*  $x \in [0, L]$  and  $t \geq 0$ . From this follows that both sides of the equation have to be constants<sup>10</sup>,

$$\frac{\partial_t T(t)}{T(t)} = C = \frac{\partial_x^2 X(x)}{X(x)} \quad (2.33)$$

and

$$\partial_t T(t) = CT(t) \quad (2.34)$$

$$\partial_x^2 X(x) = CX(x) \quad (2.35)$$

We reduced our problem to two independent ordinary differential equations!

We note that if eqs. (2.34,2.35) are true for all  $t, x$  we have

$$\partial_t \rho = \partial_t (XT) = X\partial_t T = CXT = T\partial_x^2 X = \partial_x^2 (TX) = \partial_x^2 \rho \quad (2.36)$$

---

<sup>8</sup>A German word for an educated guess.

<sup>9</sup>We will use the partial derivative symbol also when it acts on a function of a single variable.

<sup>10</sup>Imagine we have  $f(y) = g(z)$  for two independent variables  $y$  and  $z$ . Assume we have  $f(y_1) = a \neq b = f(y_2)$ , i.e.  $f$  is not constant. Now, if we pick up a value  $z_1$ , we want to have  $g(z_1) = f(y_1) = a$ , since the equality has to hold for arbitrarily picked values of  $y$  and  $z$ . But then  $g(z_1) = a \neq b = f(y_2)$  and a contradiction arises.

### 2.3 Separation of variables for the 1D heat equation

---

so that we actually do not need to worry about divisions by zero.

Eq. (2.34) is trivially solved by

$$T(t) = A'e^{Ct} \quad (2.37)$$

We proceed now to study eq. (2.35). We know that the behaviour of this equation depends on the sign of  $C$ . Let us consider the three possible cases

(I)  $C = 0$

We have

$$\partial_x^2 X(x) = 0 \Rightarrow \partial_x X(x) = \bar{A} \Rightarrow X(x) = \bar{A}x + \bar{B} \quad (2.38)$$

so that

$$\rho(t, x) = A'e^0(\bar{A}x + \bar{B}) = Ax + B \quad (2.39)$$

where we defined  $A = A'\bar{A}$ ,  $B = A'\bar{B}$ . If we want eq. (2.21) to hold, we need to have

$$\rho(t, 0) = B = 0, \quad \rho(t, \pi) = A\pi + 0 = 0 \quad \Rightarrow A = 0 \quad (2.40)$$

The  $C = 0$  case led us to the solution  $\rho = 0$ . This is trivially a solution of the heat equation with the prescribed boundary conditions, but it can satisfy eq. (2.20) only if  $f_{t=0}(x) = 0 \forall x \in [0, \pi]$ .

(II)  $C > 0$

Since  $C$  is positive, we re-name it as  $C = \lambda^2$ . Now we have from the theory of linear differential equations

$$\partial_x^2 X(x) = \lambda^2 X(x) \Rightarrow X(x) = \bar{A}e^{\lambda x} + \bar{B}e^{-\lambda x} \quad (2.41)$$

and

$$\rho(t, x) = A'e^{\lambda^2 t} X(x) = e^{\lambda^2 t} (Ae^{\lambda x} + Be^{-\lambda x}) \quad (2.42)$$

thanks to usual re-definition of the constants. The term  $e^{\lambda^2 t}$  is obviously positive, so the conditions eq. (2.21) are equivalent to ask for the term in the parenthesis to be null (we call this term  $X$  by absorbing  $A'$  in  $X$ )

$$0 = X(0) = Ae^{\lambda 0} + Be^{-\lambda 0} \Rightarrow B = -A \quad (2.43)$$

and

$$0 = X(\pi) = A(e^{\lambda \pi} + Be^{-\lambda \pi}) \Rightarrow A = 0 \quad (2.44)$$

where the last result obviously follows since for  $\lambda > 0$  we have  $\lambda \pi \neq -\lambda \pi$  and the exponential is strictly growing. The only possible solution is again  $\rho = 0$ , that would nevertheless lead to a contradiction since we had asked  $C > 0$ . No solution is thus found for  $C > 0$ .



(III)  $C < 0$

Since  $C$  is negative, we re-name it as  $C = -\lambda^2$ . Now we get

$$\partial_x^2 X(x) = -\lambda^2 X(x) \Rightarrow X(x) = \bar{A} \sin(\lambda x) + \bar{B} \cos(\lambda x) \quad (2.45)$$

and

$$\rho(t, x) = A' e^{-\lambda^2 t} X(x) = e^{-\lambda^2 t} (A \sin(\lambda x) + B \cos(\lambda x)) \quad (2.46)$$

Again, the time dependent factor is positive, so in order for eq. (2.21) to be true we need to ask  $X = 0$  (and anyway, eq. (2.21) need to be true  $\forall t > 0$ , and thus is determined by X).

$$X(0) = B = 0 \Rightarrow X(x) = A \sin(\lambda x) \quad (2.47)$$

and

$$X(\pi) = A \sin(\lambda \pi) = 0 \quad (2.48)$$

If we want to avoid to get again a null solution, we have to ask  $\lambda \in \mathbb{Z}$  so that  $\sin(\lambda \pi) = 0$  (we now see the reason of our change of variables in  $x$ ). We note anyway that  $\lambda = 0$  has to be avoided because it will lead to the  $C = 0$  case, while if  $\lambda = -n < 0$  we have  $(-n)^2 = (n^2)$  and

$$X(x) = A \sin(-nx) = -A \sin(nx) = A'' \sin(nx) \quad (2.49)$$

so that negative integers just produce the same solutions given by positive ones. We may thus limit ourselves to

$$\rho(t, x) = A e^{-n^2 t} \sin(nx), \quad n \in \mathbb{N} \quad (2.50)$$

The  $C < 0$  case just led us to solutions for our partial differential equation, eq. (2.50). This solution may also be re-written using the original variables  $(t', x')$  as<sup>11</sup>

$$\rho(t', x') = A e^{-\frac{n^2 \pi^2 \gamma t'}{L^2}} \sin\left(\frac{n\pi}{L} x'\right), \quad n \in \mathbb{N} \quad (2.51)$$

Nevertheless, eq. (2.50) gives, for  $t = 0$ ,

$$\rho(0, x) = A \sin(nx) \quad (2.52)$$

and thus cannot satisfy eq. (2.20) unless  $f_{t=0}$  is proportional to a sine function. This is better than a null function, but is not general at all!

---

<sup>11</sup>You may check that the arguments in the sine and exponential of eq. (2.51) have no dimension. This is correct and necessary (such functions may be written as power series of their arguments and thus it would make no sense to have arguments with dimensions).

## 2.3 Separation of variables for the 1D heat equation

---

### 2.3.1 Separation of variables and linearity

Nevertheless, we can use solutions in the form eq. (2.50) to build new solutions. *This is due to the fact that the heat equation is linear, and the problem that we have specified is homogeneous both in the equation and in the boundary condition.* Let us see what this means.

#### Linearity

Let us assume we have an equation such as

$$D[f(\mathbf{x})] = 0 \quad (2.53)$$

Here  $f$  is a function from  $\mathbb{R}^n \rightarrow \mathbb{R}$  and  $D$  is an operator, i.e. a function that operates on a function to produce another function (e.g. a derivative). We say that  $D$  is linear if given two functions  $f$  and  $g$  and numbers  $\alpha, \beta$ , the following holds

$$D[\alpha f(\mathbf{x}) + \beta g(\mathbf{x})] = \alpha D[f(\mathbf{x})] + \beta D[g(\mathbf{x})] \quad (2.54)$$

An example of a linear operator is the multiplication by a function  $h(\mathbf{x})$

$$\begin{aligned} D[\alpha f(\mathbf{x}) + \beta g(\mathbf{x})] &= h(\mathbf{x})[\alpha f(\mathbf{x}) + \beta g(\mathbf{x})] = \alpha h(\mathbf{x})f(\mathbf{x}) + \beta h(\mathbf{x})g(\mathbf{x}) \\ &= \alpha D[f(\mathbf{x})] + \beta D[g(\mathbf{x})] \end{aligned} \quad (2.55)$$

A partial derivative, or a second partial derivative, is obviously also linear<sup>12</sup>, and operators that include sums of derivatives of different orders and multiplications by functions are also linear, as can be easily checked.

Obviously, not all operators are linear.  $D[f] = f^2$  is clearly not linear, since  $D[\alpha f] = \alpha^2 D[f]$ . Also *adding* a function  $D[f] = f + h$  is not linear, since  $D[f + g] = f + g + h \neq D[f] + D[g] = g + f + 2h$ .

Nevertheless, our differential equation (2.28) is in the form eq. (2.53) with the *linear* operator<sup>13</sup>

$$D = \partial_t - \partial_x^2 \quad (2.56)$$

As a result, if we have two solutions for our problem,  $D[f] = 0$  and  $D[g] = 0$ , we can get a third solution just by a combination of them,  $D[\alpha f + \beta g] = 0$ .

---

<sup>12</sup>Linearity is a property of derivation.

<sup>13</sup>In section 1.6 we assumed  $\rho$  to be conserved, namely not created or destroyed. If we quit this assumption we get a diffusion equation with a source, in the form  $D'[f] = 0$ ,  $D'[f] = D[f] - h$ , and we lose linearity. Equations with such terms are called *not homogeneous*.

### Homogeneous boundary conditions

Until now we did not touch the problem of the boundary conditions. We have seen that if we have two solutions  $\rho_1$  and  $\rho_2$  that satisfy eq. (2.28), then also

$$\rho_3(t, x) = \alpha\rho_1(t, x) + \beta\rho_2(t, x) \quad (2.57)$$

will satisfy eq. (2.28). But our problem includes also boundary conditions. If such conditions were, for example, in the form

$$\rho(t, 0) = f_{x=0} \neq 0 \quad (2.58)$$

and if  $\rho_1$  and  $\rho_2$  satisfied such conditions, we would have

$$\rho_3(0, x) = \alpha\rho_1(0, x) + \beta\rho_2(0, x) = (\alpha + \beta)f_{x=0} \quad (2.59)$$

which is in general *not* equal to  $f_{x=0}$ .

The process of combining two solutions to obtain a third will thus be possible to apply only when we have *homogeneous boundary conditions* such as<sup>14</sup>

$$\rho(t, 0) = \rho(t, L) = 0 \quad (2.60)$$

or

$$\partial_x\rho(t, 0) = \partial_x\rho(t, L) = 0 \quad (2.61)$$

## 2.4 Fourier series and general solution to the 1D heat equation with homogeneous boundary conditions

### 2.4.1 General solution

For our problem we may thus combine different solutions in the form eq. (2.50) to obtain functions that are more general than simple sine functions. But, how general can we get?

At  $t = 0$ , eq. (2.50) becomes

$$\rho(0, x) = A \sin(nx), \quad n \in \mathbb{N} \quad (2.62)$$

so that the most general initial condition that we may express with the method of separation of variables is

$$\rho(0, x) = \sum_{n=1}^{\infty} A_n \sin(nx) \quad (2.63)$$

---

<sup>14</sup>We may always deal with boundary conditions in the form  $\rho(t, 0) = f_{x=0} = c$  by re-defining the zero of our function, i.e. by studying  $\rho' = \rho - c$ . This function may assume also negative values, but it is described by the same equation that describes  $\rho$ .

## 2.5 Computation of coefficients

---

An important theorem in functional analysis says that a function which is enough well behaved<sup>15</sup> and assumes  $f(0) = f(\pi) = 0$  may be approximated to any degree by the *Fourier series* eq. (2.63). We will assume our initial conditions to satisfy these criteria<sup>16</sup>. We thus state that the general solution to our problem is written as

$$\rho(t, x) = \sum_{n=1}^{\infty} A_n e^{-n^2 t} \sin(nx) \quad (2.64)$$

in a-dimensional variables, and

$$\rho(t', x') = \sum_{n=1}^{\infty} A_n e^{-\frac{n^2 \pi^2 \gamma t'}{L^2}} \sin\left(\frac{n\pi}{L} x'\right) \quad (2.65)$$

in the original, dimensional, ones.

## 2.5 Computation of coefficients

The formula (2.64), although theoretically important, is by itself scarcely useful since initial conditions are not given in the form eq. (2.64). We need thus a way to obtain the coefficients  $A_n$ , that completely specify the solution, from the initial condition. The process is based on a formal analogy with the process that we may use to obtain the coefficients of a vector in a given basis through scalar products, and is based on the definition of a vector space of functions and a corresponding scalar product. We will deal with the theory, although in a largely non rigorous way, in a subsequent chapter, but for the moment we will limit ourselves to the introduction of the computation technique.

We first notice that

$$\int_0^{\pi} \sin(nx) \sin(mx) dx = \frac{\pi}{2} \delta_{nm} = \begin{cases} \frac{\pi}{2} & \text{if } n = m \\ 0 & \text{if } n \neq m \end{cases} \quad (2.66)$$

This result is based on the following trigonometric relations<sup>17</sup>

$$\sin(x + y) = \sin(x) \cos(y) + \sin(y) \cos(x) \quad (2.67)$$

$$\cos(x + y) = \cos(x) \cos(y) - \sin(x) \sin(y) \quad (2.68)$$

---

<sup>15</sup>For example, piece-wise continuous and differentiable

<sup>16</sup>Functions in classical physics are considered in general *smooth*, i.e. possible to differentiate infinite times, although sometimes piece-wise continuous functions may be used to model some initial conditions.

<sup>17</sup>A straightforward but tedious analytical proof may be based on Euler formulae.

---

### Separation of variables in the heat equation

---

From these we obtain, by substitution and using  $\cos(-x) = \cos(x)$ ,  $\sin(-x) = -\sin(x)$

$$\cos(x-y) - \cos(x+y) = 2 \sin(x) \sin(y) \Rightarrow \sin(x) \sin(y) = \frac{1}{2} [\cos(x-y) - \cos(x+y)] \quad (2.69)$$

Our integral becomes

$$\int_0^\pi \sin(nx) \sin(mx) dx = \frac{1}{2} \int_0^\pi \cos((n-m)x) dx - \frac{1}{2} \int_0^\pi \cos((n+m)x) dx \quad (2.70)$$

Now, if  $n \neq m$ , we can use

$$\int_0^\pi \cos(kx) dx = \frac{1}{k} [\sin(kx)]_0^\pi = 0 \quad (2.71)$$

valid for integer  $k \neq 0$ , and obtain

$$\int_0^\pi \sin(nx) \sin(mx) dx = 0 \quad (2.72)$$

If  $n = m$ , eq. (2.69) becomes

$$\sin(x)^2 = \frac{1}{2} [1 - \cos(2x)] \quad (2.73)$$

so that

$$\int_0^\pi \sin^2(nx) dx = \frac{1}{2} \int_0^\pi dx - \frac{1}{2} \int_0^\pi \cos(2nx) dx = \frac{1}{2}(\pi + 0) = \frac{\pi}{2} \quad (2.74)$$

We may get rid of the factor  $\pi/2$  by defining the functions

$$s_n(x) = \sqrt{\frac{2}{\pi}} \sin(nx) \quad (2.75)$$

so that

$$\int_0^\pi s_n(x) s_m(x) dx = \delta_{nm} \quad (2.76)$$

Now, just by redefining the  $A_n$ , we may re-write eqs. (2.63,2.64) as

$$\rho(0, x) = \sum_{n=1}^{\infty} A_n s_n(x) \quad (2.77)$$

$$\rho(t, x) = \sum_{n=1}^{\infty} A_n e^{-n^2 t} s_n(x) \quad (2.78)$$

The initial condition for our problem is given by

$$\rho(0, x) = f_{t=0}(x) = \sum_{n=1}^{\infty} A_n s_n(x) \quad (2.79)$$


---

## 2.6 Time dependence

---

We know that it can be written as a Fourier series, but we explicitly have only  $f_{t=0}(x)$ . We may compute the integral

$$\int_0^\pi s_m(x) f_{t=0}(x) dx = \int_0^\pi s_m(x) \sum_{n=1}^{\infty} A_n s_n(x) dx = \sum_{n=1}^{\infty} A_n \int_0^\pi s_m(x) s_n(x) dx = A_m \quad (2.80)$$

The general solution may be thus written as

$$\rho(t, x) = \sum_{n=1}^{\infty} \left[ \int_0^\pi s_n(x) f(x)_{t=0} dx \right] e^{-n^2 t} s_n(x) \quad (2.81)$$

We used the a-dimensional equation in order to develop the theory, but we may now see how to write down a procedure to get the coefficients directly from the dimensional initial condition. Starting from eq. (2.66), and changing variables using eq. (2.22) we get, from  $dx = \alpha dx'$

$$\delta_{nm} = \frac{2}{\pi} \int_0^L \sin(n \alpha x') \sin(m \alpha x') \alpha dx' \quad (2.82)$$

or

$$\int_0^L \sin\left(n \frac{\pi}{L} x'\right) \sin\left(m \frac{\pi}{L} x'\right) dx' = \frac{L}{2} \delta_{nm} \quad (2.83)$$

that has the proper dimension (an integral over distance of a pure number give a length). Defining

$$s_n^L(x') = \sqrt{\frac{2}{L}} \sin\left(\frac{n\pi}{L} x'\right) \quad (2.84)$$

we have

$$\int_0^L s_n^L(x') s_m^L(x') dx' = \delta_{nm} \quad (2.85)$$

and, repeating the discussion above, the general solution in dimensional variables may be written as

$$\rho(t', x') = \sum_{n=1}^{\infty} \left[ \int_0^\pi s_n^L(x') f(x')_{t=0} dx' \right] e^{-\frac{n^2 \pi^2 \gamma t'}{L^2}} s_n^L(x') \quad (2.86)$$

## 2.6 Time dependence

Let us write again the general solution in the form

$$\rho(t, x) = \sum_{n=1}^{\infty} A_n e^{-n^2 t} \sin(nx) \quad (2.87)$$

We have highlighted the time dependence in blue. Since  $n \neq 0$ , all these terms go to zero in the limit  $t \rightarrow \infty$ , the terms with high  $n$  decaying faster than those with low  $n$ . This shows that

$$\lim_{t \rightarrow \infty} \rho(t, x) = 0 \quad \forall x \in [0, \pi] \quad (2.88)$$

and as a result, the identically zero function is the only stationary solution of the problem, where by stationary solution we mean a solution that does not depend on time

$$\rho(t, x) = \rho_{\text{stat}}(x) \quad (2.89)$$

This is not surprising, both from a mathematical and physical viewpoint.

Mathematically, we have

$$\partial_t \rho_{\text{stat}}(x) = 0 = \partial_x^2 \rho_{\text{stat}}(x) \Rightarrow \rho_{\text{stat}}(x) = Ax + B \quad (2.90)$$

but since we have  $\rho = 0$  at 0 and  $\pi$ , we necessarily obtain zero everywhere.

Physically, we may expect the bar to continually lose energy at the ends until the zero solution is reached<sup>18</sup>.

## 2.7 Flux boundary conditions

### 2.7.1 Stationary solution

Let us consider the solution for the stationary (time independent) problem in case the boundary conditions state that we have zero flux at the end, or (eq. 2.61)

$$\partial_x \rho(t, 0) = \partial_x \rho(t, \pi) = 0 \quad (2.91)$$

The argument leading to eq. (2.90) does not depend on boundary conditions, so that the solution is in the form  $\rho_{\text{stat}}(x) = Ax + B$ , with  $\partial_x \rho_{\text{stat}}(x) = A$ . Fixing  $A = 0$  satisfies the flux conditions at *both* ends, so that the solution for the stationary problem is

$$\rho_{\text{stat}} = B \quad (2.92)$$

There is no way to further specify the solution based on boundary conditions. If prepare the bar at constant temperature  $B$ , and insulate the ends, the

---

<sup>18</sup>The same applies obviously also if this is a “re-scaled” zero, for example if our boundary conditions for the heat equation correspond to keeping the ends at the zero of the Celsius scale, instead of the absolute zero. In this case, the bar may gain energy from the ends, since the function may assume negative values, but the validity of the result does not change.

## 2.7 Flux boundary conditions

---

bar will stay at  $B$ . This is obviously very understandable and intuitive also from a physical viewpoint. Nevertheless, in section 2.6 we have seen that the  $\rho = 0$  stationary solution is also the  $t \rightarrow \infty$  limit for *any* general solution of the problem in which the ends were kept at zero. Is also the solution eq. (2.92) a limit for large  $t$ ? And if it is, how does  $B$  relate to the initial condition?

Physical intuition suggests that since the ends are insulated, the system will conserve its energy and reach a stationary, constant solution in which  $B$  equals the average energy of the initial condition, i.e.

$$B = \langle \rho(0, x) \rangle_x = \frac{1}{\pi} \int_0^\pi \rho(0, x) dx \quad (2.93)$$

Nevertheless, we do not need to rely on physical intuition.

### 2.7.2 General solution

We use again the separation of variable method. The discussion leading to eqs. (2.34,2.35) is independent on boundary conditions, so that we can start from

$$\partial_t T(t) = CT(t) \quad (2.94)$$

with solution

$$T(t) = A'e^{Ct} \quad (2.95)$$

and

$$\partial_x^2 X(x) = CX(x) \quad (2.96)$$

We consider again the 3 possible cases for the sign of  $C$

(I)  $C > 0$

Since  $C$  is positive, we re-name it as  $C = \lambda^2$ , and solve for

$$\partial_x^2 X(x) = \lambda^2 X(x) \Rightarrow X(x) = \bar{A}e^{\lambda x} + \bar{B}e^{-\lambda x} \quad (2.97)$$

We now have<sup>19</sup>

$$\partial_x X(x) = \lambda [\bar{A}e^{\lambda x} - \bar{B}e^{-\lambda x}] \quad (2.98)$$

$$\partial_x X(0) = \lambda(\bar{A} - \bar{B}) = 0 \Rightarrow \bar{B} = \bar{A} \quad (2.99)$$

and

$$\partial_x X(\pi) = \bar{A}(e^{\lambda \pi} - e^{-\lambda \pi}) = 0 \quad (2.100)$$

But this latter equation is satisfied only for  $\bar{A} = 0$ , which would lead to  $C = 0$ , i.e., to a contradiction. There is thus no solution corresponding to  $C > 0$ .

---

<sup>19</sup>Obviously, for the conditions on  $\partial_x$  it is sufficient to analyse  $X$ .



(II)  $C = 0$

We have

$$\partial_x^2 X(x) = 0 \Rightarrow \partial_x X(x) = \bar{A} \Rightarrow X(x) = \bar{A}x + \bar{B} \quad (2.101)$$

From

$$\partial_x X(x) = \bar{A} = 0 \quad (2.102)$$

we find, with the usual re-definition of constants, the solution

$$\rho(t, x) = T(t)X(x) = e^{0t}B = B \quad (2.103)$$

which is exactly the stationary solution of section 2.7.1.

(III)  $C < 0$

Since  $C$  is negative, we re-name it as  $C = -\lambda^2$ , and find

$$\partial_x^2 X(x) = -\lambda^2 X(x) \Rightarrow X(x) = \bar{A} \sin(\lambda x) + \bar{B} \cos(\lambda x) \quad (2.104)$$

From

$$\partial_x X(x) = \lambda(\bar{A} \cos(\lambda x) - \bar{B} \sin(\lambda x)) \quad (2.105)$$

we get

$$\partial_x X(0) = \lambda \bar{A} \Rightarrow X(x) = \bar{B} \cos(\lambda x) \quad (2.106)$$

and

$$\partial_x X(\pi) = -\lambda \bar{B} \sin(\lambda \pi) = 0 \quad (2.107)$$

As before, to avoid a null solution, we have to ask  $\lambda \in \mathbb{Z}$  so that  $\sin(\lambda \pi) = 0$ .  $\lambda = 0$  corresponds to the  $C = 0$  case, while if  $\lambda = -n < 0$  we have  $(-n)^2 = (n^2)$  and

$$X(x) = \bar{B} \cos(-nx) = \bar{B} \cos(nx) \quad (2.108)$$

so that negative integers just produce the same solutions given by positive ones. We may thus limit ourselves to

$$\rho(t, x) = B e^{-n^2 t} \cos(nx), \quad n \in \mathbb{N} \quad (2.109)$$

If two solutions  $\rho_1, \rho_2$  satisfy the boundary condition  $\partial_x \rho(0) = \partial_x \rho(\pi) = 0$ , also a linear combination of them will satisfy the same condition since

$$\partial_x(\alpha \rho_1 + \beta \rho_2) = \alpha \partial_x \rho_1 + \beta \partial_x \rho_2 \quad (2.110)$$

We may thus write the general solution as

$$\rho(t, x) = \sum_{n=0}^{\infty} B_n e^{-n^2 t} \cos(nx) = B_0 + \sum_{n=1}^{\infty} B_n e^{-n^2 t} \cos(nx) \quad (2.111)$$

## 2.7 Flux boundary conditions

---

where on the rightmost expression we singled out the time independent term corresponding to  $C = 0$ . This term will obviously be the only one to survive in the  $t \rightarrow 0$  limit. As it happened for the case in which the ends temperature was kept to 0, time decay is faster for high  $n$ , but now we have also a  $t$  independent term.

This solution is, again according to the Fourier series theory, able to approximate any regular initial condition in agreement with eqs. (2.61)

### 2.7.3 Computation of coefficients

We want again to be able to compute the coefficients  $B_n$  from the initial condition  $\rho_{t=0}(x)$ . We use

$$\cos(x-y) + \cos(x+y) = 2 \cos(x) \cos(y) \Rightarrow \cos(x) \cos(y) = \frac{1}{2} \left[ \cos(x-y) + \cos(x+y) \right] \quad (2.112)$$

and

$$\int_0^\pi \cos(kx) dx = \frac{1}{k} \left[ \sin(kx) \right]_0^\pi = 0 \quad (2.113)$$

provided that  $k$  is a non zero integer, so that if  $m \neq n$

$$\int_0^\pi \cos(nx) \cos(mx) dx = \frac{1}{2} \int_0^\pi \cos((n-m)x) dx + \frac{1}{2} \int_0^\pi \cos((n+m)x) dx = 0 \quad (2.114)$$

The formula above applies also to the case  $n = 0$ ,  $\cos(nx) = 1$ .

If  $n = m \neq 0$  we have

$$\int_0^\pi \cos^2(nx) dx = \frac{1}{2} \int_0^\pi dx + \frac{1}{2} \int_0^\pi \cos(2nx) dx = \frac{1}{2}(\pi + 0) = \frac{\pi}{2} \quad (2.115)$$

while for  $n = m = 0$  we have obviously

$$\int_0^\pi dx = \pi \quad (2.116)$$

Let us then define

$$c_0(x) = \frac{1}{\sqrt{\pi}}, \quad c_n(x) = \sqrt{\frac{2}{\pi}} \cos(nx) \quad n \in \mathbb{N} \quad (2.117)$$

so that we have

$$\int_0^\pi c_n(x) c_m(x) dx = \delta_{n,m} \quad (2.118)$$

We re-write the general solution as

$$\rho(t, x) = \sum_{n=0}^{\infty} B_n e^{-n^2 t} c_n(x) \quad (2.119)$$

where (see eq. 2.80)

$$B_n = \int_0^\pi c_n(x) f_{t=0}(x) dx \quad (2.120)$$

If you prefer working with the dimensional variables, you may define

$$c_0^L(x') = \sqrt{\frac{1}{L}}, \quad c_n^L(x') = \sqrt{\frac{2}{L}} \cos\left(\frac{n\pi}{L}x'\right) \quad (2.121)$$

and

$$\rho(t', x') = \sum_{n=0}^{\infty} \left[ \int_0^\pi c_n^L(x') f(x')_{t=0} \right] e^{-\frac{n^2 \pi^2 \gamma t'}{L^2}} c_n^L(x') \quad (2.122)$$

Let us explicitly compute

$$B_0 = \int_0^\pi \frac{1}{\sqrt{\pi}} f_{t=0}(x) dx = \sqrt{\pi} \langle f_{t=0} \rangle_x \quad (2.123)$$

We thus have

$$\lim_{t \rightarrow \infty} \rho(t, x) = B_0 c_0 = \sqrt{\pi} \langle f_{t=0} \rangle_x \frac{1}{\sqrt{\pi}} = \langle f_{t=0} \rangle_x \quad (2.124)$$

confirming our physical intuition.

## 2.8 Diffusion on the circle

### 2.8.1 Periodic boundary conditions




Figure 2.2: Diffusion on the circle

Let us now imagine that, without changing the physical properties of diffusion, we may bend the bar in a circle in such a way so that the two

## 2.8 Diffusion on the circle

---

ends will be unified in a single point (Fig. 2.2). It is useful then to make a change of variables, for this problem, from

$$x' \in [0, L] \rightarrow x \in [0, 2\pi] \quad (2.125)$$

namely

$$x = \alpha x', \quad \alpha = \frac{2\pi}{L} \quad (2.126)$$

and the corresponding (see eq. 2.27)

$$t = \gamma \alpha^2 t' = \frac{4\pi^2 \gamma}{L^2} t' \quad (2.127)$$

that leads to the usual

$$\partial_t \rho = \partial_x^2 \rho \quad (2.128)$$

expression for the differential equation.

The reason for this change of variables should be clear:  $x$  (we could have called it  $\theta$ ) is the angle that gives the position on the circle. Now, since we assume it to be purely 1D, the system is completely closed, since its ends interact with each other, and are actually the same point. The boundary conditions are now fixed in a natural way by asking the solution<sup>20</sup> to be periodic

$$\rho(t, x + 2\pi) = \rho(t, x) \quad (2.129)$$

These are the so called *periodic boundary conditions*.

### 2.8.2 Separation of variables

It is straightforward that if  $\rho_1$  and  $\rho_2$  are periodic then

$$\alpha \rho_1(t, x + 2\pi) + \beta \rho_2(t, x + 2\pi) = \alpha \rho_1(t, x) + \beta \rho_2(t, x) \quad (2.130)$$

so that we will be able to combine different solutions to the problem to obtain new solutions. We proceed again with the separation method, that leads as usual to

$$\partial_t T(t) = CT(t) \quad (2.131)$$

with solution

$$T(t) = A' e^{Ct} \quad (2.132)$$

and

$$\partial_x^2 X(x) = CX(x) \quad (2.133)$$

We consider again the 3 possible cases for the sign of  $C$

---

<sup>20</sup>And by consequence all its derivatives.

---

### Separation of variables in the heat equation

---

(I)  $C = \lambda^2 > 0$

The solution

$$X(x) = \bar{A}e^{\lambda x} + \bar{B}e^{-\lambda x} \quad (2.134)$$

is not periodic, regardless of the values of the constants, so that  $C > 0$  will not satisfy the boundary conditions

(II)  $C = 0$

The solution

$$X(x) = \bar{A}x + \bar{B} \quad (2.135)$$

is periodic only for  $\bar{A} = 0$  so that we get the solution

$$\rho(t, x) = T(t)X(x) = e^{0t}B = B \quad (2.136)$$

We may already understand (and soon prove) that this stationary solution will be the  $\lim_{t \rightarrow 0}$  of  $\rho$ , and its value will correspond to the initial average energy/temperature/density<sup>21</sup>.

(III)  $C = -\lambda^2 < 0$

This time may proceed in a different (but equivalent) way. We know that the general complex solution to

$$\partial_x^2 X(x) = -\lambda^2 X(x) \quad (2.137)$$

is

$$X(x) = A'e^{i\lambda x} + B'e^{-i\lambda x} \quad (2.138)$$

This solution will be periodic if  $\lambda = n \in \mathbb{Z}$ . To ask for the solution to be real we may notice that the complex conjugate  $X^*$  is

$$X^*(x) = (A')^*e^{-inx} + (B')^*e^{inx} \quad (2.139)$$

so that the solution is real if

$$A' = (B')^* \Rightarrow (A')^* = B' \quad (2.140)$$

or

$$A' = B'' + iA'', \quad B' = B'' - iA'' \quad (2.141)$$

with  $A'', B'' \in \mathbb{R}$ , so that

$$X(x) = 2B'' \frac{e^{inx} + e^{-inx}}{2} + 2iA'' \frac{e^{inx} - e^{-inx}}{2i} = \bar{B} \cos(nx) + \bar{A} \sin(nx) \quad (2.142)$$

---

<sup>21</sup>The system is clearly conserving  $\int \rho$  since it has no contact with the exterior world. The boundary from which  $\rho$  could flow, that before corresponded to the two ends, has now been reduced to no point.

## 2.8 Diffusion on the circle

---

### 2.8.3 Complex “orthogonal” solutions

The periodic complex solutions

$$e^{int}, n \in \mathbb{Z} \quad (2.143)$$

are extremely important and may shed some light also on the behaviour of real solutions, as we will see soon. Being of a simple exponential form, they are more easy to operate on than the trigonometric functions.

We start their study by computing the following integral<sup>22</sup>

$$\int_0^{2\pi} (e^{imx})^* e^{inx} dx = \int_0^{2\pi} e^{-imx} e^{inx} dx = \int_0^{2\pi} e^{i(n-m)x} dx \quad (2.144)$$

In the  $n = m$  case we have

$$\int_0^{2\pi} dx = 2\pi \quad (2.145)$$

while if  $n \neq m$  we use

$$\int_0^{2\pi} e^{i(n-m)x} dx = \frac{1}{i(n-m)} \left[ e^{i(n-m)x} \right]_0^{2\pi} = 0 \text{ if } n, m \in \mathbb{Z} \quad (2.146)$$

If we define

$$a_n(x) = \frac{1}{\sqrt{2\pi}} e^{inx} \quad (2.147)$$

we have

$$\int_0^{2\pi} a_n^*(x) a_m(x) dx = \delta_{n,m} \quad (2.148)$$

We may also define the following real functions for  $n \in \mathbb{N}$  ( $p$  stands for periodic)

$$c_0^p(x) = a_0(x) = \frac{1}{\sqrt{2\pi}} \quad (2.149)$$

$$c_n^p(x) = \frac{1}{\sqrt{2}} (a_n(x) + a_{-n}(x)) = \frac{1}{\sqrt{\pi}} \cos(nx) \quad (2.150)$$

$$s_n^p(x) = \frac{1}{\sqrt{2}i} (a_n(x) - a_{-n}(x)) = \frac{1}{\sqrt{\pi}} \sin(nx) \quad (2.151)$$

---

<sup>22</sup>We will later learn that this integral corresponds to a scalar product in a function space.

We have<sup>23</sup>

$$\begin{aligned}
 \int_0^{2\pi} (c_n^p(x))^* c_m^p(x) dx &= \int_0^{2\pi} c_n^p(x) c_m^p(x) dx = \\
 \frac{1}{2} \int_0^{2\pi} (a_n(x) a_m(x) + a_{-n}(x) a_m(x) + a_n(x) a_{-m}(x) + a_{-n}(x) a_{-m}(x)) \\
 &= \frac{1}{2} (\delta_{n,m} + 0 + 0 + \delta_{-n,-m}) = \frac{1}{2} (\delta_{n,m} + \delta_{n,m}) = \delta_{n,m}
 \end{aligned} \tag{2.152}$$

and similarly

$$\begin{aligned}
 \int_0^{2\pi} (s_n^p(x))^* s_m^p(x) dx &= \int_0^{2\pi} s_n^p(x) s_m^p(x) dx \\
 &= \frac{1}{2} \int_0^{2\pi} (a_n(x) a_m(x) + a_{-n}(x) a_m(x) - a_n(x) a_{-m}(x) - a_{-n}(x) a_{-m}(x)) = \delta_{n,m}
 \end{aligned} \tag{2.153}$$

#### 2.8.4 General solution

If we were studying a problem in which  $\rho$  was a complex function, its general solution would be written as

$$\rho(x, t) = \sum_{n=-\infty}^{+\infty} A_n e^{-n^2 t} a_n(x) \tag{2.154}$$

with

$$A_n = \int_0^{2\pi} a_n^*(x) f_{t=0}(x) dx \tag{2.155}$$

But since our initial condition is real, we have

$$A_n^* = \int_0^{2\pi} a_n(x) f_{t=0}(x) dx = \int_0^{2\pi} a_{-n}^*(x) f_{t=0}(x) dx = A_{-n} \tag{2.156}$$

so that

$$\begin{aligned}
 \rho(x, t) &= A_0 a_0(x) + \sum_{n=1}^{\infty} (A_n a_n(x) + A_n^* a_{-n}(x)) = \\
 &= A_0 a_0(x) + \sum_{n=1}^{\infty} (2 \operatorname{Re}(A_n) (a_n(x) + a_{-n}(x)) + 2i \operatorname{Im}(A_n) (a_n(x) - a_{-n}(x)))
 \end{aligned} \tag{2.157}$$

By defining

$$A_0 = C_0, \quad C_n = 2\sqrt{2} \operatorname{Re}(A_n), \quad S_n = 2i\sqrt{2} \operatorname{Im}(A_n) \tag{2.158}$$

---

<sup>23</sup>These relations may obviously be shown to hold also through trigonometric equalities.

## 2.8 Diffusion on the circle

---

we get the general solution in a-dimensional variables

$$\rho(t, x) = \sum_{n=0}^{\infty} C_n c_n^p(x) e^{-n^2 t} + \sum_{n=1}^{\infty} S_n s_n^p(x) e^{-n^2 t} \quad (2.159)$$

with

$$C_n = \int_0^{2\pi} c_n^p(x) f_{t=0}(x), \quad S_n = \int_0^{2\pi} s_n^p(x) f_{t=0}(x) \quad (2.160)$$

or

$$C_0 = \frac{1}{\sqrt{2\pi}} \int_0^{2\pi} f_{t=0}(x) = \sqrt{2\pi} \langle f_{t=0} \rangle_x \quad (2.161)$$

$$C_n = \frac{1}{\sqrt{\pi}} \int_0^{2\pi} \cos(nx) f_{t=0}(x) \quad (2.162)$$

$$S_n = \frac{1}{\sqrt{\pi}} \int_0^{2\pi} \sin(nx) f_{t=0}(x) \quad (2.163)$$

In the  $t \rightarrow \infty$  limit we have

$$\lim_{t \rightarrow \infty} \rho(t, x) = c_0^p \sqrt{2\pi} \langle f_{t=0}(x) \rangle_x = \langle f_{t=0}(x) \rangle_x \quad (2.164)$$

These formulae that can be re-conducted to the dimensional case through eqs. (2.126, 2.127), namely defining

$$c_0^{L,p}(x') = \sqrt{\frac{1}{L}}, \quad c_n^{L,p}(x') = \sqrt{\frac{2}{L}} \cos\left(\frac{2n\pi}{L} x'\right), \quad s_n^{L,p}(x') = \sqrt{\frac{2}{L}} \sin\left(\frac{2n\pi}{L} x'\right) \quad (2.165)$$

$$C_n = \int_0^L c_n^{L,p}(x') f_{t=0}(x'), \quad S_n = \int_0^L s_n^{L,p}(x') f_{t=0}(x') \quad (2.166)$$

$$\rho(t', x') = \sum_{n=0}^{\infty} C_n e^{-\frac{4n^2\pi^2\gamma t'}{L^2}} c_n^L(x') + \sum_{n=1}^{\infty} S_n e^{-\frac{4n^2\pi^2\gamma t'}{L^2}} s_n^L(x') \quad (2.167)$$



## Chapter 3

# The Schrödinger equation

We will discuss briefly how the method of separation of variables may be used to solve simple problems in Quantum Mechanics, i.e. related to behaviours of particles at the atomic or sub-atomic level. The context for the equation describing quantum behaviour will be provided using an historical perspective, i.e. trying to describe how the leading physicists of the time developed and understood these ideas at the beginning of the 20th century. The discussion would be nevertheless extremely simplified and distorted to fit in this short chapter, and should not be considered as a loyal description of facts and opinions, but just as an invitation to the subject.

### 3.1 Historical remarks

#### 3.1.1 Plank

In 1900, the great German physicist Max Planck proposed a solution to one of the most important physical problem of the time, the law at which a non-light reflecting body (“black body”) emits radiation when it is at thermal equilibrium<sup>1</sup>. The details of the problem go beyond the purpose of this course, but in order to find the correct law Planck had to do a very revolutionary hypothesis: light (more properly, electromagnetic radiation), which Maxwell theory had shown to be a continuous wave phenomenon, had to be found in the body only under the form of discrete units (quanta). Light with wave frequency  $\nu$  could be found only in “packets” with energy

$$E = h\nu \tag{3.1}$$

---

<sup>1</sup>This may be a good model of the energy emitted by a star as a function of the star’s temperature.

### 3.1 Historical remarks

---

where  $h$  is the Planck constant. We know that energy has the dimensionality of mass multiplied by a squared velocity, while  $\nu$  is an inverse time, so that  $h$  is an *action*, i.e. a quantity with dimension

$$[h] = [M][L]^2[T]^{-1} \quad (3.2)$$

If we measure masses in kilograms, lengths in meters and time in seconds, i.e. if energy is measured in Joules, the value of  $h$  is

$$h \approx 6.63 \cdot 10^{-34} \text{ J s} \quad (3.3)$$

Planck, who was a conservative man in life and science<sup>2</sup> introduced this law not because he believed light to be formed by particles of energy  $h\nu$ , but simply because the hypothesis allowed him to find the right solution. At the time he proposed his law, it was still not understood that the hypothesis was *necessary* to find the correct law, and that a continuous distribution of energy would lead to non-sensical results (this was proven later by the British Rayleigh and Jeans, and, independently, by Einstein).

#### 3.1.2 Einstein

##### Photoelectric effect

As we have just seen, Albert Einstein had arrived to the conclusion that Planck's law was not just a mathematical trick, since he could prove that if the radiation had a continuous energy distribution, the black body would need an infinite energy before arriving at thermal equilibrium. Einstein had also learned about experiments saying that the energy of electrons released by a metal when electromagnetic radiation (light) was sent on it were independent on the radiation intensity, a very surprising fact since the energy of a wave is related to its intensity.

---

<sup>2</sup>Max Planck, who received the Nobel prize in 1918 for his work on the black body radiation, strongly disliked Nazism but tried to convince German physicists to remain to work in Germany under the regime. He never openly opposed Nazi policies, but tried (with no success) to use his authority in order to defend some Jewish colleagues, talking once to the same Hitler (these are Planck's memories of their talk "*In response to my comment that it would essentially be a self-inflicted catastrophe if valuable Jews were forced to emigrate, as we urgently need their scientific work, and that this will otherwise benefit primarily other countries, he (Hitler) did not comment any further. He turned instead to general chitchat and concluded by saying: People say that I occasionally suffer from neurasthenia. This is slander. I have nerves of steel. He then slapped his knee hard, spoke at an increasingly faster rate and worked himself up into such a rage that there was nothing else for me to do but to remain silent and leave.*"). Planck's son Erwin was sentenced to death in 1944 for taking part to a plot aimed at killing Hitler.

Einstein thus stated that light was composed of particles of energy  $h\nu$ , where  $\nu$  is the frequency of the wave. The intensity of the wave was related to the *number* of particles in the wave. Nevertheless, when interacting with individual electrons, it would be the energy of the particle (*photon*) to decide if the electron received enough energy to leave the metal. The energy of the released electron would be, according to Einstein's prediction,

$$E = h\nu - \phi \tag{3.4}$$

$\phi$  being the minimum energy needed to release the electron from the metal. Intensity, Einstein predicted, would be related to the number of released electrons, not to their energy.

Einstein wrote his theory in 1905, and his predictions were experimentally confirmed in 1916. The theory was not easily accepted. Einstein did not explain why light usually behaved like a wave, and not as a particle, in other situations. The particle theory of light, proposed in the past by Newton, had been rejected empirical data. For these reasons even the most prominent physicists of the time considered Einstein's theory as wrong, and even when data supporting it emerged, most physicists believed the quantization to be due to interaction with matter, and not intrinsic in light itself. Only after the quantum theory for particles (as electrons) was developed, and was understood that the particle-wave duality is universal in nature, was it possible to better understand Einstein's theory<sup>3</sup>.

### 3.1.3 Relativity

In the same year, Einstein proposed his relativity theory, which extends some results previously introduced by the Dutch physicist Lorentz, and largely parallels a theory introduced at the same time by the French mathematician Poincaré<sup>4</sup>. In non relativistic physics, the energy of a particle in a potential  $U$  is given by

$$E = \frac{1}{2}mv^2 + U(\mathbf{x}) \tag{3.5}$$

This is often written in physics as

$$H(\mathbf{x}, \mathbf{p}) = \frac{p^2}{2m} + U(\mathbf{x}) \tag{3.6}$$

---

<sup>3</sup>The quantum theory of light, being inherently relativistic, presents more difficulties than the quantum theory of electrons, and was developed only in a second time.

<sup>4</sup>The difference between their approaches is subtle and debated between historians. Most nevertheless agree that Einstein was the first one to state clearly that the new dynamics and kinematics derived just by asking that all laws of physics, including the value of the velocity of light, were the same in all inertial frames.

### 3.1 Historical remarks

---

where  $H$  is the Hamiltonian function of position  $\mathbf{x}$  and momentum  $\mathbf{p}$ , a function that plays a fundamental role in the theoretical study of mechanics. The momentum  $\mathbf{p}$  is, for this simple system, defined as

$$\mathbf{p} = m\mathbf{v} \quad (3.7)$$

so that Newton's second law may be written as

$$\frac{dp_i}{dt} = -\partial_{x_i} H(\mathbf{x}, \mathbf{p}) = -\partial_{x_i} U(\mathbf{x}) = F_i(\mathbf{x}) \quad (3.8)$$

In relativistic physics we still have

$$\frac{dp_i}{dt} = F_i(\mathbf{x}) \quad (3.9)$$

but now the momentum is given by

$$\mathbf{p} = \frac{m\mathbf{v}}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (3.10)$$

where  $c$  is the velocity of light (this causes the inertia of a particle to diverge when the velocity approaches  $c$ , so that no force may bring a particle to a velocity higher than  $c$ .) The energy of a free particle (i.e. in absence of force) is now defined by

$$E = \sqrt{m^2 c^4 + p^2 c^2} \quad (3.11)$$

which may be solved for

$$E = \frac{mc^2}{\sqrt{1 - \frac{v^2}{c^2}}} \quad (3.12)$$

so that

$$\frac{p}{E} = \frac{v}{c^2} \quad (3.13)$$

Let us now assume there is a particle without mass,  $m = 0$ . Eq. (3.11) becomes

$$E = pc \quad (3.14)$$

and substituting in (3.13) gives

$$v = c \quad (3.15)$$

For this reason, the "light particles" that Einstein introduced in his work on the photoelectric effect have necessarily zero mass (since they move with velocity  $c$ <sup>5</sup>). The theory of Relativity was soon accepted by the leading physicists of the time<sup>6</sup>.

---

<sup>5</sup>It is clear from eq. (3.11) that if  $m > 0$  then  $E > pc$ .

<sup>6</sup>In particular those of Einstein's generation or younger. Others, while agreeing with the mathematical results, maintained a different philosophical approach, closer to the one

### 3.1.4 De Broglie

With his 1924 Ph. D. thesis, the French physicist Luis de Broglie made an important step in understanding the relation between wave and particle behaviour.

#### About waves

Let us first start with some notation. If we have a periodic phenomenon (in our case the oscillation of a wave) with period  $T$ , its frequency is given by

$$\nu = \frac{1}{T} \quad (3.16)$$

We often express periodic phenomena through trigonometric functions<sup>7</sup>,

$$f(t) = A \cos(\omega t + \phi) = A \cos(\omega(t + T) + \phi) = f(t + T) \quad (3.17)$$

This implies

$$\omega T = 2\pi \Rightarrow \omega = 2\pi\nu \quad (3.18)$$

where  $\omega$  is called the angular velocity. Relation (3.1) may thus be written as

$$E = \hbar\omega, \quad \hbar = \frac{h}{2\pi} \quad (3.19)$$

where we have defined the useful constant  $\hbar$  (h-bar).

If a wave has velocity  $c$ , it will cover a distance  $cT$  in time  $T$ . This distance is called the wave length

$$\lambda = \frac{c}{\nu} \quad (3.20)$$

Let us write a function which is periodic in space and time

$$\Phi(t, x) = A \cos(kx - \omega t + \phi) \quad (3.21)$$

At a fixed point  $\bar{x}$  this function will be oscillating in time with a period  $T$  such that  $\omega T = 2\pi$ , as discussed above. If observed at fixed  $t$ , the function

---

of Lorentz and Poincare. It is nevertheless interesting that less important physicists had a harder time to accept relativity theory. The Nobel Prize committee received, in the years following the publishing of Einstein's Special and General (Gravitation) Relativity theory, many endorsements to give the prize to Einstein, but the members of the committee were reluctant to give the prize for relativity theory, and eventually decided to give him the prize for the photoelectric effect in 1921. Interestingly, a few of the eminent physicists proposing Einstein were actually doubtful with regards to the correctness of the revolutionary theory according to which light is composed by particles!

<sup>7</sup>  $\cos(\omega t + \phi) = \cos(\phi) \cos(\omega t) - \sin(\phi) \sin(\omega t) = C_1 \cos(\omega t) + C_2 \sin(\omega t)$

### 3.1 Historical remarks

---

looks like oscillating with a period (distance between two crests)  $\lambda$  such that  $k\lambda = 2\pi$ , or

$$k = \frac{2\pi}{\lambda} \quad (3.22)$$

where  $k$  is called the *wave number*. Is this the same  $\lambda$  in equation (3.20)?

We have mentioned above that the oscillation of a (1D) wave is described by a differential equation in the form

$$\partial_t^2 \Phi(t, x) = c^2 \partial_x^2 \Phi(t, x) \quad (3.23)$$

If we substitute  $\Phi$  from eq. (3.21) in eq. (3.23) we obtain

$$\omega^2 \Phi(t, x) = c^2 k^2 \Phi(t, x) \quad (3.24)$$

i.e. we have a solution provided that

$$\omega = \pm ck \Rightarrow 2\pi\nu = \pm c \frac{2\pi}{\lambda} \Rightarrow \lambda = \frac{c}{\nu} \quad (3.25)$$

(we have assumed to define  $\nu$  and  $\lambda$  as positive), so that  $\lambda$  appearing in the periodic solution is indeed the distance the wave covers in  $T$ .

### De Broglie's matter waves

De Broglie noticed that for a photon, using eqs. (3.14,3.20,3.22), we have

$$p = \frac{h\nu}{c} = \frac{h}{\lambda} = \hbar k \Rightarrow \lambda = \frac{h}{p} \quad (3.26)$$

and made the hypothesis that a wave length  $\lambda$  was associated to any particle of momentum  $p$  according to the law above.

De Broglie's idea was very easy to test<sup>8</sup>: it was enough to have a beam of electrons with momentum  $p$  through a crystal with lattice distance  $b$  and see if the beam generated a diffraction pattern corresponding to a wave of wavelength  $\lambda$ . It did, and de Broglie (1929) and experimental physicists Davisson and Thomson (1937, for two different experiments performed in USA and Britain) got the Nobel prize.

#### 3.1.5 Heisenberg

In 1925, the 23 year old German physicist Werner Heisenberg formulated the first successful and completely consistent description of atomic phenomena,

---

<sup>8</sup>In principle, performing actual experiments nobody did before is never easy.

through a quite obscure model called *matrix mechanics*. The model was perfected and mathematically brushed up in collaboration with Max Born<sup>9</sup> and Pascual Jordan, two other German physicists, and used by the Austrian Wolfgang Pauli<sup>10</sup> to find the spectrum (possible energies of electrons) of the Hydrogen atom.

Despite its early success, the model, who lead Heisenberg to the 1932 Nobel Prize<sup>11</sup>, was not extremely loved by contemporary physicists. Many just disliked its mathematical formalism, but a few disliked the underlying philosophy: Heisenberg claimed that there was nothing more in physics that what could be directly observed, no underlying substrate, and based his theory on this philosophical principle.

### 3.1.6 Schrödinger

Between those that did not like the philosophical principles underlying Heisenberg model there was Albert Einstein, who believed that the truth was in the direction shown by de Broglie's ideas. Through Einstein, the Austrian mathematical physicist Erwin Schrödinger learned about de Broglie waves, namely that a free (non interacting particle) could be described through relations (3.1,3.26) by a periodic function

$$\Psi(t, \mathbf{x}) = e^{i\left(\frac{\mathbf{p}\cdot\mathbf{x}}{\hbar} - \frac{Et}{\hbar}\right)} \quad (3.27)$$

Here we used the complex formulation (whose real part gives a cosine and imaginary part a sine), and generalised to the 3D case<sup>12</sup> the relations

$$\frac{E}{\hbar} = \omega, \quad \frac{\mathbf{p}}{\hbar} = \mathbf{k} \quad (3.28)$$

---

<sup>9</sup>1954 Nobel Prize for his interpretation of the statistical nature of Quantum Mechanics, to be discussed later.

<sup>10</sup>1945 Nobel Prize for his exclusion principle, that describes the impossibility of two electrons to be in the same physical state.

<sup>11</sup>Heisenberg trajectory during the dark years of National Socialism may recall the experience of the much older Planck. A proud and patriotic German, never thought of leaving the country. He was nevertheless accused of being a "white Jew", since he taught Einstein's theories in his university courses. Eventually, due to his scientific stature, he was put in charge of theoretical work related to the development of the German Atom bomb, a role that made him be partially ostracised from the scientific community at the end of the conflict, and ruined his relation with his mentor, the Danish Jew Niels Bohr. Nevertheless, Heisenberg always stated that he worked only to develop civil nuclear power. After being arrested by Allied forces, and detained in Britain with other fellow scientists, he gave to his fellow inmates a lecture on the functioning of the Hiroshima bomb, showing, according to some historians, that he actually did his best *not* to build a Nazi bomb.

<sup>12</sup>Using  $\nabla^2 e^{i\mathbf{k}\cdot\mathbf{x}} = -k^2 e^{i\mathbf{k}\cdot\mathbf{x}}$ .

### 3.1 Historical remarks

---

He started lecturing about this idea, in opposition to Heisenberg formulation, until someone commented that “a wave needs a wave equation”. Schrödinger, who was at the time 38, an age at which in general theoretical physicists have already produced their best works, decided to spend a few days in a mountain cabin on the Alps with a woman that was not exactly his wife<sup>13</sup> and used (part of) his time to write down the wave equation that takes its name. He started from the Hamiltonian function (3.6)

$$E = H(\mathbf{x}, \mathbf{p}) = \frac{p^2}{2m} + U(\mathbf{x}) \quad (3.29)$$

He noticed that, according to eq. (3.27), the energy is obtained by operating on  $\Psi$  using the following differential operator  $\hat{E}$

$$E\Psi = \hat{E}\Psi = i\hbar\partial_t\Psi \quad (3.30)$$

In an equivalent way, the ( $x$  component of the) momentum is obtained by

$$p_x\Psi = \hat{p}_x\Psi = -i\hbar\partial_x\Psi \quad (3.31)$$

He thus replaced the quantities  $E$  and  $p$  in eq. 3.29 with these differential operators acting on  $\Psi$ . In order to replace  $p^2$ , he used

$$p^2 = \mathbf{p} \cdot \mathbf{p} \Rightarrow \hat{p}^2 = \sum_i (-i\hbar\partial_i)(-i\hbar\partial_i) = -\hbar^2\nabla^2 \quad (3.32)$$

He was left with the potential  $U$ , that he decided to introduce as an operator that multiplies  $\Psi$  by  $U$ , since, as we saw with eq. (2.55), such a choice gives a linear operator.

He thus introduced the following equation that takes his name

$$\hat{E}\Psi(t, \mathbf{x}) = \hat{H}\Psi(t, \mathbf{x}) = \frac{\hat{p}^2}{2m}\Psi(t, \mathbf{x}) + U(\mathbf{x})\Psi(t, \mathbf{x}) \quad (3.33)$$

or, explicitly

$$i\hbar\partial_t\Psi(t, \mathbf{x}) = -\frac{\hbar^2}{2m}\nabla^2\Psi(t, \mathbf{x}) + U(\mathbf{x})\Psi(t, \mathbf{x}) \quad (3.34)$$

Before discussing the meaning of this equation it is better to find its solution, just as Schrödinger did.

---

<sup>13</sup>In 1939, Schrödinger moved to Dublin, invited by the Irish Prime Minister (later President) De Valera, when he had to leave Austria due to his positions against the Nazi party (smart people had a tendency to dislike Nazism), where he openly lived with his wife and the mother of one of his daughters. While in Dublin, he had other two daughters from different women.



## 3.2 A simple 1D problem in Quantum mechanics: a trapped particle

Let us suppose we have a particle trapped in a area of length  $L$ . We will assume, for reasons of mathematical simplicity, the problem to be 1D.

### 3.2.1 Boundary conditions

This trap is realised by asking that the particle moves freely ( $U = 0$ ) in the  $[0, L]$  area, but it cannot exit it. Namely we will study the equation in such interval with the boundary conditions

$$\Psi(t, 0) = \Psi(t, L) = 0 \quad (3.35)$$

Although we still did not discuss the meaning of  $\Psi$ , we are implying that  $\Psi$  somehow tells us where the particle is, for example  $\Psi$  may be related to the particle's "density" (reality is more complex, see section 3.3). Since the equation is second order in space, these conditions should be enough to fix the behaviour of the solution. The equation is first order in time, so that knowledge of

$$\Psi(0, x), \quad x \in [0, L] \quad (3.36)$$

should suffice<sup>14</sup>.

### 3.2.2 Separation of variables: the Time-independent Schrödinger equation

The equation is linear and the boundary conditions are homogeneous, so we will proceed with separation of variables.

$$\Psi(t, x) = T(t)X(x) \quad (3.37)$$

This leads to

$$X(x)(i\hbar\partial_t T(t)) = T(t) \left( -\frac{\hbar^2}{2m} \partial_x^2 X(x) \right) \quad (3.38)$$

and, dividing by  $\Psi$ ,

$$\frac{i\hbar\partial_t T(t)}{T(t)} = C = \frac{-\frac{\hbar^2}{2m} \partial_x^2 X(x)}{X(x)} \quad (3.39)$$

Let us reflect on the role of  $C$ . We introduced the operator

$$\hat{E} = i\hbar\partial_t \quad (3.40)$$

---

<sup>14</sup>Schrödinger equation is basically a diffusion equation with a  $-i$  in front of the time derivative.

### 3.2 A simple 1D problem in Quantum mechanics: a trapped particle

---

because it gave the energy when applied to  $\Psi$  in the form of a de Broglie wave, eq. (3.27). Here we have

$$\hat{E}\Psi(t, x) = i\hbar\partial_t T(t)X(x) = X(x)i\hbar\partial_t T(t) = CX(x)T(t) = C\Psi(t, x) \quad (3.41)$$

it looks thus reasonable to name  $C = E$ , the *energy of the particle*.

We are thus left with two equations

$$i\hbar\partial_t T(t) = ET(t) \quad (3.42)$$

and

$$-\frac{\hbar^2}{2m}\partial_x^2 X(x) = EX(x) \quad (3.43)$$

The first equation is solved as

$$T(t) = A'e^{-i\frac{E}{\hbar}t} \quad (3.44)$$

This is obviously the time part of eq. (3.27), but now we obtained it from a wave equation, so that Schrödinger's critics could be silenced. It should be clear that, provided the potential  $U$  is time-independent, the procedure leading to eq. (3.44) will not depend on  $U$ . The time dependence of  $\Psi$  for a particle with a given energy  $E$  will be always given by eq. (3.44). We will thus be left with the problem of finding the space dependence of  $\Psi$ , which is given by the *Time-independent Schrödinger equation*, eq. (3.43).

We may notice, given the discussion above, that, when dealing with the Schrödinger equation, the separation of variable process has a deep physical meaning, since leads us to obtaining *physical states with a defined energy*.

#### 3.2.3 Solution of the time-independent Schrödinger equation

The equation may be written as

$$\partial_x^2 X(x) = -\frac{2mE}{\hbar^2}X(x) \quad (3.45)$$

We have as usual 3 cases depending on the sign of the energy

(I)  $E < 0$ ,  $\frac{2mE}{\hbar^2} = \lambda^2$

The solution to

$$\partial_x^2 X(x) = \lambda^2 X(x) \quad (3.46)$$

is

$$X(x) = Ae^{\lambda x} + Be^{-\lambda x} \quad (3.47)$$

Clearly

$$X(0) = 0 \Rightarrow B = -A \quad (3.48)$$

and

$$X(L) = A(e^{\lambda L} - e^{-\lambda L}) \quad (3.49)$$

cannot be zero for  $A \neq 0$ , so that there is no solution for negative energy.

This is not surprising. The energy for a classical particle is

$$E = \frac{1}{2}v^2 + U(x) \geq U(x) \quad (3.50)$$

so that by fixing the potential  $U = 0$  we are (from a classical view point) forcing the energy to be  $E \geq 0$ . We have just seen that this result applies also to the quantum problem.

(II)  $E = 0, \frac{2mE}{\hbar^2} = 0$

The solution is

$$X(x) = Ax + B \quad (3.51)$$

and as usual the boundary conditions impose  $A = B = 0$ . But if  $\Psi$  tells us where the particle is,  $\Psi = 0$  means no particle at all<sup>15</sup>.

While in the classical physics case we could have zero energy states, corresponding to the particle standing still somewhere between 0 and  $L$ , this is not possible in Quantum Mechanics. According to **Heisenberg indeterminacy principle**, it is impossible to know the position and velocity of a particle at the same time, and thus it is impossible for a particle to stand still somewhere<sup>16</sup>.

(III)  $E > 0, \frac{2mE}{\hbar^2} = -\lambda^2$

Since we are dealing with a equation that includes a complex term ( $i$  in front of  $\partial_t$ ) it is better to write the solution in complex form

$$X(x) = \bar{A}e^{i\lambda x} + \bar{B}e^{-i\lambda x} \quad (3.52)$$

Anyway, from  $X(0) = 0$  we get  $B = -A$  or

$$X(x) = A' \sin(\lambda x) \quad (3.53)$$

To have  $X(L) = 0$  we require  $\lambda L = n\pi, n \in \mathbb{N}$  or

$$\lambda = \frac{n\pi}{L} \Rightarrow E = \frac{\lambda^2 \hbar^2}{2m} = \frac{n^2 \pi^2 \hbar^2}{2mL^2} = \frac{n^2 \hbar^2}{8mL^2} = n^2 E_0 \quad (3.54)$$

---

<sup>15</sup>This statement will be justified in section 3.3.

<sup>16</sup>Heisenberg's name popped out as a result of Schrödinger's theory... more on this later.

### 3.3 The Philosophy of Quantum Mechanics

---

where we defined

$$E_0 = \frac{n^2 h^2}{8mL^2} \quad (3.55)$$

Classically, the particle's energy could assume any value  $mv^2/2 \geq 0$ , but a quantum particle described by Schrödinger's equation can only assume *discrete* values of the energy, such as

$$E_0, \quad 4E_0, \quad 9E_0, \quad 16E_0, \quad 25E_0, \text{ etc.} \quad (3.56)$$

#### 3.2.4 General solution

Eq. (3.53) may be written as (refer to eq. 2.84)

$$X(x) = A' \sin\left(\frac{n\pi}{L}x\right) = A s_n^L(x) \quad (3.57)$$

so that a general solution may be written as

$$\Psi(x, t) = \sum_{n=0}^{\infty} A_n e^{-i\frac{E_n}{\hbar}t} s_n^L(x) \quad (3.58)$$

with

$$A_n = \int_0^L s_n^L(x) \Psi(0, t) dx \quad (3.59)$$

Since the functions  $s_n^L$  represent the solutions of the time independent equation corresponding to the energy  $E_n$ , we will call them also  $\psi_{E_n}$  and write the general solution as

$$\Psi(x, t) = \sum_{n=0}^{\infty} A_n e^{-i\frac{E_n}{\hbar}t} \psi_{E_n}(x) \quad (3.60)$$

But, what does this mean?

### 3.3 The Philosophy of Quantum Mechanics

What Schrödinger actually did in the days spent on the Alps with (one of) his mistress(es) was to compute the possible energy states of the hydrogen atom. As we will see in a following chapter, also these states can assume only discrete values<sup>17</sup>. That energies of electrons in atoms assume discrete values was well known by experiments. The Danish physicist Niels Bohr had proposed in 1912 a model, that was based on some ad hoc assumptions following Planck quantum hypothesis, that constrained electrons to move only

---

<sup>17</sup>From this discreteness comes the name quantum. *Quantum* is the Latin for *how much*, and it is related to the English *quantity*.

on some specific orbits, and could explain the energies of some atoms<sup>18</sup>. But now Schrödinger was able to derive the same results from a completely consistent model, and had written an equation to derive the energy of an electron in any physical problem, just by knowing the potential  $U$ . Furthermore, he was able to derive Heisenberg's theory from his own, showing, he believed, that there was no need of such a theory with its philosophical implications. For his work on Wave Mechanics<sup>19</sup> he received the 1933 Nobel prize.

But he still had to provide an interpretation for his *wave function*  $\Psi$ . The wave function is a complex function, and cannot be directly observed. One can nevertheless define a real quantity from it

$$\rho(t, \mathbf{x}) = \Psi^*(t, \mathbf{x})\Psi(t, \mathbf{x}) \quad (3.61)$$

Other real quantities could obviously be defined, such as  $\text{Re}\Psi$  and  $\text{Im}\Psi$ . Nevertheless, the quantity defined in eq. (3.61) satisfies a conservation law similar to eq. (1.99), and thus leads to Schrödinger's interpretation of it as "a particle density".

This interpretation soon showed to have a few conceptual problems<sup>20</sup>. In 1926, Max Born suggested that  $\rho$ , if better defined as

$$\rho(t, \mathbf{x}) = \frac{\Psi^*(t, \mathbf{x})\Psi(t, \mathbf{x})}{\int_V \Psi^*(t, \mathbf{x})\Psi(t, \mathbf{x})d\mathbf{x}} \quad (3.62)$$

so that its integral over all space  $V$  is one,  $\rho$  gives the *probability* of finding a particle in  $\mathbf{x}$ . The quantum world is no more *deterministic*, as the classical one, in which the initial state (position and velocity) determines all the evolution of the system. Schrödinger equations are still deterministic regarding the evolution of  $\Psi$ , but when an experiment is performed, the result of the experiment is probabilistic<sup>21</sup>.

This framework is very general. The British physicist Paul Dirac, who shared the 1933 Nobel Prize with Schrödinger, showed that both wave and matrix mechanics may be derived from a more general formalism. Let us consider for example a de Broglie wave, eq. (3.27). If we apply the operator  $\hat{p}$  to it, we obtain

$$\hat{p}\Psi = p\Psi \quad (3.63)$$

But if we apply it to eq. (3.53) we get

$$\hat{p}\Psi = -i\hbar\partial_x \sin\left(\frac{n\pi}{L}x\right) = -i\hbar\frac{n\pi}{L} \cos\left(\frac{n\pi}{L}x\right) \quad (3.64)$$

---

<sup>18</sup>Bohr won the 1922 Nobel Prize for this model.

<sup>19</sup>Quantum Mechanics using Schrödinger's formalism.

<sup>20</sup>E.g., when used to describe more than a particle.

<sup>21</sup>The dimensionality of  $\Psi$  is, in  $D$  dimensions,  $[\Psi] = [L]^{-D/2}$ . Why?

### 3.3 The Philosophy of Quantum Mechanics

---

The operator  $\hat{p}$  has no more the effect of multiplying  $\Psi$  by a number, so that the physical state has no defined momentum. What will be the result of an experiment? From eq. (3.49) we see that the sine is the sum of two terms that have a specified momentum, each of them with a different sign. An experiment that measures the momentum of the particle in a state of given energy will give one of these two values, each one with probability 1/2. If the experiment measures the position, it will give a value between 0 and  $L$  with probability proportional to  $\sin^2(\frac{n\pi}{L}x)$ . When we have the general solution eq. (3.60), that includes many different energies, the probability of an experiment to give a given energy is

$$p_{E_n} = \frac{A_n^* A_n}{\sum_n A_n^* A_n} \quad (3.65)$$

Macroscopic objects are composed of many particles, whose dynamics is described by Quantum Mechanics. So Quantum Mechanics should be more fundamental than Classical Mechanics, and should be applied to any object. The reason we do not realise the weird quantum behaviour in our everyday life is related to the small value of  $h$ . The minimum energy of a particle in a trap is discrete,

$$E_0 = \frac{n^2 h^2}{8mL^2} \quad (3.66)$$

If we use as mass the electron mass, of order  $10^{-31}$  Kg, and as  $L = 10^{-10}$  m, comparable to the size of an atom, we get  $E_0 \approx 10^{-17}$  J. This is a relevant energy for an electron, that has an electrostatic energy of  $\approx 10^{-19}$  J when in a field of 1 V. But if we consider a ball of 1 Kg moving in a trap of 1 m, we get  $E_0 \approx 10^{-67}$ , which is basically zero from a macroscopic point of view, so that the energy appears to be continuum.

The standard interpretation of Quantum Mechanics has been developed by Niels Bohr's group (including Heisenberg). It makes a distinction between the mathematical structure of the wave equation, and the results of experiments. Only the latter are observable, and questions about what happens to the particle before the experiment result are meaningless. The predictions of Quantum Mechanics are in perfect agreement with the experiments, that are the only thing that may be tested.

This interpretation baffled Schrödinger (and Einstein). Also the experimental equipment, and the experimenter themselves, are composed of particles. How can we make the distinction between Quantum and not Quantum? Schrödinger came out with this example. A nucleus may be in different energy states. The passage between two different states (two  $\psi_E$ ) corresponds to the emission of radiation. Observing the emission of radiation is an experiment, and has a probabilistic outcome. Now, says Schrödinger, let us

connect a mechanism to the emission of radiation, and let us have this mechanism kill a cat closed in a box with the atom and mechanism, in case the atom decays. Our experimental result is the cat being dead or alive. According to Schrödinger, if Bohr's interpretation is correct, we should have a quantum function describing the whole system, and thus the cat being both alive and dead, until we open the box<sup>22</sup>.

The predictions of Quantum Mechanics are very reliable. So reliable that in general for long time many physicists adopted a “compute and shut up” approach, i.e. they ignored philosophical questions, leaving them to people like Einstein, Schrödinger, Bohr and other Nobel prizes that did not need to care too much about how they used their time since they already obtained so much<sup>23</sup>. Nevertheless, physicists have come to apply Quantum Mechanics outside the realm of atoms and nuclei (for example, to study the whole universe) so that questions such as “what happens before the experiment is performed” became quite relevant. The philosophical debate is not over.

### 3.4 Getting rid of $\hbar$

To get a clear idea of what our results imply, we worked using the original variables  $t$  and  $x$  with dimensions of time and length. Nevertheless, many physicists, in particular theoretical physicists, like to work in a unit system in which  $\hbar = 1$ . This not only allows them to get rid of  $\hbar$  in their expressions, but introduces also a “natural” unit system, i.e. a system based not on human conventions but on Nature's rules<sup>24</sup>.

Let us write the 1D Schrödinger equation

$$i\hbar\partial_t\Psi(t, \mathbf{x}) = -\frac{\hbar^2}{2m}\partial_x^2\Psi(t, x) + U(\mathbf{x})\Psi(t, x) \quad (3.67)$$

---

<sup>22</sup>In his words, *One can even set up quite ridiculous cases. A cat is penned up in a steel chamber, along with the following device (which must be secured against direct interference by the cat): in a Geiger counter, there is a tiny bit of radioactive substance, so small, that perhaps in the course of the hour one of the atoms decays, but also, with equal probability, perhaps none; if it happens, the counter tube discharges and through a relay releases a hammer that shatters a small flask of hydrocyanic acid. If one has left this entire system to itself for an hour, one would say that the cat still lives if meanwhile no atom has decayed. The psi-function of the entire system would express this by having in it the living and dead cat (pardon the expression) mixed or smeared out in equal parts.*

<sup>23</sup>And in general, this approach went in favour of Bohr's school of thinking that you shouldn't ask yourself what happens in the microscopic system.

<sup>24</sup>Since it uses a universal constant as a unit measure. In an equivalent way, the velocity of light  $c$  is usually fixed to one (this accounts to measure distances in time, as for the celebrated “light year”)

### 3.4 Getting rid of $\hbar$

---

and perform the change

$$t' = \frac{t}{\hbar} \Rightarrow \partial_t = \frac{1}{\hbar} \partial_{t'} \quad (3.68)$$

$$x' = \frac{x}{\hbar} \Rightarrow \partial_x = \frac{1}{\hbar} \partial_{x'} \quad (3.69)$$

The equation in the new variables becomes

$$i\partial_{t'}\Psi(t', x') = -\frac{1}{2m}\partial_{x'}^2\Psi(t', x') + U(x')\Psi(t', x') \quad (3.70)$$

i.e. we got rid of  $\hbar$ . To better understand how this happened, let us remember (eq. 3.2) that  $\hbar$  had the dimensionality of an action, i.e. energy multiplied by time. Actions are extremely important in physics, since it may be shown that a classical (i.e., not quantum) particle moves on a trajectory that minimises<sup>25</sup> a quantity  $S$ , called indeed action, with the dimension of energy multiplied by time.

If we define a quantity  $S'$ , given by an energy multiplied by time *in the new scaled coordinates*, we have that

$$[S'] = [M][L']^2[T']^{-2}[T'] = [M][L']^2[T']^{-1} = [M][L]^2[\hbar]^{-2}[T]^{-1}[\hbar] = [S][\hbar]^{-1} = [*]^0 \quad (3.71)$$

This equation tells us two things

1. The new action is a-dimensional (the  $[*]^0$  term)
2. Its value is given in units of  $\hbar$  (the  $[S][\hbar]^{-1}$  term)

Indeed our change of coordinates introduces a “natural” system in which actions are counted as multiples of  $\hbar$ .

By working in such a system, we would have found, for the energy of the system

$$E'_n = n^2 E'_0 \quad (3.72)$$

with

$$E'_0 = \frac{\pi^2}{2mL'^2} \quad (3.73)$$

How can we regain from this expression the one including the  $\hbar$  factors?

One way of doing it is to notice that

$$L' = \frac{L}{\hbar} \quad (3.74)$$

---

<sup>25</sup>Actually extremises.



and substitute. But in some situations, we may not have such a simple expression. A more general approach is to write

$$E_0 = \frac{\pi^2}{8mL^2} \hbar^k \quad (3.75)$$

and determine  $k$  by dimensional analysis

$$[M][L]^2[T]^{-2} = [E_0] = [E]^k [T]^k [L]^{-2} [M]^{-1} = [M]^{k-1} L^{2k-2} [T]^{-k} \quad (3.76)$$

that implies  $k = 2$ .

### 3.5 The particle in a 3D box

Let us solve our first 3D problem, by asking our particle to be trapped inside a 3D box. We have

$$U(\mathbf{x}) = 0 \text{ for } x \in [0, L_x] \text{ and } y \in [0, L_y] \text{ and } z \in [0, L_z] \quad (3.77)$$

and

$$\Psi(t, 0, y, z) = \Psi(t, L_x, y, z) = \Psi(t, 0, x, y) = \Psi(t, x, L_y, z) = \Psi(t, x, y, 0) = \Psi(t, x, y, L_z) = 0 \quad (3.78)$$

Let us use the separation of variables technique by asking

$$\Psi(t, x, y, z) = T(t)X(x)Y(y)Z(z) \quad (3.79)$$

so that the Schrödinger equation becomes

$$\begin{aligned} X(x)Y(y)Z(z)(i\hbar\partial_t T(t)) = \\ - \frac{\hbar^2}{2m} \left( T(t)Y(y)Z(z)\partial_x^2 X(x) + T(t)X(x)Z(z)\partial_y^2 Y(y) + T(t)X(x)Y(y)\partial_z^2 Z(z) \right) \end{aligned} \quad (3.80)$$

we divide by  $\Psi$  and obtain as usual<sup>26</sup>

$$\frac{i\hbar\partial_t T(t)}{T(t)} = E = -\frac{\hbar^2}{2m} \left( \frac{\partial_x^2 X(x)}{X(x)} + \frac{\partial_y^2 Y(y)}{Y(y)} + \frac{\partial_z^2 Z(z)}{Z(z)} \right) \quad (3.81)$$

As we stated above, provided that  $U$  is time independent, the solution of the time function is always

$$T(t) = A' e^{-i\frac{E}{\hbar}t} \quad (3.82)$$

---

<sup>26</sup>Since the two sides of the equal are functions of unrelated variables.

### 3.5 The particle in a 3D box

---

To solve the time independent equation that gives the physical state (wave function) corresponding to the energy  $E$  we write

$$E + \frac{\hbar^2}{2m} \frac{\partial_x^2 Z(z)}{Z(z)} = -\frac{\hbar^2}{2m} \left( \frac{\partial_x^2 X(x)}{X(x)} + \frac{\partial_x^2 Y(y)}{Y(y)} \right) \quad (3.83)$$

Now, the two sides of the equality are functions of unrelated variables, and they have to be constant. We name such constant  $E_{xy}$ . We have

$$-\frac{\hbar^2}{2m} \frac{\partial_x^2 Z(z)}{Z(z)} = E - E_{xy} \equiv E_z \quad (3.84)$$

We already solved this equation in the 1D case. We get  $E_z > 0$ ,

$$E_z = n_z^2 E_0^z, \quad E_0^z = \frac{\hbar^2}{8mL_z^2} \quad (3.85)$$

with  $n_z \in \mathbb{N}$ , and<sup>27</sup>

$$Z(z) = s_{n_z}^{L_z}(z) \quad (3.86)$$

We proceed in the same way and get from

$$E_{xy} = -\frac{\hbar^2}{2m} \left( \frac{\partial_x^2 X(x)}{X(x)} + \frac{\partial_x^2 Y(y)}{Y(y)} \right) \quad (3.87)$$

the equation

$$E_{xy} + \frac{\hbar^2}{2m} \frac{\partial_x^2 Y(y)}{Y(y)} = E_x = -\frac{\hbar^2}{2m} \frac{\partial_x^2 X(x)}{X(x)} \quad (3.88)$$

and, solving for  $X$

$$E_x = n_x^2 E_0^x, \quad E_0^x = \frac{\hbar^2}{8mL_x^2} \quad (3.89)$$

with  $n_x \in \mathbb{N}$ , and

$$X(x) = s_{n_x}^{L_x}(x) \quad (3.90)$$

Finally, from

$$-\frac{\hbar^2}{2m} \frac{\partial_x^2 Y(y)}{Y(y)} = E_{xy} - E_x \equiv E_y \quad (3.91)$$

we get

$$E_y = n_y^2 E_0^y, \quad E_0^y = \frac{\hbar^2}{8mL_y^2} \quad (3.92)$$

with  $n_y \in \mathbb{N}$ , and

$$Y(y) = s_{n_y}^{L_y}(y) \quad (3.93)$$

---

<sup>27</sup>We do not write the multiplicative constant. It is usual in Quantum mechanics to use functions that have integral of their square modulus equal to 1, as our  $s$  functions are, due to the probabilistic interpretation eq. (3.62).

Clearly, from our definitions we have

$$E = E_x + E_y + E_z = n_x^2 E_0^x + n_y^2 E_0^y + n_z^2 E_0^z \quad (3.94)$$

where the  $n$  are *positive* integers. For a cube, it simplifies to

$$E = n^2 E_0, \quad n^2 = n_x^2 + n_y^2 + n_z^2 \quad (3.95)$$

The general solution<sup>28</sup> is

$$\Psi(t, x, y, z) = \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} \sum_{n_z=1}^{\infty} A_{n_x, n_y, n_z} e^{-i \frac{n_x^2 E_0^x + n_y^2 E_0^y + n_z^2 E_0^z}{\hbar} t} s_{n_x}^{L_x}(x) s_{n_y}^{L_y}(y) s_{n_z}^{L_z}(z) \quad (3.96)$$

with

$$A_{n_x, n_y, n_z} = \int_0^{L_x} dx \int_0^{L_y} dy \int_0^{L_z} dz \left( \Psi(0, x, y, z) s_{n_x}^{L_x}(x) s_{n_y}^{L_y}(y) s_{n_z}^{L_z}(z) \right) \quad (3.97)$$

---

<sup>28</sup>The process of obtaining a general solution from the products of particular solutions in the different variables is described in more detail for the 2D membrane in 4.2.

### 3.5 The particle in a 3D box

---

## Chapter 4

# Vibrating membrane

### 4.1 The wave equation

An important differential equation is the previously mentioned *wave equation*, that describes the propagation of light or the propagation of a signal (“vibration”) in a medium<sup>1</sup>

$$\partial_t^2 \Phi(t, \mathbf{x}) = c^2 \nabla^2 \Phi(t, \mathbf{x}) \quad (4.1)$$

#### 4.1.1 A “deduction”

We now present an argument that should convince that eq. (4.1) correctly describes the basic features of the propagation of a vibration or signal in a medium, and that will help in understanding the basic features of wave dynamics.

Let us consider the dynamical system of Fig. 4.1. Here we have  $n + 1$  beads located at positions  $x = 0, a, \dots, ia, \dots, na$ , connected by springs and moving vertically on some frictionless poles. The vertical position of each bead as a function of time is given by  $\Phi_i(t)$ . Assuming all springs to have elastic constant  $k$ , the beads to have mass  $m$ , and springs to have 0 rest length, and recalling the Hooke law relating the recall force to the extension of a spring

$$m \frac{d^2 x(t)}{dt^2} = -kx(t) \quad (4.2)$$

---

<sup>1</sup>The wave equation has been developed initially to describe the propagation of a signal or vibration, and this led 19th century physicists to think about light as the vibration of some kind of medium, called aether. The position of modern physicists is basically the opposite, since nowadays we believe electromagnetic forces to be more fundamental than the vibration of any medium, being the cause of forces between atoms and molecules.

## 4.1 The wave equation

---




Figure 4.1: A system of beads as a model for a vibrating string.

we may write

$$\frac{d^2\Phi_i(t)}{dt^2} = -\frac{k}{m} [(\Phi_i(t) - \Phi_{i-1}(t)) + (\Phi_i(t) - \Phi_{i+1}(t))] \quad (4.3)$$

since the dynamics of  $\Phi_i(t)$  will depend on the displacement with respect to its neighbours. We re-write it as

$$\frac{d^2\Phi_i(t)}{dt^2} = \frac{k}{m} (\Phi_{i+1}(t) - 2\Phi_i(t) + \Phi_{i-1}(t)) \quad (4.4)$$

Now, let us consider our system of beads and springs as the model for a continuous string. We send the distance  $a \rightarrow 0$  and assume we may measure the vertical displacement as a continuous function of  $x$ . Namely, remembering that  $\Phi_i$  corresponds to position  $ia$  we define

$$\Phi_i(t) = \Phi(t, ia) \rightarrow \Phi(t, x) \quad (4.5)$$

Before sending  $a \rightarrow 0$ , we re-write again our equation as

$$\frac{d^2\Phi(t, ia)}{dt^2} = \frac{k}{m} (\Phi(t, (i+1)a) - 2\Phi(t, ia) + \Phi(t, (i-1)a)) \quad (4.6)$$

We now use a Taylor expansion for the term in parentheses

$$\begin{aligned} & \Phi(t, (i+1)a) - 2\Phi(t, ia) + \Phi(t, (i-1)a) = \\ & \Phi(t, ia) + a\partial_x\Phi(t, ia) + \frac{a^2}{2}\partial_x^2\Phi(t, ia) - 2\Phi(t, ia) + \Phi(t, ia) - a\partial_x\Phi(t, ia) + \frac{a^2}{2}\partial_x^2\Phi(t, ia) \\ & + O(a^3) = a^2\partial_x^2\Phi(t, ia) + O(a^3) \end{aligned} \quad (4.7)$$

If we now assume that the following quantity has a finite limit for  $a \rightarrow 0$

$$\lim_{a \rightarrow 0} \frac{ka^2}{m} = c^2 \quad (4.8)$$

we obtain performing the limit

$$\partial_t^2 \Phi(t, x) = c \partial_x^2 \Phi(t, x) \quad (4.9)$$

A simple dimensional analysis shows that the expression in eq. (4.8) is indeed a squared velocity. From eq. (4.2) we have

$$[M][L][T]^{-2} = [k][L] \Rightarrow [k] = [M][T]^{-2} \quad (4.10)$$

and thus

$$[c] = [M][T]^{-2}[L]^2[M]^{-1} = [L]^2[T]^{-2} \quad (4.11)$$

### Physical meaning

It should be very easy to understand why the second derivative emerges on the left side, by comparing the force acting on  $\Phi_2$  and  $\Phi_3$  in Fig. 4.1. In  $\Phi_3$  we have a high (negative) value of  $\partial_x \Phi$ , but this results in a balance in the force acting on the bead. On the other hand, in  $\Phi_2$  we have a change in the derivative, and a resulting force on the bead. It is in the positions in which the string is convex or concave (non zero second derivative) that we have a strong recall function.

Our simple model should also show clearly the nature of the wave (signal, vibration) propagation. We forced our beads not to move in the  $x$  direction, but the interaction between beads causes a signal (the displacement  $\Phi$ ) to propagate in  $x$  and  $t$ .

#### 4.1.2 From string to membrane

It's not hard to generalise to a 2D membrane<sup>2</sup>. We start again with a grid of beads moving on frictionless poles. Now the vertical position of each bead is  $\Phi_{i,j}(t)$ . Each bead is connected to 4 other beads, its first neighbours in  $x$  and  $y$  (Fig. 4.2).

Now we get

$$\begin{aligned} \frac{d^2 \Phi_{i,j}(t)}{dt^2} &= -\frac{k}{m} \left[ (\Phi_{i,j}(t) - \Phi_{i-1,j}(t)) + (\Phi_{i,j}(t) - \Phi_{i+1,j}(t)) \right. \\ &\quad \left. + (\Phi_{i,j}(t) - \Phi_{i,j-1}(t)) + (\Phi_{i,j}(t) - \Phi_{i,j+1}(t)) \right] \\ &= \frac{ka^2}{m} \left[ \frac{\Phi_{i+1,j}(t) - 2\Phi_{i,j}(t) + \Phi_{i-1,j}(t)}{a^2} + \frac{\Phi_{i,j+1}(t) - 2\Phi_{i,j}(t) + \Phi_{i,j-1}(t)}{a^2} \right] \end{aligned} \quad (4.12)$$

---

<sup>2</sup>The proposed models are extremely simplified but they are very easy to handle and describe the basic physics of the problem.

## 4.1 The wave equation

---




Figure 4.2: A system of beads as a model for a vibrating membrane.

and, in the  $a \rightarrow 0$  limit

$$\partial_t^2 \Phi(t, x, y) = c^2 (\partial_x^2 \Phi(t, x, y) + \partial_y^2 \Phi(t, x, y)) \quad (4.13)$$

### 4.1.3 Getting rid of $c$

The velocity of light is the same in any inertial frame. Since nature provided us with such a universal constant, theoretical physicists like to measure their distances in seconds. For example, they would say that 1 meter is the distance that light travels in  $(3 \cdot 10^8)^{-1}$  seconds.

Actually, you will never hear a theoretical physicist saying something like that. They will state that they use a system in which  $c = 1$ . Let us see why. Measuring distances as we just stated is equal to make a change of variables

$$\begin{aligned} x' = \frac{x}{c} &\Rightarrow \partial_x \Phi = \frac{\partial \Phi}{\partial x} = \frac{\partial \Phi}{\partial x'} \frac{\partial x'}{\partial x} = \frac{1}{c} \partial_{x'} \Phi \\ &\Rightarrow \partial_x^2 \Phi = \frac{1}{c} \partial_{x'} \left( \frac{1}{c} \partial_{x'} \Phi \right) \end{aligned} \quad (4.14)$$

so that in the new variables the wave equation becomes

$$\partial_t^2 \Phi(t, x) = \partial_x^2 \Phi(t, x) \quad (4.15)$$

The dimensional terms including  $c$  may be recovered at the end of computations with the same methods used for  $\hbar$  in section 3.4.



Actually, in theoretical physics the change of variables

$$t' = ct, \quad (4.16)$$

(leaving  $\mathbf{x}$  unchanged) is preferred. You may check that this change of variables removes  $c$  from eq. (4.1).

## 4.2 Separation of variables in the vibrating membrane problem

Let us suppose we have a vibrating membrane of size  $L_x$ ,  $L_y$ , with the boundary conditions

$$\Phi(t, 0, y) = \Phi(t, L_x, y) = \Phi(t, x, 0) = \Phi(t, x, L_y) = 0 \quad (4.17)$$

namely the membrane is clamped on the boundary.

We proceed as usual, due to the linearity of the equation and the homogeneous conditions,

$$\Phi(t, x, y) = T(t)X(x)Y(y) \quad (4.18)$$

and obtain

$$c^2 \left( \frac{\partial_x^2 X(x)}{X(x)} + \frac{\partial_y^2 Y(y)}{Y(y)} \right) = K = \frac{\partial_t^2 T(t)}{T(t)} \quad (4.19)$$

Let us focus first on the left side, that we rewrite as

$$\frac{\partial_x^2 X(x)}{X(x)} = \frac{K}{c^2} - \frac{\partial_y^2 Y(y)}{Y(y)} \equiv K_x \quad (4.20)$$

The last step is obtained as usual noticing that we have an equality between functions of unrelated variables. We write also

$$\frac{\partial_y^2 Y(y)}{Y(y)} = \frac{K}{c^2} - K_x \equiv K_y \quad (4.21)$$

We should be familiar enough with these equations to write the conditions for  $K_x$ ,  $K_y$ ,

$$K_x = -\frac{n_x^2 \pi^2}{L_x^2}, \quad K_y = -\frac{n_y^2 \pi^2}{L_y^2}, \quad n_x, n_y \in \mathbb{N} \quad (4.22)$$

to which correspond the solutions

$$X_{n_x}(x) = A_{n_x} s_{n_x}^{L_x}(x), \quad Y_{n_y}(y) = s_{n_y}^{L_y}(y) \quad (4.23)$$

## 4.2 Separation of variables in the vibrating membrane problem

---

The product

$$A_{n_x} A_{n_y} s_{n_x}^{L_x}(x) s_{n_y}^{L_y}(y) \equiv \bar{A}_{n_x, n_y} s_{n_x}^{L_x}(x) s_{n_y}^{L_y}(y) \equiv \bar{A}_{n_x, n_y} s_{n_x, n_y}^{L_x, L_y}(x, y) \quad (4.24)$$

is thus a solution of eq. 4.19 with

$$K = -c^2 \left( \frac{n_x^2 \pi^2}{L_x^2} + \frac{n_y^2 \pi^2}{L_y^2} \right) = -\omega_{n_x, n_y}^2 \quad (4.25)$$

The time equation becomes

$$\partial_t^2 T(t) = -\omega_{n_x, n_y}^2 T(t) \quad (4.26)$$

Its (real) solutions are in the form

$$T_{n_x, n_y}(t) = A \sin(\omega_{n_x, n_y} t) + B \cos(\omega_{n_x, n_y} t) \quad (4.27)$$

By multiplying with eq. (4.24), and re-defining

$$A_{n_x, n_y} = A \bar{A}_{n_x, n_y}, \quad B_{n_x, n_y} = B \bar{B}_{n_x, n_y} \quad (4.28)$$

we obtain the particular solution

$$A_{n_x, n_y} \sin(\omega_{n_x, n_y} t) s_{n_x, n_y}^{L_x, L_y}(x, y) + B_{n_x, n_y} \cos(\omega_{n_x, n_y} t) s_{n_x, n_y}^{L_x, L_y}(x, y) \quad (4.29)$$

where we should remember that  $\omega$  is given by eq. (4.25). The general solution is thus

$$\Phi(t, x, y) = \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} A_{n_x, n_y} \sin(\omega_{n_x, n_y} t) s_{n_x, n_y}^{L_x, L_y}(x, y) + B_{n_x, n_y} \cos(\omega_{n_x, n_y} t) s_{n_x, n_y}^{L_x, L_y}(x, y) \quad (4.30)$$

We have

$$\Phi(0, x, y) = \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} B_{n_x, n_y} s_{n_x, n_y}^{L_x, L_y}(x, y) \quad (4.31)$$

We will assume that this Fourier series may express any physical initial condition  $\Phi_0(x, t)$ . We notice that

$$\begin{aligned} & \int_0^{L_x} dx \int_0^{L_y} dy \left( s_{n_x, n_y}^{L_x, L_y}(x, y) s_{m_x, m_y}^{L_x, L_y}(x, y) \right) = \\ & \left( \int_0^{L_x} s_{n_x}^{L_x}(x) s_{m_x}^{L_x}(x) dx \right) \left( \int_0^{L_y} s_{n_y}^{L_y}(y) s_{m_y}^{L_y}(y) dy \right) \\ & = \delta_{n_x, m_x} \delta_{n_y, m_y} \end{aligned} \quad (4.32)$$

so that

$$B_{n_x, n_y} = \int_0^{L_x} dx \int_0^{L_y} dy \left( \Phi_0(x, t) s_{n_x, n_y}^{L_x, L_y}(x, y) \right) \quad (4.33)$$

These results tell us that the initial condition  $\Phi_0(x, t)$  cannot completely specify the solution, since it provides no information on the constants  $A_{n_x, n_y}$ . Let us then compute

$$\begin{aligned} \partial_t \Phi(t, x, y) = & \\ \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} \omega_{n_x, n_y} & \left[ A_{n_x, n_y} \cos(\omega_{n_x, n_y} t) s_{n_x, n_y}^{L_x, L_y}(x, y) - B_{n_x, n_y} \sin(\omega_{n_x, n_y} t) s_{n_x, n_y}^{L_x, L_y}(x, y) \right] \end{aligned} \quad (4.34)$$

and

$$\partial_t \Phi(0, x, y) = \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} \omega_{n_x, n_y} A_{n_x, n_y} s_{n_x, n_y}^{L_x, L_y}(x, y) \quad (4.35)$$

If now if we know also the initial time derivative  $\partial_t \Phi(0, x, y) = \dot{\Phi}_0(x, t)$  we may obtain the constants  $A_{n_x, n_y}$  through

$$A_{n_x, n_y} = \frac{1}{\omega_{n_x, n_y}} \int_0^{L_x} dx \int_0^{L_y} dy \left( \dot{\Phi}_0(x, t) s_{n_x, n_y}^{L_x, L_y}(x, y) \right) \quad (4.36)$$

and completely specify the solution of our problem.

## 4.2 Separation of variables in the vibrating membrane problem

# Chapter 5

## Laplace equation

### 5.1 Introduction

When we consider the stationary, i.e. time independent, versions of the heat or wave equation, we obtain the Laplace equation

$$\nabla^2\Phi = 0 \tag{5.1}$$

#### 5.1.1 Electrostatics

This equation is extremely important in many physical problems, and in particular in electrostatics, i.e. in the study of electric fields produced by non moving<sup>1</sup> charges.

One of Maxwell equations of Electromagnetism reads as

$$\operatorname{div}\mathbf{E}(t, \mathbf{x}) = k_E \rho(t, \mathbf{x}) \tag{5.2}$$

where  $\mathbf{E}$  is the electric field,  $\rho$  the electric charge, and  $k_E$  is a constant that *depends on the choice of the unit system*.

Furthermore, *for stationary electric and magnetic fields* it is possible to obtain the electric field from a potential<sup>2</sup>

$$\mathbf{E}(\mathbf{x}) = -\nabla\Phi(\mathbf{x}) \tag{5.3}$$

---

<sup>1</sup>Also charges moving at constant velocities can be described using electrostatics, but they generate also a magnetic field.

<sup>2</sup>Maxwell equations relate magnetic and electric fields. Nevertheless, the equations can be separated in equations for the electric field and equations for the magnetic field under stationary conditions, since the terms that couple the two fields involve time derivatives.

## 5.1 Introduction

---

so that, putting the two equations together, we get

$$k_E \rho(\mathbf{x}) = \operatorname{div} \mathbf{E}(\mathbf{x}) = \operatorname{div} (-\nabla \Phi(\mathbf{x})) = -\nabla^2 \Phi(\mathbf{x}) \Rightarrow \nabla^2 \Phi(\mathbf{x}) = -k_E \rho(\mathbf{x}) \quad (5.4)$$

Equation (5.4) is called the Poisson equation, and relates the difference between the average value of  $\Phi$  in a small volume centred in  $\mathbf{x}$  to the amount of charge in  $\mathbf{x}$  (section 1.5.5)

$$\bar{\Phi}(\mathbf{x}) - \Phi(\mathbf{x}) \propto -\rho(\mathbf{x}) \quad (5.5)$$

In absence of charge (for example in vacuum) the Poisson equation turns into the Laplace one, eq. (5.1).

### A “deduction” from Coulomb’s law

The law for the force felt by a charge  $\delta q$  in presence of another charge  $q$  is, for static charges,

$$\mathbf{F} = \frac{k_E \delta q q}{4\pi r^2} \mathbf{e}_r \quad (5.6)$$

Different unit systems use different definitions of  $k_E$ . Theoreticians prefer to define  $k_E$  as a pure number (either 1 or  $4\pi$ ) so that the dimensionality of  $q$  is given by

$$[M][L][T]^{-2} = [q]^2[L]^{-2} \Rightarrow [q] = [M]^{\frac{1}{2}}[L]^{\frac{3}{2}}[T]^{-1} \quad (5.7)$$

while in practical applications a specific dimension for charge is introduced and  $k_E = 1/\epsilon_0$  is used.

We define the electric charge by considering  $\delta q$  small enough not to affect the surrounding charges, and by dividing the force by  $\delta q$  so that

$$\mathbf{E} = k_E \frac{q}{4\pi r^2} \mathbf{e}_r \quad (5.8)$$

Let us compute the flux of  $\mathbf{E}$  on a surface

$$\int_{S_V} \mathbf{E} \cdot \mathbf{n} dS = \int_V \operatorname{div} \mathbf{E} dV \quad (5.9)$$

To compute the divergence it is useful to use eq. (1.74)

$$\operatorname{div} \mathbf{E}(r, \theta, \varphi) = \frac{k_E q}{4\pi} \frac{1}{r^2} \partial_r \left[ r^2 \frac{1}{r^2} \right] = 0 \quad (5.10)$$

This equation is computed using spherical coordinates and placing the origin of the coordinate system on the particle’s position. Nevertheless, since we defined the divergence in a geometrical way, the result is valid regardless

of our choice of the coordinate system. As a result, the flux of the electric field is zero through every surface **that does not include the charged particle**.

The last remark is extremely important, since the spherical coordinate system fails on the origin (zero Jacobian) which is exactly the place where the field diverges, and thus the above computation does not apply there. How can we deal with surfaces that include the charge?

It is very easy to compute the flux on a spherical surface centred on the charge of radius  $R$

$$\int_{S_V} \mathbf{E} \cdot \mathbf{n} dS = k_E \frac{q}{4\pi} \int_0^\pi d\theta \int_0^{2\pi} d\varphi R^2 \frac{1}{R^2} = k_E q \quad (5.11)$$

We may now compute the flux through any surface as the sum of the fluxes through two surfaces: a spherical one centred on the charge, plus the surface that limits a volume given by the original total volume *minus* the ball delimited by the spherical surface. The sum of these two fluxes gives the wanted results, since the terms on the spherical surface get opposite signs (Fig. 5.1).

$$\begin{aligned} \int_{S-\text{ball}} \mathbf{E} \cdot \mathbf{n} dS &= \int_S \mathbf{E} \cdot \mathbf{n} dS + \int_{\text{ball}} \mathbf{E} \cdot (-\mathbf{e}_r) dS \Rightarrow \int_{S-\text{ball}} \mathbf{E} \cdot \mathbf{n} dS + \int_{\text{ball}} \mathbf{E} \cdot \mathbf{n} dS = \\ & \int_S \mathbf{E} \cdot \mathbf{n} dS + \int_{\text{ball}} \mathbf{E} \cdot (\mathbf{e}_r) dS - \int_{\text{ball}} \mathbf{E} \cdot (\mathbf{e}_r) dS = \int_S \mathbf{E} \cdot \mathbf{n} dS \end{aligned} \quad (5.12)$$

We may conclude that the flux through a surface is  $k_E q$  if the surface includes the charge, and 0 if it does not.

An important principle of electrodynamics is that **the field generated by a system of charges is the sum of the fields generated by the single charges**. This, together with the linearity of integration, allows us to write for a system of charges

$$\int_{S_V} \mathbf{E} \cdot \mathbf{n} dS = \sum_{q_i \text{ in } V} k_E q_i \quad (5.13)$$

Let us now assume to have a continuous distribution of charges  $\rho(\mathbf{x})$ , so that

$$\sum_{q_i \text{ in } V} q_i = \int_V \rho(\mathbf{x}) dV \quad (5.14)$$

Using Gauss theorem we get

$$\int_V \text{div} \mathbf{E} dV = \int_{S_V} \mathbf{E} \cdot \mathbf{n} dS = \sum_{q_i \text{ in } V} k_E q_i = \int_V \rho(\mathbf{x}) dV \quad (5.15)$$




Figure 5.1: The flux through a surface as the sum of the flux on the ball centred on the charge, plus the flux on the surface of the volume excluding the ball. The red arrow shows the outgoing vector  $\mathbf{n} = \mathbf{e}_r$  on the ball, while the blue arrow shows the outgoing vector for the surface of the volume excluding the ball,  $\mathbf{n} = -\mathbf{e}_r$ .

Since this result is true regardless of the choice of the volume, we obtain<sup>3</sup>

$$\operatorname{div}\mathbf{E} = \rho(\mathbf{x}) \quad (5.16)$$

The relation

$$\mathbf{E} = -\nabla\Phi \quad (5.17)$$

for a field in the form (5.6) may be verified by computing the gradient of

$$\Phi(\mathbf{x}) = \frac{k_E q}{4\pi r} \quad (5.18)$$

Substitution in eq. (1.42) gives the wanted result<sup>4</sup>.

### 5.1.2 Unicity of solution for the Laplace equation

Eq. (5.5) becomes

$$\bar{\Phi}(\mathbf{x}) = \Phi(\mathbf{x}) \quad (5.19)$$

---

<sup>3</sup>Otherwise, we could integrate on a small volume centred in an area in which the equality does not hold.

<sup>4</sup>Electric fields and potentials for continuous charge distributions may be written by first writing sums over charges and then passing to the integral, as shown in any elementary or advance Electrodynamics text.



in absence of charges, i.e. when Laplace's equation holds. This clearly implies that  $\Phi$  cannot assume a minimum or maximum in a charge free zone<sup>5</sup>.

Let assume that we want to solve the Laplace equation in a volume  $V$ , and that we know the value of  $\Phi$  on the boundary of such region. Then the solution is unique.

To show it, let us assume we have two solutions of the Laplace equation,  $\Phi_1$  and  $\Phi_2$ , assuming the same value on the boundary. The function

$$\tilde{\Phi} \equiv \Phi_1 - \Phi_2 \quad (5.20)$$

is then zero on the boundary, and satisfies Laplace's equation

$$\nabla^2 \tilde{\Phi} = \nabla^2(\Phi_1 - \Phi_2) = \nabla^2 \Phi_1 - \nabla^2 \Phi_2 = 0 \quad (5.21)$$

As a result,  $\tilde{\Phi}$  cannot assume a maximum inside the volume. But since its value is zero on the boundary, we necessarily have

$$\tilde{\Phi} = 0 \Rightarrow \Phi_1 = \Phi_2 \quad (5.22)$$

and accordingly the solution is unique.

This argument can be used also for the Poisson equation, since the difference  $\tilde{\Phi}$  satisfies Laplace equation even in presence of charges,

$$\nabla^2 \tilde{\Phi} = \nabla^2(\Phi_1 - \Phi_2) = \nabla^2 \Phi_1 + \nabla^2 \Phi_2 = k_E \rho - k_E \rho = 0 \quad (5.23)$$

This result is extremely important in physics since it tells us that if we assign a given potential to a set of conductors delimiting a given volume, the value of the potential, and thus of the electric field, will be uniquely determined in the whole volume<sup>6</sup>.

We may now proceed to study some solutions of the Laplace equation, using again the separation of variables method.

## 5.2 2D Laplace equation in a rectangular area

According to the discussion above, providing the value of  $\Phi$  on the border of a rectangle

$$\begin{aligned} \Phi(0, y) &= f(y) \\ \Phi(L_x, y) &= g(y) \\ \Phi(x, 0) &= h(x) \\ \Phi(x, L_y) &= l(x) \end{aligned} \quad (5.24)$$

---

<sup>5</sup>If, for example,  $\Phi$  assumed a maximum in  $\mathbf{x}_0$ , then there would be an area around such point in which  $\Phi(\mathbf{x}) < \Phi(\mathbf{x}_0) \forall \mathbf{x}$ , so that  $\bar{\Phi}(\mathbf{x}) < \Phi(\mathbf{x})$ , leading to a contradiction.

<sup>6</sup>This is true also for a non-finite volume, provided that the behaviour at infinity is given as a further boundary condition, as we will see in some examples.

## 5.2 2D Laplace equation in a rectangular area

---

is given, the potential is uniquely determined inside the area.

This is clearly not a homogeneous boundary condition, but we will nevertheless be able to solve it with the separation of variables, though the definition of four sub-problems.

### 5.2.1 A simpler problem

Let us consider the following simpler problem, i.e. boundary conditions

$$\begin{aligned}\Phi(0, y) &= f(y) \\ \Phi(L_x, y) &= \Phi(x, 0) = \Phi(x, L_y) = 0\end{aligned}\tag{5.25}$$

The conditions are again not homogeneous at  $x = 0$ , but this is not a problem. As we could handle the initial condition at  $t = 0$  through a Fourier series for the heat, wave and Schrödinger equations, we will be able to deal with the  $x = 0$  condition in the Laplace problem.

Let us write

$$\Phi(x, y) = X(x)Y(y)\tag{5.26}$$

The Laplace equation implies

$$\frac{\partial_x^2 X(x)}{X(x)} = C = -\frac{\partial_y^2 Y(y)}{Y(y)}\tag{5.27}$$

We have

$$X(x)Y(0) = X(x)Y(L_y) = 0 \quad \forall x \Rightarrow Y(0) = Y(L_y) = 0\tag{5.28}$$

from which we get, as usual

$$-C = -\frac{n^2 \pi^2}{L_y^2}, \quad n \in \mathbb{N} \Rightarrow C = \frac{n^2 \pi^2}{L_y^2} > 0\tag{5.29}$$

to which corresponds the particular solution

$$Y_n(y) = A_n s_n^{L_y}(y)\tag{5.30}$$

Since  $C$  is positive let us define  $C = \gamma^2$ . We have, regarding the  $X$  equation

$$\partial_x^2 X(x) = \gamma^2 X(x)\tag{5.31}$$

whose general solution is

$$X(x) = A \sinh \gamma x + B \cosh \gamma x\tag{5.32}$$

We want  $X(x)$  to be zero in  $x = L_x$ . It is thus easier to define

$$x' = L_x - x\tag{5.33}$$

so that

$$\partial_{x'}^2 X(x') = \gamma^2 X(x') \quad (5.34)$$

and

$$X(x') = A \sinh \gamma x' + B \cosh \gamma x' \quad (5.35)$$

Now we ask

$$X(x' = 0) = 0 \Rightarrow B = 0 \quad (5.36)$$

and obtain the solution

$$X(x') = A \sinh \gamma x' \quad (5.37)$$

or

$$X(x) = A \sinh(\gamma[L_x - x]) \quad (5.38)$$

The general solution of the Laplace equation that assumes 0 in  $x = L_x$ ,  $y = 0$ ,  $y = L_y$  is thus

$$\Phi(x, y) = \sum_{n=1}^{\infty} A_n \sinh(\gamma[L_x - x]) s_n^{L_y}(y), \quad \gamma = \frac{n\pi}{L_y} \quad (5.39)$$

from which we have

$$\Phi(0, y) = \sum_{n=1}^{\infty} A_n \sinh(\gamma L_x) s_n^{L_y}(y) = f(y) \quad (5.40)$$

and finally

$$A_n = \frac{1}{\sinh(\gamma L_x)} \int_0^{L_y} f(y) s_n^{L_y}(y) \quad (5.41)$$

### 5.2.2 Solution to the full problem

We may use the procedure above to solve the following four problems

$$\nabla^2 \Phi_I = 0, \quad \Phi_I(0, y) = f(y), \quad \Phi_I(L_x, y) = \Phi_I(x, 0) = \Phi_I(x, L_y) = 0 \quad (5.42)$$

$$\nabla^2 \Phi_{II} = 0, \quad \Phi_{II}(L_x, y) = g(y), \quad \Phi_{II}(0, y) = \Phi_{II}(x, 0) = \Phi_{II}(x, L_y) = 0 \quad (5.43)$$

$$\nabla^2 \Phi_{III} = 0, \quad \Phi_{III}(x, 0) = h(x), \quad \Phi_{III}(0, y) = \Phi_{III}(L_x, y) = \Phi_{III}(x, L_y) = 0 \quad (5.44)$$

$$\nabla^2 \Phi_{IV} = 0, \quad \Phi_{IV}(x, L_y) = l(x), \quad \Phi_{IV}(0, y) = \Phi_{IV}(L_x, y) = \Phi_{IV}(x, 0) = 0 \quad (5.45)$$

After we found the four solutions, we have obtained the solution to our problem in the form

$$\Phi = \Phi_I + \Phi_{II} + \Phi_{III} + \Phi_{IV} \quad (5.46)$$

### 5.3 Laplace equation in polar coordinates

---

Indeed

$$\nabla^2\Phi = \nabla^2\Phi_I + \nabla^2\Phi_{II} + \nabla^2\Phi_{III} + \nabla^2\Phi_{IV} = 0 \quad (5.47)$$

and

$$\begin{aligned} \Phi(0, y) &= f(y) + 0 + 0 + 0 = f(y) \\ \Phi(L_x, y) &= 0 + g(y) + 0 + 0 = g(y) \\ \Phi(x, 0) &= 0 + 0 + h(x) + 0 = h(x) \\ \Phi(x, L_y) &= 0 + 0 + 0 + l(x) = l(x) \end{aligned} \quad (5.48)$$

### 5.3 Laplace equation in polar coordinates

Let us assume we want to know the solution of the 2D Laplace inside a circle of radius  $\bar{r}$ , given the boundary condition on the circle

$$\Phi(\bar{r}, \theta) = h(\theta) \quad (5.49)$$

The symmetry of the problem suggests obviously the use of polar coordinates. We recall (eq. 1.82 at fixed  $z$ )

$$\nabla^2\Phi(r, \theta) = \frac{1}{r}\partial_r(r\partial_r\Phi(r, \theta)) + \frac{1}{r^2}\partial_\theta^2\Phi(r, \theta) = 0 \quad (5.50)$$

We try to solve the problem again with the method of separation of variables (it will be clear soon why it works) setting

$$\Phi(r, \theta) = \Theta(\theta)R(r) \quad (5.51)$$

We get

$$\Theta(\theta)\frac{1}{r}\partial_r(r\partial_rR(r)) + \frac{R(r)}{r^2}\partial_\theta^2\Theta(\theta) = 0 \quad (5.52)$$

Dividing by  $\Phi$  we get

$$\frac{1}{rR(r)}\partial_r(r\partial_rR(r)) = -\frac{1}{\Theta(\theta)r^2}\partial_\theta^2\Theta(\theta) \quad (5.53)$$

The variable  $r$  still appears on the right, but if we multiply by  $r^2$  we get

$$\frac{r}{R(r)}\partial_r(r\partial_rR(r)) = C = -\frac{1}{\Theta(\theta)}\partial_\theta^2\Theta(\theta) \quad (5.54)$$

The equation for  $\Theta$  is easier and familiar

$$\partial_\theta^2\Theta(\theta) = -C\Theta(\theta) \quad (5.55)$$

When dealing with the angle  $\theta$ , the boundary conditions are determined by the need to have

$$\Theta(\theta + 2n\pi) = \Theta(\theta) \quad (5.56)$$

We treated this problem in section 2.2. The particular solutions are

$$\Theta_0(\theta) = \bar{A}_0 c_0^p, \quad C = 0 \quad (5.57)$$

and

$$\Theta_n(\theta) = \bar{A}_n c_n^p(\theta) + \bar{B}_n s_n^p(\theta), \quad C = n^2 \quad (5.58)$$

For each value of  $n$ , the radial equation becomes

$$r\partial_r(r\partial_r R_n(r)) = n^2 R_n(r) \quad (5.59)$$

We guess a function in the form  $R_n(r) = A' r^\alpha$ . From

$$\begin{aligned} \partial_r r^\alpha &= \alpha r^{\alpha-1} \\ r\partial_r r^\alpha &= \alpha r^\alpha \\ \partial_r(r\partial_r r^\alpha) &= \alpha^2 r^{\alpha-1} \\ r\partial_r(r\partial_r r^\alpha) &= \alpha^2 r^\alpha \end{aligned} \quad (5.60)$$

we get

$$A' \alpha^2 r^\alpha = A' n^2 r^\alpha \quad (5.61)$$

so that we we have an identity for  $\alpha = \pm n$ . We are dealing with a second order differential equation in  $r$ , and for  $n \neq 0$  we have found a solution depending on two constants

$$R_n(r) = A'_n r^n + B'_n r^{-n} \quad (5.62)$$

For  $n = 0$  we found a single solution,  $R_0 = A'_0$ . Nevertheless, we have

$$r\partial_r \left[ r\partial_r R(r) \right] = 0 \quad (5.63)$$

so that a solution may be found also by asking that the term in square parentheses is constant, so that

$$r\partial_r R(r) = \text{const} = B'_0 \Rightarrow \partial_r R(r) = \frac{B'_0}{r} \Rightarrow R(r) = B'_0 \ln r \quad (5.64)$$

We thus obtain, by redefining the constants (e.g.  $A_n = \bar{A}_n A'_n$ ,  $\tilde{A}_n = \bar{A}_n B'_n$ , etc.) the general solution for the problem as

$$\Phi(r, \theta) = (A_0 + \tilde{A}_0 \ln r) c_0^p + \sum_{n=1}^{\infty} \left( A_n c_n^p(\theta) r^n + \tilde{A}_n c_n^p(\theta) r^{-n} + B_n s_n^p(\theta) r^n + \tilde{B}_n c_n^p(\theta) r^{-n} \right) \quad (5.65)$$

We obtain at  $\bar{r}$

$$\Phi(\bar{r}, \theta) = (A_0 + \tilde{A}_0 \ln \bar{r}) c_0^p + \sum_{n=1}^{\infty} \left( A_n c_n^p(\theta) \bar{r}^n + \tilde{A}_n c_n^p(\theta) \bar{r}^{-n} + B_n s_n^p(\theta) \bar{r}^n + \tilde{B}_n c_n^p(\theta) \bar{r}^{-n} \right) \quad (5.66)$$

### 5.3 Laplace equation in polar coordinates

---

We appear to have a problem, since the general expression for the function  $h(\theta)$  giving the boundary condition at  $\bar{r}$  is

$$h(\theta) = A_0 c_0^p + \sum_{n=1}^{\infty} \left( A_n c_n^p(\theta) + B_n s_n^p(\theta) \right) \quad (5.67)$$

so that the number of constants in eq. (5.66) doubles those of eq. (bcsingle).

This is due to the fact that when we specify the boundary condition for a problem that is defined on an area extending to infinite distance (in opposition to the limited area studied in section 5.2), we need to specify also the behaviour at infinity. Namely, in studying the solution outside the circle, we may ask the potential not to diverge when  $r \rightarrow \infty$ <sup>7</sup>.

Since

$$\lim_{r \rightarrow \infty} \ln r = \infty, \quad \lim_{r \rightarrow \infty} r^n = \infty \quad (5.68)$$

we have

$$\tilde{A}_0 = A_n = B_n = 0 \quad (5.69)$$

The general solution outside the circle reads

$$\Phi(r, \theta) = A_0 c_0^p + \sum_{n=1}^{\infty} \left( \tilde{A}_n c_n^p(\theta) r^{-n} + \tilde{B}_n s_n^p(\theta) r^{-n} \right) \quad (5.70)$$

with

$$A_0 = \int_0^{2\pi} h(\theta) c_0^p d\theta, \quad \tilde{A}_n = \frac{\int_0^{2\pi} h(\theta) c_n^p(\theta) d\theta}{\bar{r}^{-n}}, \quad \tilde{B}_n = \frac{\int_0^{2\pi} h(\theta) s_n^p(\theta) d\theta}{\bar{r}^{-n}} \quad (5.71)$$

If you read carefully, you probably noticed that this is not the solution that we were seeking, because we wanted to find a solution *inside* the circle! But as we discussed before, the polar coordinate system fails at  $r = 0$ . This means that the  $r = 0$  point has to be treated in a special way, since if we do not treat it carefully the solution would be ill defined in the origin.

To find a solution inside the circle, we ask the potential not to diverge in the origin, and since

$$\lim_{r \rightarrow 0} \ln r = -\infty, \quad \lim_{r \rightarrow 0} r^{-n} = \infty \quad (5.72)$$

we have

$$\tilde{A}_0 = \tilde{A}_n = \tilde{B}_n = 0 \quad (5.73)$$

---

<sup>7</sup>This is not the only possible condition. For example, a term growing like  $r$  in the potential can express a field that behaves as constant at infinity. Although a real physical field will never have this property, such a boundary condition may be useful in solving specific problems (e.g., when the field is constant at distances large when compared to the scale at which we study the problem).

The general solution inside the circle reads

$$\Phi(r, \theta) = A_0 c_0^p + \sum_{n=1}^{\infty} \left( A_n c_n^p(\theta) r^n + B_n c_n^p(\theta) r^{-n} \right) \quad (5.74)$$

with

$$A_0 = \int_0^{2\pi} h(\theta) c_0^p d\theta, \quad A_n = \frac{\int_0^{2\pi} h(\theta) c_n^p(\theta) d\theta}{\bar{r}^n}, \quad B_n = \frac{\int_0^{2\pi} h(\theta) s_n^p(\theta) d\theta}{\bar{r}^n} \quad (5.75)$$

A general solution reads as eq. (5.70) for  $r > \bar{r}$ , and eq. (5.70) for  $r < \bar{r}$ . Continuity at  $\bar{r}$  is obtained by construction, since both solutions are equal to  $h(\theta)$  on the boundary.

### 5.3 Laplace equation in polar coordinates

---



## Chapter 6

# A short introduction to function spaces and Dirac formalism

### 6.1 Introduction

We have often written the general solution of our partial differential equation as

$$\Phi(t, x) = \sum_{n=1}^{\infty} A_n f_n(t) s_n(x) \quad (6.1)$$

where the time dependent functions  $f_n(t)$  are uniquely determined by  $n$ . The knowledge of the particular solution implies knowing the  $A_n$ , that we obtain from the initial condition  $\Phi(0, x) = g(x)$  as

$$A_n = \int g(x) s_n(x) dx \quad (6.2)$$

through

$$\int s_m(x) s_n(x) dx = \delta_{n,m} \quad (6.3)$$

This procedure should recall us of another mathematical problem. Let us assume we have a symmetric  $n$  by  $n$  matrix<sup>1</sup>,  $A$ , such that this matrix admits an orthonormal set of eigenvalues  $\mathbf{v}_{\lambda_i}$  which is a basis for the vector

---

<sup>1</sup>A more general class of matrix, normal matrices, would still have the same properties.

## 6.2 Vector spaces

---

space, i.e. vectors such that

$$\begin{aligned} A\mathbf{v}_{\lambda_i} &= \lambda_i\mathbf{v}_{\lambda_i} \\ \mathbf{v}_{\lambda_i} \cdot \mathbf{v}_{\lambda_j} &= \delta_{i,j} \\ \mathbf{a} &= \sum_{i=1}^n \alpha_i \mathbf{v}_{\lambda_i} \quad \forall \mathbf{a} \in \mathbb{R}^n \end{aligned} \tag{6.4}$$

A very straightforward way of computing  $A\mathbf{a}$  is thus

$$A\mathbf{a} = A \sum_{i=1}^n \alpha_i \mathbf{v}_{\lambda_i} = \sum_{i=1}^n \alpha_i A\mathbf{v}_{\lambda_i} = \sum_{i=1}^n \alpha_i \lambda_i \mathbf{v}_{\lambda_i} \tag{6.5}$$

In order to know the coefficients  $\alpha_i$ , we compute

$$\mathbf{v}_{\lambda_j} \cdot \mathbf{a} = \sum_{i=1}^n \alpha_i \mathbf{v}_{\lambda_i} \cdot \mathbf{v}_{\lambda_j} = \sum_{i=1}^n \alpha_i \delta_{i,j} = \alpha_j \tag{6.6}$$

The formal analogy should be clear. Both procedures use indeed the properties of vector spaces provided with a scalar product, the main difference being, as it will be clear soon, that in one case the vector space has finite dimension, while in the other case it has infinite dimension. Since this mathematical structure is extremely powerful, we will study its basic principles in this chapter, learning in the process an extremely useful formalism due to the British physicist and Nobel Prize Paul Dirac<sup>2</sup>.

## 6.2 Vector spaces

### 6.2.1 Basic geometry and physics

Vectors are usually introduced in basic physics through a graphic representation as arrows connecting two points in 3D space. They have magnitude and direction, may be summed through the parallelogram rule, and multiplied by a number, that scales their magnitude without changing their direction if positive, while scales the magnitude and reverses the orientation if negative.

---

<sup>2</sup>Whose many contributions include the development of a formalism from which Heisenberg's and Schrödinger's theories may be derived, an equation to describe the dynamics of a relativistic electron, the theoretical prediction of anti-matter, the statistical theory of many electrons (independently developed also by the Italian Fermi), the seminal works in the fields of Quantum Field Theory and Path Integral formulation of Quantum Mechanics, and the introduction of the Dirac delta function, based on previous works by Fourier and Heaviside, an ill-defined mathematical concept that will be later formalised into distribution theory. A popular book about Dirac's life is entitled "The strangest man".

A scalar product is introduced as an operation that gives a number from two vectors

$$\mathbf{a} \cdot \mathbf{b} = a b \cos \theta \quad (6.7)$$

$a$  and  $b$  being the magnitudes of the vectors and  $\theta$  the angle between them.

Once we choose a Cartesian reference frame, the vector may be represented by its 3 components

$$\mathbf{x} = (x, y, z) = (x_1, x_2, x_3) \quad (6.8)$$

vector sum by

$$\mathbf{x} + \mathbf{y} = (x_1 + y_1, x_2 + y_2, x_3 + y_3) \quad (6.9)$$

product by a scalar by

$$\alpha \mathbf{x} = (\alpha x_1, \alpha x_2, \alpha x_3) \quad (6.10)$$

and scalar product by

$$\mathbf{x} \cdot \mathbf{y} = x_1 y_1 + x_2 y_2 + x_3 y_3 \quad (6.11)$$

### 6.2.2 Generalisation

$\mathbb{R}^n$

We may easily generalise these concepts to the  $n$  dimensional case

$$\mathbf{x} = (x_1, \dots, x_n) \quad (6.12)$$

$$(\mathbf{x} + \mathbf{y})_i = x_i + y_i \quad (6.13)$$

$$(\alpha \mathbf{x})_i = \alpha x_i \quad (6.14)$$

$$\mathbf{x} \cdot \mathbf{y} = \sum_i x_i y_i \quad (6.15)$$

#### Abstract vector space

But we can do better. Let us call a vector space  $V$  a set provided with the operations of sum and a product by a scalar, such that if  $\mathbf{x}, \mathbf{y} \in V$  and  $\alpha$  is a number, then

$$\mathbf{x} + \mathbf{y} = \mathbf{z} \in V, \quad \alpha \mathbf{x} = \mathbf{w} \in V \quad (6.16)$$

We basically call vector space a set on which we may define a sum and product by a number in such a way that the result still belongs to the set. This is clearly related to the studies we have been performing up to now. Consider for example the set of periodic functions,

$$f(\theta + 2n\pi) = f(\theta) \quad (6.17)$$

## 6.2 Vector spaces

---

If we define the sum of two functions  $f$  and  $g$  as the function that assumes in  $\theta$  the value

$$(f + g)(\theta) = f(\theta) + g(\theta) \quad (6.18)$$

and as the product of  $f$  with the number  $\alpha$  the function that assumes in  $\theta$  the value

$$(\alpha f)(\theta) = \alpha f(\theta) \quad (6.19)$$

we may trivially verify that the resulting functions are still members of the set. Other examples of interest are functions that assume zero value on boundaries, functions that have zero derivative on the boundaries, and solutions of linear differential equations<sup>3</sup>.

### Complex numbers

Until now we have been talking of a product with “a number”  $\alpha$ . This vague concept was used to hint that  $\alpha \in \mathbb{R}$ .<sup>4</sup> But we may actually be interested to generalise to  $\alpha \in \mathbb{C}$ .<sup>5</sup>

The main reason for using complex numbers is the *Fundamental theorem of algebra*, that states that any polynomial of degree  $n > 0$  with complex coefficients has at least a complex root (and actually, if counted with multiplicity, exactly  $n$  roots). Namely, when we are dealing with complex polynomial equations, we may always find a solution, and actually all the solutions we need. Two examples will help us in understanding better this issue.

Let us consider the second order differential equation

$$\frac{d^2 f}{dx^2} = a \frac{df}{dx} + bf \quad (6.20)$$

If we guess a solution in the form  $e^{\lambda x}$  we get

$$(\lambda^2 - a\lambda - b)e^{\lambda x} = 0 \quad (6.21)$$

A solution is then found by solving the algebraic equation

$$\lambda^2 - a\lambda - b = 0 \quad (6.22)$$

The equation above does not necessarily have real solutions, while it has exactly two complex solutions, providing the correct number of integration

---

<sup>3</sup>Examples of sets that are not vector spaces are functions assuming non zero value on boundaries, or solutions of non-linear differential equations.

<sup>4</sup>In such a case we say that  $V$  is a vector space on the real field.

<sup>5</sup>In such a case we say that  $V$  is a vector space on the complex field.

constants for a second order equation. For example, the most important version of this equation is probably the harmonic oscillator one,  $a = 0$ ,  $b = -1$ , whose solutions are complex,  $\lambda = \pm i$ . By working with complex numbers we may obtain the solutions

$$e^{\pm ix} \tag{6.23}$$

and from their linear combination the real solutions  $\sin x$  and  $\cos x$ .

The reader should be already familiar with the concept of eigenvectors and eigenvalues of a matrix (that will be reviewed soon and related to the problem of solving differential equations). Let us consider the simple matrix (rotation of  $\pi/2$ )

$$A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \tag{6.24}$$

To find its eigenvalues, we compute

$$\det(A - \lambda \mathbb{1}) = \lambda^2 + 1 = 0 \tag{6.25}$$

There is no real solution, as understandable since no real vector is left unchanged by such a rotation. Nevertheless, as we have seen above (eq. 6.4), it is useful to operate on eigenvectors of the matrix, so it is important to know that to the complex eigenvalues  $\pm i$  correspond the eigenvectors

$$\mathbf{v}_i = \begin{pmatrix} \alpha \\ i\alpha \end{pmatrix} \quad \mathbf{v}_{-i} = \begin{pmatrix} \alpha \\ -i\alpha \end{pmatrix} \tag{6.26}$$

### 6.2.3 The scalar or inner product

A scalar or inner product on a Vector space is an operator that takes two vectors and gives a complex number

$$\langle \mathbf{x}, \mathbf{y} \rangle \in \mathbb{C} \tag{6.27}$$

with the following properties

1. Complex conjugation under exchange:

$$\langle \mathbf{x}, \mathbf{y} \rangle = (\langle \mathbf{y}, \mathbf{x} \rangle)^* \tag{6.28}$$

2. Positive definiteness:

$$\langle \mathbf{x}, \mathbf{x} \rangle \geq 0, \quad \langle \mathbf{x}, \mathbf{x} \rangle = 0 \text{ iff } \mathbf{x} = 0 \tag{6.29}$$

## 6.2 Vector spaces

---

3. Linearity:

$$\langle \mathbf{x}, \alpha \mathbf{y} \rangle = \alpha \langle \mathbf{x}, \mathbf{y} \rangle, \quad \langle \mathbf{x}, \mathbf{y} + \mathbf{z} \rangle = \langle \mathbf{x}, \mathbf{y} \rangle + \langle \mathbf{x}, \mathbf{z} \rangle \quad (6.30)$$

The definition eq. (6.15) satisfies the above conditions *when applied to real vectors*, but needs to be modified to

$$\mathbf{x} \cdot \mathbf{y} = \sum_i x_i^* y_i \quad (6.31)$$

to satisfy them in the complex case. One may wonder then why we asked the scalar product to satisfy eq. (6.28) in the first place. First of all, such a property automatically assures

$$\langle \mathbf{x}, \mathbf{x} \rangle \in \mathbb{R} \quad (6.32)$$

Furthermore, coupled with linearity, we have

$$\langle \alpha \mathbf{x}, \mathbf{y} \rangle = (\langle \mathbf{y}, \alpha \mathbf{x} \rangle)^* = (\alpha \langle \mathbf{y}, \mathbf{x} \rangle)^* = \alpha^* (\langle \mathbf{y}, \mathbf{x} \rangle)^* = \alpha^* \langle \mathbf{x}, \mathbf{y} \rangle \quad (6.33)$$

so that

$$\langle \alpha \mathbf{x}, \alpha \mathbf{x} \rangle = \alpha^* \alpha \langle \mathbf{x}, \mathbf{x} \rangle = |\alpha|^2 \langle \mathbf{x}, \mathbf{x} \rangle \quad (6.34)$$

so that the required property eq. (6.28) allows us to define a positive definite scalar product. Indeed, for the specific definition eq. (6.31) we have

$$\mathbf{x} \cdot \mathbf{x} = \sum_i |x_i|^2 \geq 0 \quad (6.35)$$

A positive definite scalar product allows us to use the scalar product to define a norm

$$\|\mathbf{x}\| = \sqrt{\langle \mathbf{x}, \mathbf{x} \rangle} \quad (6.36)$$

### 6.2.4 Linear independence, bases, dimension and the like

#### Linear independence

We say that a set of vectors of  $n$  vectors  $\mathbf{v}_i \in V$  is linearly independent if

$$\sum_i c_i \mathbf{v}_i = 0 \Rightarrow c_i = 0 \forall i \quad (6.37)$$

(or, equivalently, if it is impossible to write one of them as a combination of some of the others).

### Basis and dimension

Linear combinations of these vectors forms a subspace of  $V$ , i.e. a vector space  $V' \subset V$ . If  $V' = V$ , i.e. if any vector in  $V$  may be written as a linear combination of the  $\mathbf{v}_i$ , we say that the vectors form a *basis* of the vector space, and that the *dimension* of the vector space is  $n$ . It may indeed be shown that if the vector space has a basis of size  $n$ , all its bases will necessarily have the same size<sup>6</sup>.

Obviously, not all vector spaces admit a basis of dimension  $n$ . If they do, we say that the space is finite dimensional. Finite dimension spaces are “easier” and we will focus on them for a while.

#### 6.2.5 Orthonormalisation

If a (non-zero) vector  $\mathbf{v}_1$  is given, it is trivial to obtain a *normalised* vector from it, i.e.

$$\mathbf{e}_1 = \frac{1}{\sqrt{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle}} \mathbf{v}_1 \quad (6.38)$$

so that

$$\langle \mathbf{e}_1, \mathbf{e}_1 \rangle = \frac{1}{\sqrt{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle}} \langle \mathbf{v}_1, \mathbf{v}_1 \rangle = 1 \quad (6.39)$$

Let us now consider a second vector  $\mathbf{v}_2$ , and assume  $\mathbf{v}_1$  and  $\mathbf{v}_2$  to be linearly independent. We now compute

$$\mathbf{v}'_2 = \mathbf{v}_2 - \langle \mathbf{e}_1, \mathbf{v}_2 \rangle \mathbf{e}_1 \neq 0 \quad (6.40)$$

The result is necessarily non-zero, otherwise it would mean that  $\mathbf{v}_2$  may be written as a linear combination of  $\mathbf{v}_1$ . We may now compute

$$\langle \mathbf{e}_1, \mathbf{v}'_2 \rangle = \langle \mathbf{e}_1, \mathbf{v}_2 \rangle - \langle \mathbf{e}_1, \mathbf{v}_2 \rangle \langle \mathbf{e}_1, \mathbf{e}_1 \rangle = \langle \mathbf{e}_1, \mathbf{v}_2 \rangle - \langle \mathbf{e}_1, \mathbf{v}_2 \rangle = 0 \quad (6.41)$$

We may now define

$$\mathbf{e}_2 = \frac{1}{\sqrt{\langle \mathbf{v}'_2, \mathbf{v}'_2 \rangle}} \mathbf{v}'_2 \quad (6.42)$$

so that

$$\langle \mathbf{e}_1, \mathbf{e}_1 \rangle = \langle \mathbf{e}_2, \mathbf{e}_2 \rangle = 1, \quad \langle \mathbf{e}_1, \mathbf{e}_2 \rangle = 0 \quad (6.43)$$

---

<sup>6</sup>The logic behind the proof is the following. Assume you have a basis  $\mathbf{x}_i$  of size  $n$ , and a basis  $\mathbf{y}_i$  of size  $m > n$ . You can write  $\mathbf{y}_1$  as a combination of the  $\mathbf{x}_i$ , and use the obtained formula to express one of the  $\mathbf{x}_i$ , let us call it  $\mathbf{x}_1$  (a re-ordering may be necessary), as a combination of  $\mathbf{y}_1, \mathbf{x}_i, i > 1$ . This set of  $n$  vectors is thus a basis, that we use to express  $\mathbf{y}_2$ . By repeating the process, you finally obtain that the first  $n$   $\mathbf{y}_i$  are a basis, in contradiction with the hypothesis.

---

## 6.2 Vector spaces

---

If now a third vector  $\mathbf{v}_3$ , forming a linearly independent set with  $\mathbf{v}_1$  and  $\mathbf{v}_2$ , is given, we define

$$\mathbf{v}'_3 = \mathbf{v}_3 - \langle \mathbf{e}_1, \mathbf{v}_3 \rangle \mathbf{e}_1 - \langle \mathbf{e}_2, \mathbf{v}_3 \rangle \mathbf{e}_2 \neq 0 \quad (6.44)$$

This vector is again non-zero, since  $\mathbf{e}_1$  and  $\mathbf{e}_2$  are linear combinations of  $\mathbf{v}_1$  and  $\mathbf{v}_2$ , and thus if we had a zero we could write

$$\mathbf{v}_3 = \langle \mathbf{e}_1, \mathbf{v}_3 \rangle \mathbf{e}_1 - \langle \mathbf{e}_2, \mathbf{v}_3 \rangle \mathbf{e}_2 = c_1 \mathbf{v}_1 + c_2 \mathbf{v}_2 \quad (6.45)$$

in contradiction with the hypothesis. Now we have

$$\begin{aligned} \langle \mathbf{e}_1, \mathbf{v}'_3 \rangle &= \langle \mathbf{e}_1, \mathbf{v}_3 \rangle - \langle \mathbf{e}_1, \mathbf{v}_3 \rangle \langle \mathbf{e}_1, \mathbf{e}_1 \rangle - \langle \mathbf{e}_2, \mathbf{v}_3 \rangle \langle \mathbf{e}_1, \mathbf{e}_2 \rangle = \\ &\langle \mathbf{e}_1, \mathbf{v}_3 \rangle - \langle \mathbf{e}_1, \mathbf{v}_3 \rangle - 0 = 0 \end{aligned} \quad (6.46)$$

and

$$\langle \mathbf{e}_2, \mathbf{v}'_3 \rangle = \langle \mathbf{e}_2, \mathbf{v}_3 \rangle - \langle \mathbf{e}_1, \mathbf{v}_3 \rangle \langle \mathbf{e}_2, \mathbf{e}_1 \rangle - \langle \mathbf{e}_2, \mathbf{v}_3 \rangle \langle \mathbf{e}_2, \mathbf{e}_2 \rangle = 0 \quad (6.47)$$

By defining

$$\mathbf{e}_3 = \frac{1}{\sqrt{\langle \mathbf{v}'_3, \mathbf{v}'_3 \rangle}} \mathbf{v}'_3 \quad (6.48)$$

we get

$$\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \delta_{i,j}, \quad i, j = 1, 2, 3 \quad (6.49)$$

The generalisation

$$\mathbf{v}'_k = \mathbf{v}_k - \sum_{i=1}^{k-1} \langle \mathbf{e}_i, \mathbf{v}_k \rangle \mathbf{e}_i \neq 0 \quad (6.50)$$

$$\mathbf{e}_k = \frac{1}{\sqrt{\langle \mathbf{v}'_k, \mathbf{v}'_k \rangle}} \mathbf{v}'_k \quad (6.51)$$

leads clearly to the construction of a orthonormal basis

$$\langle \mathbf{e}_i, \mathbf{e}_j \rangle = \delta_{i,j} \quad (6.52)$$

starting from any given basis.

### 6.2.6 Scalar product and orthonormal bases

Given vectors  $\mathbf{x}, \mathbf{y}$  in a finite dimensional vector space  $V$ , we have, as shown above,

$$\mathbf{x} = \sum_i x_i \mathbf{e}_i, \quad \mathbf{y} = \sum_i y_i \mathbf{e}_i \quad (6.53)$$



where the  $x_i$  and  $y_i$  are the coefficients that give the vectors as linear combination of the vectors in the basis (the components in that basis), and the basis has been chosen to be orthonormal with respect to the scalar product of the vector space.

We then have

$$\begin{aligned} \langle \mathbf{x}, \mathbf{y} \rangle &= \left\langle \sum_i x_i \mathbf{e}_i, \sum_j y_j \mathbf{e}_j \right\rangle = \sum_j y_j \left( \left\langle \sum_i x_i \mathbf{e}_i, \mathbf{e}_j \right\rangle \right) = \sum_j y_j \left( \sum_i \langle x_i \mathbf{e}_i, \mathbf{e}_j \rangle \right) = \\ &= \sum_j y_j \left( \sum_i x_i^* \langle \mathbf{e}_i, \mathbf{e}_j \rangle \right) = \sum_j y_j \left( \sum_i x_i^* \delta_{i,j} \right) = \sum_j x_j^* y_j \end{aligned} \quad (6.54)$$

We thus see that once we assign a orthonormal basis to a vector space of dimension  $n$ , we may establish (eq. 6.53) a one-to-one correspondence with the space  $\mathbb{C}^n$ , that preserves (eq. 6.54) the form of the scalar product eq. (6.31).

We may also show that if we have, in a space of dimension  $n$ ,  $n$  orthonormal vectors, they are independent, and thus a base. Indeed, if we assume

$$\mathbf{e}_i = \sum_{j \neq i} c_j \mathbf{e}_j \quad (6.55)$$

we get the contradiction that all  $c_j$  are zero

$$0 = \langle \mathbf{e}_k, \mathbf{e}_i \rangle = \sum_{j \neq i} c_j \langle \mathbf{e}_k, \mathbf{e}_j \rangle = c_k \quad (6.56)$$

## 6.3 Linear operators

### 6.3.1 Definition

Let us consider an *linear operator*  $A$  from  $V$  to  $V$ , i.e. an operator such that, if  $\mathbf{x}, \mathbf{v}$  are in  $V$  then we have

$$A\mathbf{x} = \mathbf{z} \in V \quad (6.57)$$

$$A(\alpha\mathbf{x}) = \alpha A\mathbf{x} \quad (6.58)$$

$$A(\mathbf{x} + \mathbf{y}) = A\mathbf{x} + A\mathbf{y} \quad (6.59)$$

## 6.3 Linear operators

---

### 6.3.2 Linear operators on a finite dimensional space

Let us consider a finite dimensional space with orthonormal basis  $\mathbf{e}_i$ . When a linear operator  $A$  acts on a member of the basis  $\mathbf{e}_i$ , it produces a vector in  $V$ , that may again be written as a linear combination of the members of the basis

$$A\mathbf{e}_i = \mathbf{z} = \sum_{j=1}^n z_j \mathbf{e}_j \quad (6.60)$$

Now, since the  $z_j$  are determined by  $A$  and  $\mathbf{e}_i$  we may call them  $A_{ji}$ . Let us now have  $A$  to act on a generic vector  $\mathbf{x}$

$$\begin{aligned} A\mathbf{x} &= A\left(\sum_i x_i \mathbf{e}_i\right) = \sum_i x_i A\mathbf{e}_i = \sum_i x_i \sum_j A_{ji} \mathbf{e}_j = \sum_j \left(\sum_i A_{ji} x_i\right) \mathbf{e}_j \\ &= \mathbf{z} = \sum_j z_j \mathbf{e}_j \end{aligned} \quad (6.61)$$

The action of the linear operator is thus to take a vector with components  $x_i$  and to generate a vector with components  $\sum_i A_{ji} x_i$ . We have thus created a correspondence between linear operators and matrices acting on  $\mathbb{R}^n$ .

What is exactly  $A_{ji}$ ? We can see it easily by computing

$$\langle \mathbf{e}_j, A\mathbf{e}_i \rangle = \left\langle \mathbf{e}_j, \sum_k A_{ki} \mathbf{e}_k \right\rangle = \sum_k A_{ki} \langle \mathbf{e}_j, \mathbf{e}_k \rangle = \sum_k A_{ki} \delta_{jk} = A_{ji} \quad (6.62)$$

What happens when a linear operator acts on the result of another linear operator?

$$\begin{aligned} BA\mathbf{x} &= B(A\mathbf{x}) = B\left(\sum_j \left(\sum_i A_{ji} x_i\right) \mathbf{e}_j\right) = \sum_j B\left(\left(\sum_i A_{ji} x_i\right) \mathbf{e}_j\right) = \\ &= \sum_j \left(\sum_i A_{ji} x_i (B\mathbf{e}_j)\right) = \sum_j \left(\sum_i A_{ji} x_i \left(\sum_k B_{kj} \mathbf{e}_k\right)\right) = \sum_k \left(\sum_j \sum_i B_{kj} A_{ji} x_i\right) \mathbf{e}_k \end{aligned} \quad (6.63)$$

The components of the new vector are given by  $\sum_j \sum_k B_{kj} A_{ji} x_i$  so that also the rule of matrix product is obtained for operators on general vector spaces. This means that

$$\langle \mathbf{e}_k, BA\mathbf{e}_i \rangle = \sum_j \langle \mathbf{e}_k, B\mathbf{e}_j \rangle \langle \mathbf{e}_j, A\mathbf{e}_i \rangle \quad (6.64)$$

a formula that will result trivial once we introduce the Dirac formalism.

### 6.3.3 Eigenvalues and eigenvectors

In general, the action of a linear operator on a vector is more complex than a simple multiplication by a complex number

$$A\mathbf{x} \neq \alpha\mathbf{x} \quad (6.65)$$

and indeed an operator for which the above equation would hold *for all*  $\mathbf{x}$  would be just a multiple of the identity operator  $\mathbf{1}$ .

Nevertheless, for each operator  $A$ , there may be some *eigenvalues*  $\lambda$  and *eigenvectors*  $\mathbf{v}_\lambda$  such that

$$A\mathbf{v}_\lambda = \lambda\mathbf{v}_\lambda \quad (6.66)$$

The knowledge of such eigenvalues and eigenvectors is extremely important, as already discussed above (eq. 6.6). In order to explicitly compute the eigenvalues and eigenvectors, the matrix representation of linear operators is extremely valuable. Let us first rewrite eq. (6.66) as

$$(A - \lambda\mathbf{1})\mathbf{v}_\lambda = 0 \quad (6.67)$$

Using an explicit basis, we get

$$\sum_j (A - \lambda\mathbf{1})_{i,j} (\mathbf{v}_\lambda)_j = 0 \quad (6.68)$$

From the theory of linear equations, we know that this is possible only if

$$\det(A - \lambda\mathbf{1})_{i,j} = \det(A_{i,j} - \lambda\delta_{i,j}) = 0 \quad (6.69)$$

The above equation is a polynomial of order  $n$  in  $\lambda$ , and its solution gives (since we are working with complex numbers)  $n$  roots  $\lambda_i$ , (possibly some of them being equal, i.e. of multiplicity larger than one). The solution of the linear equations

$$\sum_j A_{k,j} (\mathbf{v}_{\lambda_i})_j = \lambda_i (\mathbf{v}_{\lambda_i})_k \quad (6.70)$$

will provide the explicit form of the eigenvector  $\mathbf{v}_{\lambda_i}$ . Obviously, the solution of eq. (6.70) is *not* unique. We know this both from the theory of linear equations (since  $\det(A - \lambda_i\mathbf{1}) = 0$ ) and from general theory of linear operators

$$A\mathbf{v} = \lambda\mathbf{v} \Rightarrow A(\alpha\mathbf{v}) = \alpha A\mathbf{v} = \lambda(\alpha\mathbf{v}) \quad (6.71)$$

so that if  $\mathbf{v}$  is an eigenvector, also  $\alpha\mathbf{v}$  is an eigenvector with the same eigenvalue.

## 6.3 Linear operators

---

### 6.3.4 Inverse operator

Let us consider an operator  $A$  such that each  $\mathbf{y}$  in  $V$  may be written as

$$\mathbf{y} = A\mathbf{x} \quad (6.72)$$

We define the operator  $A^{-1}$  as the operator

$$A^{-1}A = \mathbb{1} \Rightarrow A^{-1}\mathbf{y} = \mathbf{x} \quad (6.73)$$

We may easily verify that

$$AA^{-1} = \mathbb{1} \quad (6.74)$$

Indeed

$$AA^{-1}\mathbf{y} = AA^{-1}(A\mathbf{x}) = A(A^{-1}A\mathbf{x}) = A\mathbf{x} = \mathbf{y} \quad (6.75)$$

In matrix representation we have

$$\delta_{i,j} = \langle \mathbf{e}_i, \mathbb{1}\mathbf{e}_j \rangle = \langle \mathbf{e}_i, A^{-1}A\mathbf{e}_j \rangle = \sum_k A_{i,k}^{-1}A_{k,j} \quad (6.76)$$

### 6.3.5 Adjoint operator

#### Definition

Given the linear operator  $A$ , we define its adjoint  $A^\dagger$  by asking

$$\langle \mathbf{x}, A\mathbf{y} \rangle = \langle A^\dagger\mathbf{x}, \mathbf{y} \rangle, \quad \forall \mathbf{x}, \mathbf{y} \quad (6.77)$$

Let us first notice that, once we assign an orthonormal basis  $\mathbf{e}_i$  we have

$$\begin{aligned} \langle \mathbf{x}, A\mathbf{y} \rangle &= \sum_i x_i^* \langle \mathbf{e}_i, A\mathbf{y} \rangle = \sum_i x_i^* \left\langle \mathbf{e}_i, \sum_j \sum_k A_{j,k} y_k \mathbf{e}_j \right\rangle \\ &= \sum_i \sum_j \sum_k x_j^* A_{j,k} y_k \langle \mathbf{e}_i, \mathbf{e}_j \rangle = \sum_i \sum_j \sum_k x_j^* A_{j,k} y_k \delta_{i,j} = \sum_j \sum_k x_j^* A_{j,k} y_k \end{aligned} \quad (6.78)$$

and

$$\langle A^\dagger\mathbf{x}, \mathbf{y} \rangle = \left( \langle \mathbf{y}, A^\dagger\mathbf{x} \rangle \right)^* = \left( \sum_j \sum_k y_k^* A_{k,j}^\dagger x_j \right)^* = \sum_j \sum_k x_j^* \left( A_{k,j}^\dagger \right)^* y_k \quad (6.79)$$

so that, since the relation has to be true for each  $\mathbf{x}$  and  $\mathbf{y}$ , the adjoint operator's matrix representation is given by the the complex conjugate of the transpose

$$\left( A_{k,j}^\dagger \right)^* = A_{j,k} \Rightarrow A_{k,j}^\dagger = (A_{j,k})^* \quad (6.80)$$

We may easily verify that

$$(AB)^\dagger = B^\dagger A^\dagger \quad (6.81)$$

Indeed

$$\langle \mathbf{x}, AB\mathbf{y} \rangle = \langle A^\dagger \mathbf{x}, B\mathbf{y} \rangle = \langle B^\dagger A^\dagger \mathbf{x}, \mathbf{y} \rangle \quad (6.82)$$

### Self adjoint operators

We say that an operator is self-adjoint if, for any  $\mathbf{x}, \mathbf{y}$ , the following holds

$$\langle \mathbf{x}, A\mathbf{y} \rangle = \langle A\mathbf{x}, \mathbf{y} \rangle \quad (6.83)$$

Namely the operator is self-adjoint if we can move it on the other side of the scalar product without changing it.

According to the theory developed above, a linear operator on a finite dimensional vector space is self-adjoint if its matrix representation satisfies

$$A_{k,j} = (A_{j,k})^* \quad (6.84)$$

### Eigenvalues and eigenvectors of a self-adjoint operator

Self-adjoint operators (symmetric operators in the real case) have properties that make them extremely important. First of all, their eigenvalues  $\lambda$  such that

$$A\mathbf{v}_\lambda = \lambda\mathbf{v}_\lambda \quad (6.85)$$

are real,  $\lambda \in \mathbb{R}$ . Indeed

$$\begin{aligned} \lambda \langle \mathbf{v}, \mathbf{v} \rangle &= \langle \mathbf{v}, \lambda\mathbf{v} \rangle = \langle \mathbf{v}, A\mathbf{v} \rangle = \langle A^\dagger \mathbf{v}, \mathbf{v} \rangle = \langle A\mathbf{v}, \mathbf{v} \rangle = \langle \lambda\mathbf{v}, \mathbf{v} \rangle = \\ &\lambda^* \langle \mathbf{v}, \mathbf{v} \rangle \Rightarrow \lambda = \lambda^* \end{aligned} \quad (6.86)$$

Furthermore, two eigenvalues corresponding two different eigenvectors of a self-adjoint operator are orthogonal. Indeed if

$$A\mathbf{v}_{\lambda_1} = \lambda_1\mathbf{v}_{\lambda_1}, \quad A\mathbf{v}_{\lambda_2} = \lambda_2\mathbf{v}_{\lambda_2}, \quad \lambda_1 \neq \lambda_2 \quad (6.87)$$

we have, using  $A = A^\dagger$  and  $\lambda = \lambda^*$ ,

$$\begin{aligned} \lambda_1 \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle &= \langle A\mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle = \langle \mathbf{v}_{\lambda_1}, A\mathbf{v}_{\lambda_2} \rangle = \lambda_2 \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle \\ &\Rightarrow (\lambda_1 - \lambda_2) \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle \Rightarrow \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle = 0 \end{aligned} \quad (6.88)$$

This is not enough to assure that  $A$  has  $n$  orthogonal eigenvectors, since the multiplicity of some eigenvalues may be higher than one (the sub-space corresponding to some eigenvalue  $\lambda$  is more than one.). Nevertheless, it is

### 6.3 Linear operators

---

possible to show that a self-adjoint  $A$  has an orthonormal basis of eigenvectors.

We can show it using the principle of induction. For  $n = 1$  it is clear that any operator (simply the multiplication by a number) has any vector as an eigenvector, and such vector may be normalised and be, by itself, a basis for the vector space. Let us assume that on any space of dimension  $n - 1$  a self-adjoint operator  $A$  admits an orthonormal basis. For a  $n$  dimensional space, we start by finding an eigenvector  $\mathbf{v}_1$ , since we may always do that, as shown above, due to the fundamental theorem of algebra. Let us call the corresponding eigenvalue  $\lambda_1$ . This vector may be normalised as

$$\mathbf{e}_1 = \frac{1}{\sqrt{\langle \mathbf{v}_1, \mathbf{v}_1 \rangle}} \mathbf{v}_1 \quad (6.89)$$

We may always now find  $n - 1$  vectors  $\mathbf{v}_i$ ,  $i > 1$  that form a basis together with  $\mathbf{e}_1$ . We may orthonormalise them in the basis  $\{\mathbf{e}_1, \bar{\mathbf{e}}_i\}$ . The linear combinations of the  $\bar{\mathbf{e}}_i$  form obviously a subspace of vectors orthogonal to  $\mathbf{e}_1$ . The important point, in which the fact that  $A$  is self-adjoint is crucial, is that the action of  $A$  on vectors in the subspace, stays in the subspace, since  $A\bar{\mathbf{e}}_i$  is orthogonal to  $\mathbf{e}_1$

$$\langle A\bar{\mathbf{e}}_i, \mathbf{e}_1 \rangle = \langle \bar{\mathbf{e}}_i, A\mathbf{e}_1 \rangle = \lambda_1 \langle \bar{\mathbf{e}}_i, \mathbf{e}_1 \rangle = 0 \quad (6.90)$$

Now we can use the induction hypothesis, and find  $n - 1$  orthonormal eigenvectors on the  $n - 1$  space orthogonal to  $\mathbf{e}_1$ , that we may call  $\mathbf{e}_i$  and that, being in a subspace orthogonal to  $\mathbf{e}_1$ , form with it an orthonormal basis.

Symmetric real matrices

$$A_{i,j} = A_{j,i} \quad (6.91)$$

are clearly self-adjoint and thus admit a orthonormal basis of eigenvectors with real eigenvalues.

#### Anti-adjoint operator

It is interesting to check what happens to an anti-adjoint operator

$$A^\dagger = -A \quad (6.92)$$

Their eigenvalues  $\lambda$  such that

$$A\mathbf{v}_\lambda = \lambda\mathbf{v}_\lambda \quad (6.93)$$

are *purely imaginary*. Indeed

$$\begin{aligned} \lambda \langle \mathbf{v}, \mathbf{v} \rangle &= \langle \mathbf{v}, \lambda \mathbf{v} \rangle = \langle \mathbf{v}, A\mathbf{v} \rangle = \langle A^\dagger \mathbf{v}, \mathbf{v} \rangle = \langle -A\mathbf{v}, \mathbf{v} \rangle = \langle -\lambda \mathbf{v}, \mathbf{v} \rangle = \\ &= -\lambda^* \langle \mathbf{v}, \mathbf{v} \rangle \Rightarrow \lambda = -\lambda^* \end{aligned} \quad (6.94)$$

Furthermore, two eigenvalues corresponding two different eigenvectors are orthogonal. Indeed if

$$A\mathbf{v}_{\lambda_1} = \lambda_1 \mathbf{v}_{\lambda_1}, \quad A\mathbf{v}_{\lambda_2} = \lambda_2 \mathbf{v}_{\lambda_2}, \quad \lambda_1 \neq \lambda_2 \quad (6.95)$$

we have, using  $A^\dagger = -A$  and  $\lambda^* = -\lambda$ ,

$$\begin{aligned} \lambda_1 \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle &= \langle -\lambda_1 \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle = \langle -A\mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle = \langle \mathbf{v}_{\lambda_1}, A\mathbf{v}_{\lambda_2} \rangle = \lambda_2 \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle \\ &\Rightarrow (\lambda_1 - \lambda_2) \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle \Rightarrow \langle \mathbf{v}_{\lambda_1}, \mathbf{v}_{\lambda_2} \rangle = 0 \end{aligned} \quad (6.96)$$

To show that these operators admit a basis of orthonormal eigenvectors, we just need to repeat the proof above using this time

$$\langle A\bar{\mathbf{e}}_i, \mathbf{e}_1 \rangle = \langle \bar{\mathbf{e}}_i, -A\mathbf{e}_1 \rangle = -\lambda_1 \langle \bar{\mathbf{e}}_i, \mathbf{e}_1 \rangle = 0 \quad (6.97)$$

Purely imaginary symmetric matrices are anti-adjoint, so that they share these properties.

### 6.3.6 Commuting operators

In general, linear operators on a vector space *do not commute*. Let us consider for example a  $n = 2$  space with an orthonormal basis  $\mathbf{e}_1, \mathbf{e}_2$ . Let us consider the operator  $A$  such that

$$A\mathbf{e}_1 = \mathbf{e}_2, \quad A\mathbf{e}_2 = 0 \quad (6.98)$$

whose matrix representation is

$$A = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} \quad (6.99)$$

and the operator  $B$  such that

$$B\mathbf{e}_1 = 0, \quad B\mathbf{e}_2 = \mathbf{e}_1 \quad (6.100)$$

whose matrix representation is

$$B = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (6.101)$$

### 6.3 Linear operators

---

Both by considering their action on the basis, or by matrix multiplication, we have

$$AB = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \quad BA = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix}, \quad (6.102)$$

namely  $AB$  leaves  $\mathbf{e}_1$  in itself and deletes  $\mathbf{e}_2$ , while  $BA$  has the opposite effect.

We may define the *commutator*

$$[A, B] = AB - BA = -[B, A] \quad (6.103)$$

and we will have in general  $[A, B] \neq 0$ . We will study later the properties of the commutator, when we will study the solution of the Laplace equation in spherical coordinates and its relation to Quantum Mechanics. In the meanwhile, we study some properties of self-adjoint and anti-adjoint commuting operators, i.e. such that

$$[A, B] = 0 \quad (6.104)$$

Let us assume  $A$  and  $B$  admit a orthonormal basis of eigenvectors, as for example self-adjoint and anti-adjoint operators (or, more in general, normal operators, as we will show later) do, and that they commute  $AB = BA$ . If  $\mathbf{v}_\lambda$  is an eigenvector of  $A$  with eigenvalue  $\lambda$ , we have

$$A(B\mathbf{v}_\lambda) = AB\mathbf{v}_\lambda = BA\mathbf{v}_\lambda = \lambda(B\mathbf{v}_\lambda) \quad (6.105)$$

This equation tells us that  $B\mathbf{v}_\lambda$  is an eigenvector of  $A$ , with eigenvalue  $\lambda$ , i.e. the subspace generated by the linear combinations of all the eigenvectors corresponding to the eigenvalue  $\lambda$  is left invariant by the action of  $B$ . Since  $B$  is self-adjoint (or anti-adjoint), we may diagonalise it on the subspace, i.e. find an orthonormal basis of eigenvectors of  $B$  in the subspace. Each of these eigenvectors will be at the same time an eigenvector of  $A$  (with eigenvalue  $\lambda$ , since it is a vector of the subspace) and of  $B$ . We may do that for all the eigenvalues  $\lambda_i$  of  $A$  and obtain a *common basis of eigenvectors* of  $A$  and  $B$ . These vectors form a orthonormal basis

$$\begin{aligned} \mathbf{e}_{\lambda_i, \gamma_j}, \quad A\mathbf{e}_{\lambda_i, \gamma_j} &= \lambda_i \mathbf{e}_{\lambda_i, \gamma_j}, \quad B\mathbf{e}_{\lambda_i, \gamma_j} = \gamma_j \mathbf{e}_{\lambda_i, \gamma_j} \\ \langle \mathbf{e}_{\lambda_i, \gamma_j}, \mathbf{e}_{\lambda_k, \gamma_l} \rangle &= \delta_{i,k} \delta_{j,l} \end{aligned} \quad (6.106)$$

Since in general to the eigenvalues of a (self-adjoint) operator may correspond a subspace with dimension larger than one, although it is possible to find a orthonormal basis of eigenvalues corresponding to it, it is not possible to identify in a unique way all the vector of the basis through the eigenvalue. By using enough commuting operators, it is possible to identify all vectors



in the basis through a set of eigenvalues corresponding to each operator, a process which is extremely useful in Quantum Mechanics (as we will see in our analysis of the Hydrogen atom).

### 6.3.7 Normal operators

The most general class of operators that admits an orthonormal basis of eigenvectors is that of normal operators

$$[A, A^\dagger] = 0 \quad (6.107)$$

To prove it, we first see that *any operator*  $A$  may be written as the sum of a self-adjoint and anti-adjoint operator

$$A = A_{SA} + A_{AA} = \frac{1}{2}[(A + A^\dagger) + (A - A^\dagger)] \quad (6.108)$$

since

$$(A_{SA})^\dagger = \frac{1}{2}(A + A^\dagger)^\dagger = \frac{1}{2}(A^\dagger + A) = A_{SA} \quad (6.109)$$

and

$$(A_{AA})^\dagger = \frac{1}{2}(A - A^\dagger)^\dagger = \frac{1}{2}(A^\dagger - A) = -A_{AA} \quad (6.110)$$

Now, if  $A$  is normal, we have

$$\begin{aligned} A_{SA}A_{AA} &= \frac{1}{4}(A^2 - (A^\dagger)^2 - AA^\dagger + A^\dagger A) = \frac{1}{4}(A^2 - (A^\dagger)^2) \\ &= \frac{1}{4}(A^2 - (A^\dagger)^2 - A^\dagger A + AA^\dagger) = A_{AA}A_{SA} \end{aligned} \quad (6.111)$$

Since  $A_{SA}$  and  $A_{AA}$  admit an orthonormal basis of eigenvalues (one being self-adjoint, and the other one being anti-adjoint), and commute, they admit a *common* orthonormal basis of eigenvalues

$$\begin{aligned} \mathbf{e}_{\lambda_i, \gamma_j}, \quad A_{SA}\mathbf{e}_{\lambda_i, \gamma_j} &= \lambda_i\mathbf{e}_{\lambda_i, \gamma_j}, \quad A_{AA}\mathbf{e}_{\lambda_i, \gamma_j} = \gamma_j\mathbf{e}_{\lambda_i, \gamma_j} \\ \langle \mathbf{e}_{\lambda_i, \gamma_j}, \mathbf{e}_{\lambda_k, \gamma_l} \rangle &= \delta_{i,k}\delta_{j,l} \end{aligned} \quad (6.112)$$

so that these vectors are an orthonormal basis of eigenvalues of  $A$

$$A\mathbf{e}_{\lambda_i, \gamma_j} = (A_{SA} + A_{AA})\mathbf{e}_{\lambda_i, \gamma_j} = (\lambda_i + \gamma_j)\mathbf{e}_{\lambda_i, \gamma_j} \quad (6.113)$$

Since the  $\lambda$  are real, and the  $\gamma$  are imaginary, in general the eigenvalues of a normal operator are complex.

## 6.3 Linear operators

---

### 6.3.8 Unitary operator

An important class of normal operators are the Unitary operators  $U$  such that

$$U^\dagger = U^{-1} \Rightarrow U^\dagger U = U U^\dagger = \mathbf{1} \quad (6.114)$$

They have the property that

$$\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, U^\dagger U \mathbf{y} \rangle = \langle U \mathbf{x}, U \mathbf{y} \rangle \quad (6.115)$$

namely  $U$  conserves the scalar product.

By being normal,  $U$  assumes a basis of eigenvectors. Its eigenvalues  $\lambda$  are such that

$$\langle \mathbf{v}_\lambda, \mathbf{v}_\lambda \rangle = \langle U \mathbf{v}_\lambda, U \mathbf{v}_\lambda \rangle = \langle \lambda \mathbf{v}_\lambda, \lambda \mathbf{v}_\lambda \rangle = \lambda^* \lambda \langle \mathbf{v}_\lambda, \mathbf{v}_\lambda \rangle \quad (6.116)$$

So that  $|\lambda| = 1$ , or  $\lambda = e^{i\theta}$ .

Let us consider an orthonormal basis  $\mathbf{e}_i$ , and a unitary operator  $U$ . If we define

$$\tilde{\mathbf{e}}_i = U \mathbf{e}_i \quad (6.117)$$

we see that the  $\tilde{\mathbf{e}}_i$  are a new orthonormal basis

$$\langle \tilde{\mathbf{e}}_i, \tilde{\mathbf{e}}_j \rangle = \langle \mathbf{e}_i, \mathbf{e}_j \rangle = \delta_{i,j} \quad (6.118)$$

On the other end, let us assume we know two different orthonormal bases,  $\mathbf{e}_i$  and  $\mathbf{u}_i$ . For example, the  $\mathbf{u}_i$  may be eigenvectors of  $A$ , a normal operator. Each  $\mathbf{u}$  is obviously possible to write as a combination of the  $\mathbf{e}_i$

$$\mathbf{u}_i = \sum_j u_{i,j} \mathbf{e}_j \quad (6.119)$$

The operator whose matrix representation is  $u_{i,j}$  is, by construction the operator that generates  $\mathbf{u}_i$  when acting on  $\mathbf{e}_i$ . We may show that this operator is unitary. Indeed, calling  $U$  the operator, we have

$$(U U^\dagger)_{i,j} = \sum_k U_{i,k} (U^\dagger)_{k,j} = \sum_k u_{i,k} u_{j,k}^* = \langle \mathbf{u}_i, \mathbf{u}_j \rangle = \delta_{i,j} \quad (6.120)$$

It is also trivial to show that, if as stated above the  $\mathbf{u}_i$  are the eigenvectors of the normal operator  $A$ ,  $U^\dagger A U$  is diagonal in the original basis

$$\begin{aligned} (U^\dagger A U)_{i,j} &= \langle \mathbf{e}_i, U^\dagger A U \mathbf{e}_j \rangle = \langle U \mathbf{e}_i, A U \mathbf{e}_j \rangle = \\ &= \langle \mathbf{u}_i, A \mathbf{u}_j \rangle = \langle \mathbf{u}_i, \lambda_j \mathbf{u}_j \rangle = \lambda_j \langle \mathbf{u}_i, \mathbf{u}_j \rangle = \lambda_j \delta_{i,j} \end{aligned} \quad (6.121)$$

## 6.4 Dirac notation

### 6.4.1 Bras, kets, and scalar products

After this quick revision of (finite dimensional) vector spaces, we are ready to learn Dirac's notation and how it simplifies computations.

We defined the scalar product as an operator that takes two vectors and gives a number

$$\langle *, * \rangle \rightarrow \mathbb{C} \quad (6.122)$$

where the  $*$  represent slots that may be filled with any vector.

We may also decide to fill the first slot with a vector,

$$\langle \mathbf{v}, * \rangle \rightarrow \mathbb{C} \quad (6.123)$$

and obtain an operator that associates to each vector  $*$  a complex number. We will have an operator *for each*  $\mathbf{v}$ . Dirac notation is based on providing a different name and symbol for the vectors and for their version as operators that give numbers when applied on vectors, and to express all mathematical expressions in vector spaces through a combination of such symbols<sup>7</sup>.

Let us denote the vectors in the space through a symbol called *ket*, i.e. the vector  $\mathbf{v}$  is represented by

$$|v\rangle \quad (6.124)$$

The corresponding operator  $\langle \mathbf{v}, * \rangle$  is represented through a symbol named *bra*<sup>8</sup>

$$\langle v| \quad (6.125)$$

The scalar product between  $\mathbf{x}$  and  $\mathbf{y}$ ,  $\langle \mathbf{x}, \mathbf{y} \rangle$  is a *braket*

$$\langle x|y\rangle \quad (6.126)$$

We may write the product of a ket by a complex number  $\alpha$  equivalently as<sup>9</sup>

$$\alpha|v\rangle = |\alpha v\rangle \quad (6.127)$$

The linear combination of two kets may be written as

$$\alpha|v\rangle + \beta|u\rangle = |\alpha v\rangle + |\beta u\rangle = |\alpha v + \beta u\rangle \quad (6.128)$$

---

<sup>7</sup>Although in eq. (6.123) we defined the operator through the scalar product, in Dirac's notations such operators are more fundamental than the scalar product and define it.

<sup>8</sup>At a dinner with other academics in Cambridge that were discussing about who between them introduced new words in the English language, Dirac told them that he invented the word "bra". He never explained to the unbelieving colleagues that he did not refer to lingerie.

<sup>9</sup>We will use Greek letters for complex numbers and Latin letters for bras and kets.

## 6.4 Dirac notation

---

Bras are obviously linear

$$\langle x | (\alpha |v\rangle + \beta |u\rangle) = \alpha \langle x | v \rangle + \beta \langle x | u \rangle \quad (6.129)$$

The bra corresponding to  $|\alpha u\rangle$  is

$$\langle \alpha u | = \alpha^* \langle u | \quad (6.130)$$

### 6.4.2 Ketbras, projectors and representations of unity

#### Ketbra

What happens if we write a *ketbra*?

$$|v\rangle\langle u| \quad (6.131)$$

Is it a meaningful symbol? It is, and it represents a linear operator on the vector space. Let us apply it on a ket (vector)

$$(|v\rangle\langle u|)|x\rangle = |v\rangle\langle u|x\rangle = (\langle u|x\rangle)|v\rangle \quad (6.132)$$

The bracket  $\langle u|x\rangle$  is just a number, so we moved it on the front, just to show that the effect of the operator is to produce a ket (vector) parallel to  $|v\rangle$  but multiplied by the scalar  $\langle u|x\rangle$ .

Since a ketbra is an operator, we may wonder how to write its adjoint. We may show that if  $A = |x\rangle\langle y|$  then

$$(|x\rangle\langle y|)^\dagger = |y\rangle\langle x| \quad (6.133)$$

Indeed (see eq. 6.144 above for the Dirac operator notation)

$$\begin{aligned} (\langle v|y\rangle\langle x|u\rangle)^* &= \langle u|x\rangle\langle y|v\rangle = \langle u|Av\rangle = \langle A^\dagger u|v\rangle = \left(\langle v|A^\dagger u\rangle\right)^* \\ &\Rightarrow \langle v|y\rangle\langle x|u\rangle = \langle v|A^\dagger u\rangle \end{aligned} \quad (6.134)$$

#### projector

A very important case is when  $|v\rangle$  is normalised, and we consider the *projector*

$$|v\rangle\langle v| \quad (6.135)$$

that projects a vector along  $|v\rangle$ , i.e.

$$|v\rangle\langle v|u\rangle = (\langle v|u\rangle)|v\rangle \quad (6.136)$$

It is easier to understand its meaning by taking  $|v\rangle$  corresponding to  $\mathbf{e}_i$ , and

$$\mathbf{u} = \sum_j u_j \mathbf{e}_j \quad (6.137)$$

so that the action of the projector is

$$\langle i|u\rangle = \langle \mathbf{e}_i, \mathbf{u} \rangle \mathbf{e}_i = u_i \mathbf{e}_i \quad (6.138)$$

If we consider a few orthonormal vectors (not necessarily a basis for the whole space)  $|i\rangle$  (representing  $\mathbf{e}_i$ ), then

$$\sum_i |i\rangle \langle i| \quad (6.139)$$

is a projector on the the subspace generated by the linear combinations of the  $\mathbf{e}_i$  (by being the sum of the projectors corresponding to each basis vector).

### Representations of identity

If we take the sum over all the basis, we get a projector on the whole space, which is just the identity!

$$\sum_{i=1}^n |i\rangle \langle i| = \mathbf{1} \quad (6.140)$$

To see it in explicit terms let us remember

$$\mathbf{u} = \sum_i u_i \mathbf{e}_i, \quad u_i = \langle \mathbf{e}_i, \mathbf{u} \rangle \quad (6.141)$$

Written in bra-ket notation, the latter reads

$$|u\rangle = \sum_i \langle i|u\rangle |i\rangle = \left( \sum_i |i\rangle \langle i| \right) |u\rangle \quad (6.142)$$

This is true for each  $|u\rangle$ , provided that the  $|i\rangle$  are a basis, so that eq. (6.140) follows.

## 6.4.3 Operators

### General notation

The action of operator  $A$  on ket  $|v\rangle$  is

$$|Av\rangle = A|v\rangle \quad (6.143)$$

## 6.4 Dirac notation

---

while the scalar product  $\langle \mathbf{x}, A\mathbf{y} \rangle$  may be written as

$$\langle x|Ay \rangle = \langle x|A|y \rangle \quad (6.144)$$

the latter notation being more used. The Dirac notation is asymmetric in its treatment on operators on the two sides of the scalar product, since we have for  $\langle A\mathbf{x}, \mathbf{y} \rangle$  only

$$\langle Ax|y \rangle \quad (6.145)$$

This asymmetry may be the only unpleasant feature of the notation. I like to write

$$\langle Ax|y \rangle = \langle x|A||y \rangle \quad (6.146)$$

but it is only a personal notation.

The matrix representation of  $A$  in the  $|i\rangle$  basis is clearly given by

$$A_{i,j} = \langle i|A|j \rangle \quad (6.147)$$

We are now ready to see how the Dirac formalism make some computations trivial. The matrix representation of  $BA$  is

$$\langle i|BA|j \rangle \quad (6.148)$$

We may insert an identity operator where ever we want and easily obtain eq. (6.64)

$$\langle i|B\mathbf{1}A|j \rangle = \langle i|B| \sum_k |k\rangle \langle k|A|j \rangle = \sum_k \langle i|B|k\rangle \langle k|A|j \rangle = \sum_k B_{i,k}A_{k,j} \quad (6.149)$$

It is a useful exercise to rewrite all the equations above using the new formalism, for example

$$A_{i,j}^\dagger = \langle i|A^\dagger|j \rangle = \langle i|A|j \rangle = (\langle j|A|i \rangle)^* = (A_{j,i})^* \quad (6.150)$$

### Normal operators

A normal operator admits an orthonormal basis of eigenvectors. We may enumerate them  $i = 1, \dots, n$ , and call the eigenvalues  $\lambda_i$  (we may have degeneracy,  $\lambda_i = \lambda_j$  for  $i \neq j$ ). Then the normal operator may be written as

$$A = \sum_i \lambda_i |i\rangle \langle i| \quad (6.151)$$

Indeed, in the basis of its eigenvectors, the matrix representation is

$$A_{i,j} = \sum_k \lambda_k \langle i|k\rangle \langle k|j \rangle = \sum_k \lambda_k \delta_{i,k} \delta_{k,j} = \lambda_i \delta_{i,j} \quad (6.152)$$

or, in an equivalent way

$$A|i\rangle = \sum_j \lambda_j |j\rangle \langle j|i\rangle = \sum_j \lambda_j |j\rangle \delta_{i,j} = \lambda_i |i\rangle \quad (6.153)$$

When defined on a basis,  $A$  is uniquely defined on the whole space, so that eq. (6.151) gives the wanted operator.

### Unitary operators

A unitary operator brings an orthonormal basis  $|i\rangle$  in another orthonormal basis  $|i'\rangle$ , so that it is by construction

$$U = \sum_i |i'\rangle \langle i| \quad (6.154)$$

where the prime in  $i'$  shows a change in the vector, not in the indexes, i.e. we may write

$$\langle i'|j'\rangle = \delta_{i,j} \quad (6.155)$$

and

$$U|j\rangle = \sum_i |i'\rangle \langle i|j\rangle = \sum_i |i'\rangle \delta_{i,j} = |j'\rangle \quad (6.156)$$

as wanted. We have

$$U^\dagger = \sum_i |i\rangle \langle i'| \quad (6.157)$$

and

$$U^\dagger U = \sum_j |j\rangle \langle j'| \sum_i |i'\rangle \langle i| = \sum_i \sum_j |j\rangle \langle j'|i'\rangle \langle i| = \sum_i \sum_j \delta_{i,j} |j\rangle \langle i| = \sum_i |i\rangle \langle i| = \mathbb{1} \quad (6.158)$$

The matrix representation of  $U$  is

$$U_{i,j} = \sum_k \langle i|k'\rangle \langle k|j\rangle = \sum_k \delta_{k,j} \langle i|k'\rangle = \langle i|j'\rangle \quad (6.159)$$

The matrix representation of an operator  $A$  in the basis  $|i'\rangle$  is

$$A'_{i,j} = \langle i'|A|j'\rangle = \sum_{k,l} \langle i'|k\rangle \langle k|A|l\rangle \langle l|j'\rangle = U_{i,k}^\dagger A_{k,l} U_{l,j} \quad (6.160)$$

### 6.5 Function spaces

#### 6.5.1 Introduction

We have seen that the concept of vector space may be applied also to functions, as for example by defining a vector space of periodic functions, or a vector space of solutions of a differential equations. Furthermore, a differential operators, as for example  $\partial_t$ ,  $\partial_t^2$ , or  $\nabla^2$ , are linear operators on such vector spaces, since

$$\nabla^2(\alpha f(\mathbf{x}) + \beta g(\mathbf{x})) = \alpha \nabla^2 f(\mathbf{x}) + \beta \nabla^2 g(\mathbf{x}) \quad (6.161)$$

It is thus clear that we are interested in developing the theory of these *functional vector spaces* as part of our theory of partial differential equations. Nevertheless, as hinted by the fact that our general solutions in the form of Fourier series included in general *infinite terms*, these spaces are not finite dimensional. The theory of functional analysis, dealing with these vector spaces, is not trivial and goes beyond the limits of this course. Nevertheless, we may learn the basic tools and methods of the theory, that we will develop, without proofs, using Dirac's notation and in analogy with the finite dimensional case.

If  $f(x)$  is a function in our vector space, for example a periodic function between 0 and  $L$ , we may denote the corresponding ket as  $|f\rangle$ . We may formally define the sum and product by a complex number, and we identify

$$|\alpha f + \beta g\rangle \quad (6.162)$$

as the function

$$\alpha f(x) + \beta g(x) \quad (6.163)$$

We may also formally define the bra

$$\langle f| \quad (6.164)$$

and ask it to satisfy the formal properties that define

$$\langle f|g\rangle \quad (6.165)$$

as a scalar product. But it is clear that if we want our theory to be of any practical value, we need to find a way to express this bracket in an explicit way.



### 6.5.2 Continuum limit

Let us first deal with some partial information. We may compute the value of the function in  $i = 1, \dots, n$  points,

$$f_i = f(x_i), \quad x_i = -\frac{\Delta x}{2} + i\Delta x, \quad \Delta x = \frac{L}{n} \quad (6.166)$$

This is just a  $n$  dimensional vector, so that we may treat it with the usual methods

$$f_i = f(x_i) = \langle x_i | f \rangle \quad (6.167)$$

$$\langle x_i | x_j \rangle = \delta_{i,j} \quad (6.168)$$

$$\sum_i |x_i\rangle \langle x_i| = \mathbb{1} \quad (6.169)$$

and

$$\langle f | g \rangle = \sum_i \langle f | x_i \rangle \langle x_i | g \rangle = \sum_i (f(x_i))^* g(x_i) \quad (6.170)$$

This description works at fixed  $n$ , but for finite  $n$  we may have only an approximate information, and eventually we want to perform the continuum limit,  $n \rightarrow \infty$  and  $\Delta x \rightarrow 0$ . But if we perform the limit with this naive approach, we soon meet a problem.

Let us consider a constant function  $f(x) = C$ . We have

$$\langle f | f \rangle = nC \quad (6.171)$$

a result that would diverge as  $n \rightarrow \infty$ . Let us then define

$$\sum_i |x_i\rangle \langle x_i| \Delta x = \mathbb{1} \quad (6.172)$$

so that the scalar product is

$$\langle f | g \rangle = \sum_i \langle f | x_i \rangle \langle x_i | g \rangle \Delta x = \sum_i (f(x_i))^* g(x_i) \Delta x \quad (6.173)$$

Now the scalar product of the constant function is

$$\langle f | f \rangle = nC\Delta x = nC\frac{L}{n} = CL \quad (6.174)$$

a  $n$  independent result. We may now easily see that a candidate for the eigen-kets  $|x_i\rangle$  are functions

$$\delta_i(x) = \begin{cases} \frac{1}{\sqrt{\Delta x}} & \text{for } x \in [(i-1)\Delta x, i\Delta x) \\ 0 & \text{elsewhere} \end{cases} \quad (6.175)$$

## 6.5 Function spaces

---

We finally send  $n$  to infinity. The kets  $|x\rangle$  now represent an infinitely narrow and high function, a *Dirac delta function*<sup>10</sup>. The representation of the identity is

$$\int_0^L |x\rangle\langle x| dx = \mathbb{1} \quad (6.176)$$

so that

$$\langle x|y\rangle = \delta(x - y) \quad (6.177)$$

$$f(x) = \langle x|f\rangle \quad (6.178)$$

and

$$\langle f|g\rangle = \int_0^L \langle f|x\rangle\langle x|g\rangle dx = \int_0^L (f(x))^* g(x) dx \quad (6.179)$$

The scalar product between  $f$  and  $g$  becomes thus an integral of the product of the complex conjugate of  $f$  and  $g$ .

What is the meaning of the Dirac delta function? It may be defined by the following expression

$$f(x) = \langle x|f\rangle = \int_0^L \langle x|y\rangle\langle y|f\rangle dy = \int_0^L \delta(x - y) f(y) dy \quad (6.180)$$

When we integrate the product of a function  $f(y)$  and the delta function  $\delta(x - y)$ , the result is the value of  $f$  in  $x$  (if  $x \in [0, L]$ ). We will study these functions in deeper later.

### 6.5.3 Adjoint operator

An adjoint operator is defined by

$$\langle f, Ag\rangle = \langle A^\dagger f, g\rangle \quad (6.181)$$

for finite or infinite dimension spaces.

Let us consider the operator  $D = \frac{d}{dx}$  on the space of periodic functions. We have

$$\begin{aligned} \langle f, Dg\rangle &= \int_0^L (f(x))^* \left( \frac{d}{dx} g(x) \right) dx = \int_0^L \frac{d}{dx} ((f(x))^* g(x)) dx + \int_0^L \left( -\frac{d}{dx} (f(x))^* \right) g(x) dx \\ &= [(f(x))^* g(x)]_0^L + \int_0^L \left( -\frac{d}{dx} f(x) \right)^* g(x) dx \Rightarrow D^\dagger = -\frac{d}{dx} = -D \end{aligned} \quad (6.182)$$

where we performed an integration by part and used the fact that the functions are periodic to delete the total derivative term.

---

<sup>10</sup>As stated above, this is an ill defined function, but a formal definition may be provided as part of the theory of distributions.

We may now consider the operator  $P = -i\frac{d}{dx}$ , again *on the space of periodic functions*. We have

$$\begin{aligned} \langle f, Pg \rangle &= \int_0^L (f(x))^* \left( -i\frac{d}{dx}g(x) \right) dx = \\ &= -i \int_0^L \frac{d}{dx} ((f(x))^*g(x)) dx + i \int_0^L \left( \frac{d}{dx}(f(x))^* \right) g(x) dx \quad (6.183) \\ &= -i [(f(x))^*g(x)]_0^L + \int_0^L \left( -i\frac{d}{dx}f(x) \right)^* g(x) dx \Rightarrow P^\dagger = P \end{aligned}$$

The  $i$  in front of the operator changes sign with conjugation, and allows the operator to be self adjoint. In both cases, the operator's adjoint had a simple form because we had the correct boundary conditions, that thus have a very important role in infinite dimensional vector spaces.

Since

$$\frac{d^2}{dx^2} \equiv D^2 = -P^2 \Rightarrow (D^2)^\dagger = D^\dagger D^\dagger = (-)^2 D^2 \quad (6.184)$$

the second derivative operator is self-adjoint. This has an important effect on the possible eigenvalues of the operator  $\nabla^2$ .

#### 6.5.4 Enumerable basis

We have seen that the operator  $P = -i\frac{d}{dx}$  is self-adjoint. In the finite dimensional case, a self-adjoint operator has a orthonormal basis of eigenvectors with real eigenvalues. You can easily check that in the proof regarding the eigenvalue being real we did not use the fact that the space was finite or not, so that a self-adjoint operator *always* has real eigenvalues<sup>11</sup>.

What is an eigenvalue of a linear operator like  $P$ ? It is the solution of the differential equation

$$P|\alpha\rangle = \alpha|\alpha\rangle \quad (6.185)$$

We already know that a solution is given by the a ket such that

$$\langle x|n\rangle = \sqrt{\frac{1}{L}} e^{i\frac{2\pi n}{L}x} \quad (6.186)$$

the eigenvalue being

$$\frac{2\pi n}{L}, \quad n \in \mathbb{Z} \quad (6.187)$$

---

<sup>11</sup>In quantum mechanics, these eigenvalues play the role of the values assumed by physical observables, and for this reason only self-adjoint operators may be used to build physical observables. This is the reason the momentum operator in Schrödinger's equation includes the  $i$  term.

## 6.5 Function spaces

---

We have already shown that

$$\frac{1}{L} \int_0^L e^{-i\frac{2\pi m}{L}} e^{i\frac{2\pi n}{L}} dx = \langle m|n \rangle = \delta_{m,n} \quad (6.188)$$

This is an enumerable basis for the functional vector space, since each function that is smooth enough may be written as

$$|f\rangle = \sum_n |n\rangle \langle n|f\rangle \quad (6.189)$$

The coefficients of the Fourier expansion are the

$$|f\rangle = \langle n|f\rangle = \sqrt{\frac{1}{L}} \int_0^L e^{i\frac{2\pi n}{L}} f(x) dx \quad (6.190)$$

The Dirac notations helps us in derive, remember and manipulate these relations. We may, for example, remember that the functions

$$s_{n_x}^{L_x}(x) s_{n_y}^{L_y}(y) s_{n_z}^{L_z}(z) = \langle x, y, z | n_x, n_y, n_z \rangle \quad (6.191)$$

are the eigenkets of the the operator  $\nabla^2$  with zero boundary conditions (the quantum particle in the box). As a result

$$\sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} \sum_{n_z=1}^{\infty} |n_x, n_y, n_z\rangle \langle n_x, n_y, n_z| = \mathbb{1} \quad (6.192)$$

so that

$$|f\rangle = \sum_{n_x=1}^{\infty} \sum_{n_y=1}^{\infty} \sum_{n_z=1}^{\infty} |n_x, n_y, n_z\rangle \langle n_x, n_y, n_z|f\rangle \quad (6.193)$$

You may compare this to eqs. (3.96,3.97).

## Chapter 7

# Laplace equation in spherical coordinates

### 7.1 Introduction

We now study a very important problem in mathematical physics: the Laplace equation in spherical coordinates. Recalling eq. (1.83), we have

$$\frac{1}{r^2} \partial_r (r^2 \partial_r \Phi(r, \theta, \varphi)) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta \Phi(r, \theta, \varphi)) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \Phi(r, \theta, \varphi) = 0 \quad (7.1)$$

### 7.2 Axial symmetry

Let us first study a simpler problem, by considering a function that does not depend on  $\varphi$

$$\Phi = \Phi(r, \theta) \quad (7.2)$$

so that the equation becomes

$$\frac{1}{r^2} \partial_r (r^2 \partial_r \Phi(r, \theta)) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta \Phi(r, \theta)) = 0 \quad (7.3)$$

The boundary condition may be the value assumed on a sphere of radius  $\bar{r}$

$$\Phi(\bar{r}, \theta) = f(\theta) \quad (7.4)$$

We play our usual trick

$$\Phi = R(r)\Theta(\theta) \quad (7.5)$$

and get

$$\Theta(\theta) \frac{1}{r^2} \partial_r (r^2 \partial_r R(r)) = -R(r) \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta \Theta(\theta)) \quad (7.6)$$

## 7.2 Axial symmetry

---

multiply by  $r^2/\Phi$  and get

$$\frac{1}{R(r)}\partial_r(r^2\partial_r R(r)) = C = -\frac{1}{\Theta(\theta)\sin\theta}\partial_\theta(\sin\theta\partial_\theta\Theta(\theta)) \quad (7.7)$$

### 7.2.1 Radial equation

Let us first study the radial equation

$$\partial_r(r^2\partial_r R(r)) = CR(r) \quad (7.8)$$

As we did in the polar coordinate case (section 5.3), we try a power function

$$R(r) = r^l \quad (7.9)$$

as solution. We have

$$\begin{aligned} \partial_r r^l &= lr^{l-1} \\ r^2(\partial_r r^l) &= lr^{l+1} \\ \partial_r(r^2(\partial_r r^l)) &= l(l+1)r^l \end{aligned} \quad (7.10)$$

This means that  $r^l$  is a solution of eq. (7.8) provided that

$$l(l+1) = C \quad (7.11)$$

### 7.2.2 Angular equation

The angular equation is then

$$\frac{1}{\sin\theta}\partial_\theta(\sin\theta\partial_\theta\Theta(\theta)) = -l(l+1)\Theta(\theta) \quad (7.12)$$

The equation is simplified if we define

$$u = \cos\theta \quad (7.13)$$

from which we get

$$u^2 = \cos^2\theta \Rightarrow \sin^2\theta = 1 - \cos^2\theta = 1 - u^2 \quad (7.14)$$

and

$$\frac{\partial}{\partial\theta} = \frac{\partial u}{\partial\theta} \frac{\partial}{\partial u} \Rightarrow \partial_\theta = -\sin\theta\partial_u \quad (7.15)$$

The equation in  $u$  becomes

$$\begin{aligned} \frac{1}{\sin\theta}\partial_\theta(\sin\theta(-\sin\theta)\partial_u\Theta(u)) &= -l(l+1)\Theta(u) \\ \Rightarrow \frac{1}{\sin\theta}(-\sin\theta)\partial_u((u^2-1)\partial_u\Theta(u)) &= -l(l+1)\Theta(u) \\ \Rightarrow -\partial_u[(u^2-1)\partial_u\Theta(u)] &= -l(l+1)\Theta(u) \\ \Rightarrow \partial_u[(u^2-1)\partial_u\Theta(u)] &= l(l+1)\Theta(u) \end{aligned} \quad (7.16)$$

Let us try again a power function

$$\Theta(u) = u^n \quad (7.17)$$

so that

$$\begin{aligned} \partial_u u^n &= nu^{n-1} \\ (u^2 - 1)\partial_u u^n &= nu^{n+1} - nu^{n-1} = nu^{n-1}[u^2 - 1] \\ \partial_u [(u^2 - 1)\partial_u \Theta(u)] &= n(n+1)u^n - n(n-1)u^{n-2} \end{aligned} \quad (7.18)$$

and thus

$$n(n+1)u^n - n(n-1)u^{n-2} = l(l+1)u^n \quad (7.19)$$

Different powers of  $u$  are found on the left and right side. This means that a single power function of  $u$  will not satisfy the equation. Nevertheless, we may still try to use a power series, hoping that if we correctly choose the coefficients the series will satisfy the equation. We try then

$$\Theta(u) = \sum_{n=0}^{\infty} a_n u^n \quad (7.20)$$

so that

$$\begin{aligned} \sum_{n=0}^{\infty} a_n n(n+1)u^n - \sum_{n=0}^{\infty} a_n n(n-1)u^{n-2} &= \sum_{n=0}^{\infty} a_n l(l+1)u^n \\ \Rightarrow \sum_{n=0}^{\infty} a_n u^n [l(l+1) - n(n+1)] + \sum_{n=0}^{\infty} a_n n(n-1)u^{n-2} &= 0 \end{aligned} \quad (7.21)$$

since

$$n(n-1) = 0 \quad \text{for } n = 0, 1 \quad (7.22)$$

we may write the last equation as

$$\sum_{n=0}^{\infty} a_n u^n [l(l+1) - n(n+1)] + \sum_{n=2}^{\infty} a_n n(n-1)u^{n-2} = 0 \quad (7.23)$$

or, using

$$n' = n - 2 \Rightarrow n = n' + 2 \quad (7.24)$$

$$\begin{aligned} \sum_{n=0}^{\infty} a_n u^n [l(l+1) - n(n+1)] + \sum_{n'=0}^{\infty} a_{n'+2} (n'+2)(n'+1)u^{n'} &= 0 \\ \Rightarrow \sum_{n=0}^{\infty} a_n u^n [l(l+1) - n(n+1)] = - \sum_{n=0}^{\infty} a_{n+2} u^n [(n+2)(n+1)] & \quad (7.25) \\ \Rightarrow \sum_{n=0}^{\infty} u^n \left( a_n [l(l+1) - n(n+1)] + a_{n+2} [(n+2)(n+1)] \right) &= 0 \end{aligned}$$

## 7.2 Axial symmetry

---

satisfied if

$$a_n[l(l+1) - n(n+1)] + a_{n+2}[(n+2)(n+1)] = 0 \quad \forall n \quad (7.26)$$

or

$$a_{n+2} = -a_n \frac{l(l+1) - n(n+1)}{(n+2)(n+1)} \quad (7.27)$$

Eq. (7.27) is a recursion equation that gives the value of the coefficient  $a_{n+2}$  once the  $a_n$  is given. Namely, if  $a_0$  and  $a_1$  are provided, we have the whole series.

### 7.2.3 Legendre polynomials

We also notice that, for high enough  $n$  we have

$$a_{n+2} \approx a_n \frac{n(n+1)}{(n+2)(n+1)} \approx a_n \quad (7.28)$$

so that for  $\theta = 0$ ,  $u = 1$ , the series diverges

$$\sum_{n=0}^{\infty} a_n \rightarrow \infty \quad (7.29)$$

How can we avoid this problem? If we have, for some  $\bar{n}$

$$l(l+1) = \bar{n}(\bar{n}+1) \Rightarrow a_{\bar{n}+2} = 0 \Rightarrow a_{\bar{n}+2k} = 0 \quad (7.30)$$

then we may get all the even (if  $n$  is even) or odd (if  $n$  is odd) coefficients larger than  $n$  to be zero. The  $n$  are positive integers, and thus to satisfy eq. (7.30) we need  $l \in \mathbb{Z}$ . Namely

$$\begin{aligned} l(l+1) = \bar{n}(\bar{n}+1) &\Rightarrow (l^2 - \bar{n}^2) + (l - \bar{n}) = 0 \Rightarrow (l + \bar{n})(l - \bar{n}) + l - \bar{n} = 0 \\ &\Rightarrow (l - \bar{n})(l + \bar{n} + 1) = 0 \Rightarrow l = \bar{n} \text{ or } l = -\bar{n} - 1 \end{aligned} \quad (7.31)$$

So that for each  $n$  we will have 2 possible  $l$  value, one  $l \geq 0$ , and one  $l < 0$ .

If  $l(l+1) = \bar{n}(\bar{n}+1)$  is odd, we may not use eq. (7.30) to set any  $a_n = 0$  for even  $n$ , and thus we will choose  $a_0 = 0$ . On the other hand, for even  $l(l+1) = \bar{n}(\bar{n}+1)$  we choose  $a_1 = 0$ . We call these solutions the Legendre polynomials,  $P_l(u)$ .

For example,  $P_0$  is characterised by having

$$l(l+1) = \bar{n}(\bar{n}+1) = 0 \Rightarrow \bar{n} = 0, l = 0 \text{ or } l = -1 \quad (7.32)$$



---

### Laplace equation in spherical coordinates

---

$P_0$  (as all the  $P_l$ ) is determined only by the value of  $\bar{n}$ , so that we will not talk about  $P_l$  for negative  $l$ . We assign a given  $a_0$  and set  $a_1 = 0$ , and we get all the other  $a_n = 0$ , so that

$$P_0(u) = a_0 = \text{const} \quad (7.33)$$

For  $P_1$  we assign  $a_1$ , set  $a_0 = 0$ , and get all the other coefficients as zero, so that

$$P_1(u) = a_1 u \quad (7.34)$$

For  $P_2$  we assign  $a_0$ , set  $a_1 = 0$  and get all the other coefficients as zero, with the exclusion of

$$a_2 = -a_0 \frac{l(l+1) - 0}{(1)(2)} = -3 \quad (7.35)$$

so that

$$P_2(u) = a_0(1 - 3u^2) \quad (7.36)$$

etc. These polynomials are usually defined so that  $P(1) = 1$ , namely

$$P_0(u) = 1 \quad P_1(u) = u \quad P_2(u) = \frac{3}{2}u^2 - \frac{1}{2} \quad (7.37)$$

This choice *does not* normalise the function  $P_l$  in the sense of the previous chapter

$$\int (P_l(u))^* P_l(u) du = 1 \quad (7.38)$$

Indeed, since  $u = \cos(\theta)$  varies in  $[-1, 1]$ , we have, for example

$$\int_{-1}^1 (P_0(u))^* P_0(u) du = \int_{-1}^1 du = 2 \quad (7.39)$$

We have anyway

$$\int_{-1}^1 (P_0(u))^* P_1(u) du \propto \int_{-1}^1 u du = 0 \quad (7.40)$$

$$\int_{-1}^1 (P_2(u))^* P_1(u) du \propto \int_{-1}^1 (u - 3u^3) du = 0 \quad (7.41)$$

$$\int_{-1}^1 (P_0(u))^* P_2(u) du \propto \int_{-1}^1 (1 - 3u^2) du = 2 - 3 \left[ \frac{u^3}{3} \right]_{-1}^1 = 2 - 3 \left( \frac{1}{3} + \frac{1}{3} \right) = 0 \quad (7.42)$$

regardless of the choice of the normalisation constant.

## 7.2 Axial symmetry

---

This is not by chance. As we will show in the next chapter, the  $P_l(u)$  are solutions of the eigenvalue problem

$$L P_l(u) = l(l+1)P_l(u) \quad L \equiv \partial_u((u^2-1)\partial_u) \quad (7.43)$$

for the *self-adjoint operator*  $L$ , and thus they are *necessarily* orthogonal (eq. 6.87). To make them orthonormal, we may just compute the normalised polynomials  $\tilde{P}_l$

$$\int_{-1}^1 (P_0(u))^* P_0(u) du = \int_{-1}^1 du = 2 \Rightarrow \tilde{P}_0(u) = \frac{1}{\sqrt{2}} \quad (7.44)$$

$$\int_{-1}^1 (P_1(u))^* P_1(u) du = \int_{-1}^1 u^2 du = \frac{2}{3} \Rightarrow \tilde{P}_1(u) = \sqrt{\frac{3}{2}} u \quad (7.45)$$

$$\begin{aligned} \int_{-1}^1 (1-3u^2)^2 du &= \int_{-1}^1 (1-6u^2+9u^4) du = 2 - 6\frac{2}{3} + 9\left[\frac{u^5}{5}\right]_{-1}^1 = \frac{8}{5} \\ \Rightarrow \tilde{P}_2(u) &= \sqrt{\frac{5}{8}}(1-3u^2) \end{aligned} \quad (7.46)$$

etc., so to obtain

$$\int_{-1}^1 (\tilde{P}_l(u))^* \tilde{P}_{l'}(u) du = \delta_{l,l'} \quad (7.47)$$

The  $P_l$  can be defined (and usually are defined) to be real, so that we may drop the complex conjugation. Changing variables back to  $\theta$ , the orthogonality relation becomes

$$\int_0^\pi \tilde{P}_l(\cos\theta) \tilde{P}_{l'}(\cos\theta) \sin\theta d\theta = \delta_{l,l'} \quad (7.48)$$

*By being eigenvectors of a self-adjoint operator, a combination (series) of these polynomials may represent each regular function in  $[-1, 1]$ , so that we will be able to express a general solution through them, as we did with the Fourier series.*

### 7.2.4 General solution

For each value of  $C = l(l+1)$  we have the solution

$$\Phi(r, \theta) = A r^l \tilde{P}_l(\cos\theta) + B r^{-l-1} \tilde{P}_l(\cos\theta) \quad (7.49)$$

As we did for the solution of the Laplace equation in polar coordinates, the boundary conditions will be completed by requiring the function to be

regular at infinity and at 0. For example, if we ask  $\Phi$  to be finite at zero, we will set  $B = 0$  since for  $l \geq 0$

$$\lim_{r \rightarrow 0} r^{-l-1} = \infty \quad (7.50)$$

so that the general solution is

$$\sum_{l=0}^{\infty} A_l r^l \tilde{P}_l(\cos \theta) \quad (7.51)$$

On the other end, if we ask  $\Phi$  to go to zero at infinity, we set  $A = 0$  and obtain the general solution

$$\sum_{l=0}^{\infty} B_l r^{-l-1} \tilde{P}_l(\cos \theta) \quad (7.52)$$

The boundary condition on the sphere at  $\bar{r}$  is (for example in the eq. 7.51 case,  $r \leq \bar{r}$ )

$$\Phi(\bar{r}, \theta) = \sum_{l=0}^{\infty} A_l \bar{r}^l \tilde{P}_l(\cos \theta) = f(\theta) \quad (7.53)$$

giving, through eq. (7.48)

$$A_l = \frac{1}{\bar{r}^l} \int_0^\pi P_l(\cos \theta) f(\theta) \sin \theta d\theta \quad (7.54)$$

In the solution is given for  $r \geq \bar{r}$  we get the same expression with

$$A_l = \bar{r}^{l+1} \int_0^\pi P_l(\cos \theta) f(\theta) \sin \theta d\theta \quad (7.55)$$

.

### 7.3 Introduction to the general case

In case  $\Phi$  depends also on  $\varphi$ , to solve

$$\frac{1}{r^2} \partial_r (r^2 \partial_r \Phi(r, \theta, \varphi)) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta \Phi(r, \theta, \varphi)) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \Phi(r, \theta, \varphi) = 0 \quad (7.56)$$

we ask

$$\Phi = R(r)\Theta(\theta)\phi(\varphi) \quad (7.57)$$

so that

$$\frac{\Theta(\theta)\phi(\varphi)}{r^2} \partial_r (r^2 \partial_r R(r)) + \frac{R(r)\phi(\varphi)}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta \Theta(\theta)) + \frac{R(r)\Theta(\theta)}{r^2 \sin^2 \theta} \partial_\varphi^2 \phi(\varphi) = 0 \quad (7.58)$$

### 7.3 Introduction to the general case

---

Let us multiply everything by

$$\frac{r^2 \sin^2 \theta}{R(r)\Theta(\theta)\phi(\varphi)} \quad (7.59)$$

to get

$$\frac{\sin^2 \theta}{R(r)} \partial_r(r^2 \partial_r R(r)) + \frac{\sin \theta}{\Theta(\theta)} \partial_\theta(\sin \theta \partial_\theta \Theta(\theta)) = -\frac{1}{\phi(\varphi)} \partial_\varphi^2 \phi(\varphi) \quad (7.60)$$

As in the best separation of variables tradition, we have an equality between functions of unrelated variable, so that we may set

$$\frac{1}{\phi(\varphi)} \partial_\varphi^2 \phi(\varphi) = C \quad (7.61)$$

$\varphi$  is an angle varying between 0 and  $2\pi$ , so that its boundary conditions are necessarily

$$\phi(\theta) = \phi(\theta + 2n\pi) \quad (7.62)$$

and we can use the results of section 2.8

$$\phi(\theta) = e^{im\phi}, \quad m \in \mathbb{Z} \quad (7.63)$$

Substituting in eq. (7.60) we get

$$\frac{\sin^2 \theta}{R(r)} \partial_r(r^2 \partial_r R(r)) + \frac{\sin \theta}{\Theta(\theta)} \partial_\theta(\sin \theta \partial_\theta \Theta(\theta)) = m^2 \quad (7.64)$$

or

$$\frac{1}{R(r)} \partial_r(r^2 \partial_r R(r)) = K = -\frac{1}{\Theta(\theta) \sin \theta} \partial_\theta(\sin \theta \partial_\theta \Theta(\theta)) + \frac{m^2}{\sin^2 \theta} \quad (7.65)$$

$K$  is again a constant, since we have on the left a function of  $r$ , and on the right a function of  $\theta$ . The radial equation,

$$\partial_r(r^2 \partial_r R(r)) = KR(r) \quad (7.66)$$

is the same we studied in section 7.2.1. We just set  $K = l(l+1)$  and write the solution as

$$R(r) = r^l \quad (7.67)$$

The  $\theta$  equation becomes

$$-\frac{1}{\sin \theta} \partial_\theta(\sin \theta \partial_\theta \Theta(\theta)) + \frac{m^2}{\sin^2 \theta} \Theta(\theta) = l(l+1)\Theta(\theta) \quad (7.68)$$

---

## Laplace equation in spherical coordinates

---

Using the same change of variables of eq. (7.16) we get

$$\begin{aligned} \frac{1}{\sin \theta} \partial_{\theta} (\sin \theta \partial_{\theta} \Theta(\theta)) &\rightarrow -\partial_u [(u^2 - 1) \partial_u \Theta(u)] \\ \frac{m^2}{\sin^2 \theta} \Theta(\theta) &\rightarrow \frac{m^2}{1 - u^2} \Theta(u) \end{aligned} \quad (7.69)$$

and thus

$$\partial_u [(u^2 - 1) \partial_u \Theta(u)] + \frac{m^2}{1 - u^2} \Theta(u) = l(l + 1) \Theta(u) \quad (7.70)$$

The solutions to this equation are called associated Legendre polynomials (see Byron-Fuller page 256)

$$P_{l,m}(\theta) \equiv (1 - u)^{\frac{m}{2}} \frac{d^m P_l(\theta)}{du^m} \quad (7.71)$$

with  $l \in \{0, \mathbb{N}\}$ ,  $m = \{-l, -l + 1, \dots, l - 1, l\}$ . The functions (spherical harmonics)

$$Y_{l,m}(\theta, \varphi) \propto P_{l,m}(\theta) e^{im\varphi} \quad (7.72)$$

i.e. the solutions of the angular part of the Laplace equation, satisfy the orthogonal relation

$$\int_0^{2\pi} d\varphi \int_0^{\pi} d\theta \sin \theta (Y_{l',m'}(\theta, \varphi))^* Y_{l,m}(\theta, \varphi) = \delta_{l,l'} \delta_{m,m'} \quad (7.73)$$

It is possible again to show that these functions are eigenvectors of a self-adjoint operator, so that a combination of these functions may be used to represent each regular function on the sphere. We thus have that a general solution for  $r \leq \bar{r}$  (see the discussion leading to eq. 7.51) is

$$\Phi(r, \theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l A_{l,m} r^l Y_{l,m}(\theta, \varphi) \quad (7.74)$$

while a general solution for  $r \geq \bar{r}$  is

$$\Phi(r, \theta, \varphi) = \sum_{l=0}^{\infty} \sum_{m=-l}^l A_{l,m} r^{-l-1} Y_{l,m}(\theta, \varphi) \quad (7.75)$$

In the former case, if a boundary condition on the sphere is given as

$$\Phi(\bar{r}, \theta, \varphi) = f(\theta, \varphi) \quad (7.76)$$

we get

$$A_{l,m} = \frac{1}{\bar{r}^l} \int_0^{2\pi} d\varphi \int_0^{\pi} d\theta \sin \theta (Y_{l,m}(\theta, \varphi))^* f(\theta, \varphi) \quad (7.77)$$

### 7.3 Introduction to the general case

---

while in the latter

$$A_{l,m} = \bar{r}^{l+1} \int_0^{2\pi} d\varphi \int_0^\pi d\theta \sin \theta (Y_{l,m}(\theta, \varphi))^* f(\theta, \varphi) \quad (7.78)$$

We will come back on the properties of spherical harmonics in the next chapters.

## Chapter 8

# Orthogonality (an invitation to Sturm-Liouville theory)

### 8.1 An other approach to orthogonality

In previous chapters we studied some orthogonal systems of functions. For example, we have seen (eqs. 2.75,2.76) that defining

$$s_n(x) = \sqrt{\frac{2}{\pi}} \sin(nx) \quad (8.1)$$

we have

$$\int_0^\pi s_n(x)s_m(x)dx = \delta_{nm} \quad (8.2)$$

Let us study these relations from a different view point. Let us compute, using only the definition of the  $s_n$

$$\begin{aligned} n^2 \int_0^\pi s_n(x)s_m(x)dx &= - \int_0^\pi \left( \frac{d^2}{dx^2} s_n(x) \right) s_m(x)dx = \\ &- \int_0^\pi \frac{d}{dx} \left( \left( \frac{d}{dx} s_n(x) \right) s_m(x) \right) dx + \int_0^\pi \left( \frac{d}{dx} s_n(x) \right) \left( \frac{d}{dx} s_m(x) \right) dx \end{aligned} \quad (8.3)$$

where we have used differentiation by parts and the trivial differentiation properties of sine. Now we may use

$$\int_0^\pi \frac{d}{dx} \left( \left( \frac{d}{dx} s_m(x) \right) s_m(x) \right) dx = \left[ \left( \frac{d}{dx} s_m(x) \right) s_m(x) \right]_0^\pi = 0 \quad (8.4)$$

(since  $s_m(0) = s_m(\pi) = 0$ ) and obtain

$$\begin{aligned} n^2 \int_0^\pi s_n(x)s_m(x)dx &= \dots = \int_0^\pi \left( \frac{d}{dx} s_n(x) \right) \left( \frac{d}{dx} s_m(x) \right) dx = \\ &- \int_0^\pi s_n(x) \left( \frac{d^2}{dx^2} s_m(x) \right) dx + \int_0^\pi \frac{d}{dx} \left( s_n(x) \left( \frac{d}{dx} s_m(x) \right) \right) dx \end{aligned} \quad (8.5)$$

## 8.1 An other approach to orthogonality

---

Again we have

$$\int_0^\pi \frac{d}{dx} \left( s_n(x) \left( \frac{d}{dx} s_m(x) \right) \right) dx = \left[ s_n(x) \left( \frac{d}{dx} s_m(x) \right) \right]_0^\pi = 0 \quad (8.6)$$

so that

$$n^2 \int_0^\pi s_n(x) s_m(x) dx = \dots = - \int_0^\pi s_n(x) \left( \frac{d^2}{dx^2} s_m(x) \right) dx = m^2 \int_0^\pi s_n(x) s_m(x) dx \quad (8.7)$$

or

$$(n^2 - m^2) \int_0^\pi s_n(x) s_m(x) dx = 0 \quad (8.8)$$

If  $n \neq m$ , we obtain

$$\int_0^\pi s_n(x) s_m(x) dx = 0 \quad (8.9)$$

i.e., we found another way to obtain eq. (2.76).

More in general, operating similarly to section 6.5.3, we may consider the operator  $L = \frac{d^2}{dx^2}$  on the space of functions  $f(x)$  such that  $f(0) = f(\pi) = 0$  and see, using multiple times differentiation by parts and the fact that functions are equal to zero in 0 and  $\pi$ , that

$$\begin{aligned} \langle f, Lg \rangle &= \int_0^\pi (f(x))^* \left( \frac{d^2}{dx^2} g(x) \right) dx = \\ &= \int_0^\pi \frac{d}{dx} \left( (f(x))^* \left( \frac{d}{dx} g(x) \right) \right) dx - \int_0^\pi \left( \frac{d}{dx} (f(x))^* \right) \left( \frac{d}{dx} g(x) \right) dx \\ &= \left[ (f(x))^* \left( \frac{d}{dx} g(x) \right) \right]_0^\pi - \int_0^\pi \left( \frac{d}{dx} (f(x))^* \right) \left( \frac{d}{dx} g(x) \right) dx = \\ &= 0 - \int_0^\pi \frac{d}{dx} \left( \left( \frac{d}{dx} (f(x))^* \right) g(x) \right) dx + \int_0^\pi \left( \frac{d^2}{dx^2} (f(x))^* \right) g(x) dx = \quad (8.10) \\ &= - \left[ \left( \frac{d}{dx} (f(x))^* \right) g(x) \right]_0^\pi + \int_0^\pi \left( \frac{d^2}{dx^2} (f(x))^* \right) g(x) dx = \\ &= 0 + \int_0^\pi \left( \frac{d^2}{dx^2} (f(x))^* \right) g(x) dx = \langle Lf, g \rangle \\ &\Rightarrow L^\dagger = L \end{aligned}$$

It is thus not surprising that the *eigenfunctions* of  $L$  are orthogonal, since eigenvectors corresponding to different eigenvalues of a self-adjoint operator are orthogonal (eq. 6.87).



## 8.2 The Sturm-Liouville problem

### 8.2.1 Definition

Let us assume that we are interested in solving the differential equation

$$\alpha(x) \frac{d^2}{dx^2} \psi(x) + \beta(x) \frac{d}{dx} \psi(x) + \gamma(x) \psi(x) = \lambda \psi(x) \quad (8.11)$$

where  $\alpha(x)$ ,  $\beta(x)$  and  $\gamma(x)$  are real functions. This problem may be considered an *eigenvalue problem*,

$$\hat{L}\psi_\lambda(x) = \lambda\psi_\lambda(x) \quad (8.12)$$

if we define the operator

$$\hat{L} \equiv \alpha(x) \frac{d^2}{dx^2} + \beta(x) \frac{d}{dx} + \gamma(x) \quad (8.13)$$

The function  $\psi(x)$  is defined on an interval  $[a, b]$  and satisfies some boundary conditions in  $a$  and  $b$  (e.g., periodic conditions, zero value, zero derivative, etc.). These properties define a functional vector space.

### 8.2.2 Scalar product

To study this problem we introduce a more general definition of scalar product

$$\langle \phi, \psi \rangle_w = \int_a^b \phi^*(x) \psi(x) w(x) dx \quad (8.14)$$

where  $w(x)$  is a real function with  $w(x) \geq 0$ , while  $\phi, \psi$  are functions in the considered vector space. We see that we have

$$\langle \psi, \phi \rangle_w = \int_a^b \psi^*(x) \phi(x) w(x) dx = \langle \phi, \psi \rangle_w^* \quad (8.15)$$

and

$$\begin{aligned} \langle \psi, \alpha\phi_1 + \beta\phi_2 \rangle_w &= \int_a^b \psi^*(x) (\alpha\phi_1(x) + \beta\phi_2(x)) w(x) dx = \\ &= \alpha \int_a^b \psi^*(x) \phi_1(x) w(x) dx + \beta \int_a^b \psi^*(x) \phi_2(x) w(x) dx = \alpha \langle \psi, \phi_1 \rangle_w + \beta \langle \psi, \phi_2 \rangle_w \end{aligned} \quad (8.16)$$

We also have

$$\langle \psi, \psi \rangle_w = \int_a^b |\psi|^2 w(x) dx \geq 0 \quad (8.17)$$

## 8.2 The Sturm-Liouville problem

---

since the integrand is positive,  $|\psi|^2 w(x) \geq 0$ .

We have thus defined a scalar product if

$$\langle \psi, \psi \rangle_w = 0 \Rightarrow \psi = 0 \quad (8.18)$$

Since the integrand is  $|\psi|^2 w(x) dx \geq 0$ ,  $\langle \psi, \psi \rangle_w$  can be zero only if the integrand is zero everywhere<sup>1</sup>.  $\psi(x)$  may thus differ from zero only where  $w(x) = 0$ . We will say thus that  $\psi = 0$  is equal to zero with respect to the measure (the weighted integral) defined by  $w$ . In the following we will introduce  $w$  in such a way to obtain some interesting results concerning the eigenvalue problem (8.12), and we will be content with this definition of  $\psi = 0$  since it will serve our purposes.

### 8.2.3 Adjoint operator

Given a linear operator  $A$ , we define the adjoint operator  $A_w^\dagger$  (although we will drop in the following the subscript  $w$ ) as the linear operator such that  $\forall \psi, \phi$  we have

$$\langle \psi, A\phi \rangle_w = \langle A_w^\dagger \psi, \phi \rangle_w \quad (8.19)$$

or

$$\int_a^b \psi^*(x) A\phi(x) w(x) dx = \int_a^b \left( A_w^\dagger \psi(x) \right)^* \phi(x) w(x) dx \quad (8.20)$$

### 8.2.4 Defining $w$ so that $\hat{L}$ is self-adjoint

We know that self-adjoint operators have interesting properties, between them the fact that their eigenvectors are orthogonal, and may be used to define an orthonormal basis. We will now show that for the Sturm-Liouville problem (8.11) we may define a suitable  $w(x)$  such that the operator  $\hat{L}$  is self-adjoint with respect to  $\langle \cdot, \cdot \rangle_w$ .

Let us compute

$$\begin{aligned} \int_a^b \psi^*(x) \hat{L}\phi(x) w(x) dx &= \int_a^b \psi(x)^* \alpha(x) \left( \frac{d^2}{dx^2} \phi(x) \right) w(x) dx + \\ \int_a^b \psi(x)^* \beta(x) \left( \frac{d}{dx} \phi(x) \right) w(x) dx &+ \int_a^b \psi(x)^* \gamma(x) \phi(x) w(x) dx = (*) \end{aligned} \quad (8.21)$$

---

<sup>1</sup>More properly, different from zero on a set of zero measure.

We now use derivation by parts

$$\begin{aligned}
 (*) &= \int_a^b \frac{d}{dx} \left[ \psi(x)^* \alpha(x) \left( \frac{d}{dx} \phi(x) \right) w(x) \right] dx - \int_a^b \frac{d}{dx} (\psi(x)^* \alpha(x) w(x)) \left( \frac{d}{dx} \phi(x) \right) dx + \\
 &\int_a^b \frac{d}{dx} [\psi(x)^* \beta(x) \phi(x) w(x)] dx - \int_a^b \phi(x) \frac{d}{dx} (\psi(x)^* \beta(x) w(x)) dx + \int_a^b \psi(x)^* \gamma(x) \phi(x) w(x) dx = (**)
 \end{aligned}
 \tag{8.22}$$

and

$$\begin{aligned}
 (**) &= \left[ \psi(x)^* \alpha(x) \left( \frac{d}{dx} \phi(x) \right) w(x) \right]_a^b + [\psi(x)^* \beta(x) \phi(x) w(x)]_a^b \\
 &- \int_a^b \left( \frac{d}{dx} \psi(x)^* \right) \left( \frac{d}{dx} \phi(x) \right) \alpha(x) w(x) dx - \int_a^b (\psi(x)^*) \frac{d}{dx} (\alpha(x) w(x)) \left( \frac{d}{dx} \phi(x) \right) dx \\
 &- \int_a^b \left( \frac{d}{dx} \psi(x)^* \right) \beta(x) w(x) \phi(x) dx - \int_a^b \psi(x)^* \frac{d}{dx} (\beta(x) w(x)) \phi(x) dx \\
 &+ \int_a^b \psi(x)^* \gamma(x) \phi(x) w(x) dx = (***)
 \end{aligned}
 \tag{8.23}$$

If one of the following holds

1.  $\psi$  and  $\phi$  are 0 in  $a$  and  $b$
2. the derivatives of  $\psi$  and  $\phi$  are 0 in  $a$  and  $b$
3.  $\psi$ ,  $\phi$ ,  $\alpha$  and  $w$  are periodic in  $[a, b]$
4.  $w$  is 0 in  $a$  and  $b$
5.  $\alpha$  is 0 in  $a$  and  $b$

we have for the first term

$$\left[ \psi(x)^* \alpha(x) \left( \frac{d}{dx} \phi(x) \right) w(x) \right]_a^b = 0
 \tag{8.24}$$

Let us assume this to be true, and get

$$\begin{aligned}
 \langle \psi, \hat{L}\phi \rangle_w &= \int_a^b \psi(x)^* \hat{L}\phi(x) w(x) dx = [\psi(x)^* \beta(x) \phi(x) w(x)]_a^b \\
 &- \int_a^b \left( \frac{d}{dx} \psi(x)^* \right) \left( \frac{d}{dx} \phi(x) \right) \alpha(x) w(x) dx - \int_a^b \psi(x)^* \frac{d}{dx} (\alpha(x) w(x)) \left( \frac{d}{dx} \phi(x) \right) dx \\
 &- \int_a^b \left( \frac{d}{dx} \psi(x)^* \right) \beta(x) w(x) \phi(x) dx - \int_a^b \psi(x)^* \frac{d}{dx} (\beta(x) w(x)) \phi(x) dx \\
 &+ \int_a^b \psi(x)^* \gamma(x) \phi(x) w(x) dx
 \end{aligned}
 \tag{8.25}$$

## 8.2 The Sturm-Liouville problem

---

Let us now compute

$$\begin{aligned} \langle \hat{L}\psi, \phi \rangle_w &= \int_a^b \left( \hat{L}\psi(x) \right)^* \phi(x) w(x) dx = \int_a^b \phi(x) \alpha(x) \left( \frac{d^2}{dx^2} \psi(x) \right)^* w(x) dx + \\ &\int_a^b \phi(x) \beta(x) \left( \frac{d}{dx} \phi(x) \right)^* w(x) dx + \int_a^b \psi(x)^* \gamma(x) \phi(x) w(x) dx \end{aligned} \quad (8.26)$$

The only difference with the previous case is that now derivatives are acting on the complex conjugate of  $\psi$  (due to linearity, the complex conjugate of the derivative is the derivative of the complex conjugate). As a consequence, also in the result the only change will be in where the derivative acts, and thus

$$\begin{aligned} \langle \hat{L}\psi, \phi \rangle_w &= \int_a^b (L\psi(x))^* \phi(x) w(x) dx = [\psi(x)^* \beta(x) \phi(x) w(x)]_a^b \\ &- \int_a^b \left( \frac{d}{dx} \psi(x)^* \right) \left( \frac{d}{dx} \phi(x) \right) \alpha(x) w(x) dx - \int_a^b \left( \frac{d}{dx} \psi(x) \right)^* \frac{d}{dx} (\alpha(x) w(x)) \phi(x) dx \\ &- \int_a^b \psi(x)^* \beta(x) w(x) \left( \frac{d}{dx} \phi(x) \right) dx - \int_a^b \psi(x)^* \frac{d}{dx} (\beta(x) w(x)) \phi(x) dx \\ &+ \int_a^b \psi(x)^* \gamma(x) \phi(x) w(x) dx \end{aligned} \quad (8.27)$$

A few terms are left unchanged, so that

$$\begin{aligned} \langle \hat{L}\psi, \phi \rangle_w - \langle \psi, \hat{L}\phi \rangle_w &= \\ &- \int_a^b \left( \frac{d}{dx} \psi(x) \right)^* \frac{d}{dx} (\alpha(x) w(x)) \phi(x) dx + \int_a^b \psi(x)^* \frac{d}{dx} (\alpha(x) w(x)) \left( \frac{d}{dx} \phi(x) \right) dx \\ &- \int_a^b \psi(x)^* \beta(x) w(x) \left( \frac{d}{dx} \phi(x) \right) dx + \int_a^b \left( \frac{d}{dx} \psi(x)^* \right) \beta(x) w(x) \phi(x) dx \end{aligned} \quad (8.28)$$

or

$$\begin{aligned} \langle \hat{L}\psi, \phi \rangle_w - \langle \psi, \hat{L}\phi \rangle_w &= \\ &\int_a^b \left( \frac{d}{dx} \psi(x) \right)^* \phi(x) \left[ \beta(x) w(x) - \frac{d}{dx} (\alpha(x) w(x)) \right] dx \\ &+ \int_a^b \psi(x)^* \left( \frac{d}{dx} \phi(x) \right) \left[ \frac{d}{dx} (\alpha(x) w(x)) - \beta(x) w(x) \right] dx \end{aligned} \quad (8.29)$$

In order for this term to be zero, and thus, under the hypothesis (8.24), for having  $L^\dagger = L$ , we need to choose  $w$  so that

$$\frac{d}{dx} (\alpha(x) w(x)) = \beta(x) w(x) \quad (8.30)$$

We may try to solve this latter differential equation in order to obtain  $w(x)$ . We use again partial differentiation

$$\left(\frac{d}{dx}\alpha(x)\right)w(x) + \left(\frac{d}{dx}w(x)\right)\alpha(x) = \beta(x)w(x) \quad (8.31)$$

and we rearrange it as

$$\frac{d}{dx}w(x) = w(x)\frac{\beta(x) - \frac{d}{dx}\alpha(x)}{\alpha(x)} \quad (8.32)$$

Let us define

$$g(x) \equiv \frac{\beta(x) - \alpha'(x)}{\alpha(x)} \quad (8.33)$$

(the prime being a short-hand for differentiation with respect to  $x$ ) which requires that  $\alpha(x) \neq 0$ , and obtain

$$\frac{d}{dx}w(x) = g(x)w(x) \quad (8.34)$$

This equation is solved by

$$w(x) = C \exp \left[ \int_a^x g(\tau) d\tau \right] \quad (8.35)$$

since

$$\frac{d}{dx} \left( C \exp \left[ \int_a^x g(\tau) d\tau \right] \right) = g(x) C \exp \left[ \int_a^x g(\tau) d\tau \right] = g(x)w(x) \quad (8.36)$$

We thus have found an expression of  $w(x)$  that gives us  $\hat{L}^\dagger = \hat{L}$ .

### 8.2.5 A different formulation of the problem

We do not necessarily need to perform an integration to find  $w$ . Let us write

$$\frac{1}{w(x)} \left( \frac{d}{dx} \left( w(x)\alpha(x) \frac{d}{dx}\psi(x) \right) \right) + \gamma(x)\psi(x) = \lambda\psi(x) \quad (8.37)$$

Using

$$\frac{d}{dx} (\alpha(x)w(x)) = \beta(x)w(x) \quad (8.38)$$

and partial differentiation we have

$$\frac{1}{w(x)}w(x)\alpha(x) \left( \frac{d^2}{dx^2}\psi(x) \right) + \frac{1}{w(x)} \left( \frac{d}{dx} (w(x)\alpha(x)) \right) \frac{d}{dx}\psi(x) + \gamma(x)\psi(x) = \lambda\psi(x) \quad (8.39)$$

## 8.2 The Sturm-Liouville problem

---

from which we get

$$\frac{1}{w(x)}w(x)\alpha(x)\left(\frac{d^2}{dx^2}\psi(x)\right) + \frac{1}{w(x)}\beta(x)w(x)\frac{d}{dx}\psi(x) + \gamma(x)\psi(x) = \lambda\psi(x) \quad (8.40)$$

which is equivalent to our original problem

$$\alpha(x)\left(\frac{d^2}{dx^2}\psi(x)\right) + \beta(x)\frac{d}{dx}\psi(x) + \gamma(x)\psi(x) = \lambda\psi(x) \quad (8.41)$$

This means that if we define

$$\eta(x) = w(x)\alpha(x) \quad (8.42)$$

the differential problem

$$\frac{1}{w(x)}\left(\frac{d}{dx}\left(\eta(x)\frac{d}{dx}\psi(x)\right)\right) + \gamma(x)\psi(x) = \lambda\psi(x) \quad (8.43)$$

is equivalent to the Sturm-Liouville one.

### 8.2.6 Orthogonality

If now we have two eigenfunctions of our Sturm-Liouville problem, i.e

$$\hat{L}\psi_\lambda(x) = \lambda\psi_\lambda(x), \quad \hat{L}\psi_{\lambda'}(x) = \lambda'\psi_{\lambda'}(x) \quad (8.44)$$

we may easily verify (repeating our general analysis of section 6.3.5, and using the fact that  $\langle \cdot, \cdot \rangle_w$  satisfies all the scalar product properties) that

$$\begin{aligned} \lambda\langle\psi_\lambda, \psi_\lambda\rangle_w &= \langle\psi_\lambda, \lambda\psi_\lambda\rangle_w = \langle\psi_\lambda, \hat{L}\psi_\lambda\rangle_w = \\ \langle\hat{L}^\dagger\psi_\lambda, \psi_\lambda\rangle_w &= \langle\hat{L}\psi_\lambda, \psi_\lambda\rangle_w = \langle\lambda\psi_\lambda, \psi_\lambda\rangle_w = \lambda^*\langle\psi_\lambda, \psi_\lambda\rangle_w \\ \Rightarrow \lambda &= \lambda^* \end{aligned} \quad (8.45)$$

assuring us that all eigenvalues are real. Furthermore, using this last result

$$\begin{aligned} \lambda\langle\psi_{\lambda'}, \psi_\lambda\rangle_w &= \langle\psi_{\lambda'}, \lambda\psi_\lambda\rangle_w = \langle\psi_{\lambda'}, \hat{L}\psi_\lambda\rangle_w = \\ \langle\hat{L}^\dagger\psi_{\lambda'}, \psi_\lambda\rangle_w &= \langle\hat{L}\psi_{\lambda'}, \psi_\lambda\rangle_w = \langle\lambda'\psi_{\lambda'}, \psi_\lambda\rangle_w = \lambda'\langle\psi_{\lambda'}, \psi_\lambda\rangle_w \\ \Rightarrow (\lambda - \lambda')\langle\psi_{\lambda'}, \psi_\lambda\rangle_w &= 0 \end{aligned} \quad (8.46)$$

From this follows that two eigenfunctions corresponding to different eigenvalues,  $\lambda \neq \lambda'$ , are orthogonal

$$\langle\psi_{\lambda'}, \psi_\lambda\rangle_w = 0 \quad (8.47)$$

The task of showing, for an infinite dimensional vector space, that the eigenfunctions are a basis for the space is a non-trivial problem in functional analysis, that we will just assume to be true.

### 8.3 Application to Legendre polynomials

In our study of the Laplace equation in spherical coordinates we studied (eq. 7.12) the equation

$$\frac{1}{\sin \theta} \frac{d}{d\theta} \left( \sin \theta \frac{d}{d\theta} \Theta(\theta) \right) = \lambda \Theta(\theta), \quad \theta \in [0, \pi] \quad (8.48)$$

By comparison with eq. (8.43) we see that this is a Sturm-Liouville problem with

$$w(\theta) = \sin \theta, \quad \eta(\theta) = \alpha(\theta)w(\theta) = \alpha(\theta) \sin \theta = \sin \theta \Rightarrow \alpha(\theta) = 1 \quad (8.49)$$

We see that  $\alpha \neq 0$ , and  $w$  assumes zero value in 0 and  $\pi$  (and it is positive in  $(0, \pi)$ ), so that all the requirements for the Sturm-Liouville theory are satisfied. This means that the eigenvalues  $\lambda$  are necessarily real, and<sup>2</sup>

$$\int_0^\pi \Theta_\lambda(\theta) \Theta_{\lambda'} \sin \theta d\theta = \delta_{\lambda, \lambda'} \quad (8.50)$$

The factor  $\sin \theta$  in the integral comes from the weight  $w(\theta)$ . It is extremely useful, since we want our eigenfunctions to be orthogonal when integrated on the sphere, i.e. with integration volume  $dx^3 = r^2 \sin \theta dr d\theta d\varphi$ .

---

<sup>2</sup>Assuming the functions  $\Theta$  to be defined as real, something that we know to be possible from our previous study of Legendre polynomials

---

### 8.3 Application to Legendre polynomials

---



## Chapter 9

# The hydrogen atom

### 9.1 Introduction

In chapter 3, we discussed how the Austrian physicist Erwin Schrödinger introduced a partial differential equation to study the motion of extremely small particles, such as electrons (Quantum Mechanics). We then applied the theory to compute the possible energies of particles trapped in a potential box.

That was a simple problem, very suitable to a first application of the theory, but it wasn't obviously the problem that Schrödinger had developed his theory for. Has he wrote down his equation, he tried to apply it to a real natural system, one of the simplest and more common physical systems in nature: the hydrogen atom.

The hydrogen atom is composed of two particles, a proton and an electron, with opposite charge. It is similar to a small planetary system, in which the heavier proton plays the role of the Sun, and the lighter electron the role of the Earth. We will assume that the electron and proton are point particles without inner structure (this is true, as far as we know, for the electron, while the proton is composed of quarks, held together by a force, the Strong Force of Quantum Chromodynamics, whose mathematical description is extremely complex and still not fully understood), that their dynamics may be described by the non-relativistic Schrödinger equation (quantum generalisation of Newtonian dynamics), without taking in account relativistic effects (Dirac equation, quantum generalisation of Einstein dynamics), and that the electric field giving the interaction between the two particles can be described by the classical Coulomb force, without taking in account the quantum nature of the field itself.

These assumptions make the study of the system *extremely* simpler.

## 9.2 The equation

---

They are nevertheless good approximations when studying the hydrogen atom<sup>1</sup>, so that when Schrödinger applied his theory to the hydrogen atom, was extremely satisfied of the agreement between theory and observations<sup>2</sup>.

The assumption makes the study of the system simpler, but we will still need some effort to arrive at a solution. The study of the hydrogen atom will represent the highest hurdle in our study of separation of variable methods, and will lead us to introduce some new techniques, as for example the algebraic treatment of angular momentum. It may be interesting to see how hard is to find a solution to a simplified version of one of Nature's simplest systems. It gives the perspective that maybe some of the extremely simple and straightforward solutions proposed for real world systems in economy, social systems, etc., may not be as comprehensive as their proponents like to state.

## 9.2 The equation

Let us first assume that, by being much more massive<sup>3</sup>, the proton may be considered as fixed<sup>4</sup>. The electron moves then in a Coulomb potential

$$\Phi(r) = \frac{e}{r} \quad (9.1)$$

$\mathbf{r}$  being the position of the electron with respect to the proton (located in the origin), and  $e$  being the charge of the proton. Since the charge of the electron is  $-e$ , the (classical) energy of the electron is given as a function of its velocity  $\mathbf{v}$  and position  $\mathbf{r}$  as

$$E = \frac{1}{2}mv^2 - \frac{e^2}{r} \quad (9.2)$$

---

<sup>1</sup>For example, the energy of the fundamental (lowest) state of the system, is negligible with respect to the energy that corresponds to the mass of the electron corresponding to the celebrated  $E = mc^2$  equation, and thus relativistic effects may be ignored in a first computation

<sup>2</sup>Obviously, the Dirac equation, Quantum Electrodynamics (quantum treatment of the electromagnetic field] and Quantum Chromodynamics where developed after Schrödinger's theory. Nevertheless, Schrödinger first tried to develop an equation taking in account Relativistic effects, but the equation did not agree with observations, and he decided not to publish it. The equation is nowadays known as the Klein-Gordon equation, from the names of the physicists that re-discovered it. The equation describes a particle without spin, and thus is not apt to describe the relativistic dynamics of the electron, as the Dirac equation thus

<sup>3</sup>The ratio between the proton and electron mass is  $\approx 1836$ .

<sup>4</sup>This approximation is studied in appendix 9.B (you may want to read first appendix 9.A).

$m$  being the electron mass <sup>5</sup>. Remembering that Schrödinger's theory connects the energy to time derivative, and the momentum to space derivative, we recall the classical relation

$$\mathbf{p} = m\mathbf{v} \quad (9.3)$$

and obtain

$$E = \frac{p^2}{2m} - \frac{e^2}{r} \quad (9.4)$$

Now we follow Schrödinger's recipe for a quantum equation, namely we replace the energy with a time derivative (multiplied by  $i\hbar$ ,  $i$  being the imaginary unit and  $\hbar$  the Plank constant divided by  $2\pi$ ), acting on the wave function  $\Psi(t, \mathbf{x})$

$$E \rightarrow i\hbar\partial_t \Psi(t, \mathbf{x}) \quad (9.5)$$

We then replace each component of the momentum  $p_j$  with  $-i\hbar\partial_j$ , or

$$\frac{p^2}{2m} \rightarrow -\frac{\hbar^2}{2m}\nabla^2\Psi(t, \mathbf{x}) \quad (9.6)$$

and finally replace the potential with a multiplicative operator

$$\frac{e^2}{r} \rightarrow \frac{e^2}{r}\Psi(t, \mathbf{x}) \quad (9.7)$$

The Schrödinger equation for the hydrogen atom is then

$$i\hbar\partial_t\Psi(t, \mathbf{x}) = -\frac{\hbar^2}{2m}\nabla^2\Psi(t, \mathbf{x}) - \frac{e^2}{r}\Psi(t, \mathbf{x}) \quad (9.8)$$

Since the potential depends only on  $r$ , is better to use spherical coordinates, i.e. (refer to eq. 1.83)

$$i\hbar\partial_t\Psi(t, r, \theta, \varphi) = -\frac{\hbar^2}{2m}\left[\frac{1}{r^2}\partial_r(r^2\partial_r) + \frac{1}{r^2\sin\theta}\partial_\theta(\sin\theta\partial_\theta) + \frac{1}{r^2\sin^2\theta}\partial_\varphi^2\right]\Psi(t, r, \theta, \varphi) - \frac{e^2}{r}\Psi(t, r, \theta, \varphi) \quad (9.9)$$

### 9.3 Time independent Schrödinger equation

Let us use as usual the separation of variables technique and assume

$$\Psi(t, r, \theta, \varphi) = T(t)\psi(r, \theta, \varphi) \quad (9.10)$$

---

<sup>5</sup>We are using the definition of charge that allows us to write potentials and forces without having constants such as  $\epsilon_0$  appear. See section 9.5.2

## 9.4 The angular solution

---

We get

$$\begin{aligned} \psi(r, \theta, \varphi) i\hbar T(t) = \\ - T(t) \left( \frac{\hbar^2}{2m} \left[ \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \right] \psi(r, \theta, \varphi) - \frac{e^2}{r} \psi(r, \theta, \varphi) \right) \end{aligned} \quad (9.11)$$

We divide each side by  $\Psi$  and obtain

$$\begin{aligned} \frac{1}{T(t)} i\hbar T(t) = E = \\ - \frac{1}{\psi(r, \theta, \varphi)} \left( \frac{\hbar^2}{2m} \left[ \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \right] \psi(r, \theta, \varphi) - \frac{e^2}{r} \psi(r, \theta, \varphi) \right) \end{aligned} \quad (9.12)$$

where  $E$  is a constant (the energy of the time independent solution), due as usual to the fact that we obtained functions of independent variables on the two sides of the equality.

The equation

$$i\hbar T(t) = ET(t) \quad (9.13)$$

is trivially solved by

$$T(t) = e^{-i\frac{E}{\hbar}t} \quad (9.14)$$

The corresponding spatial solution, that we may call  $\psi_E$ , is given by the *time independent Schrödinger equation*

$$\begin{aligned} - \left( \frac{\hbar^2}{2m} \left[ \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \right] - \frac{e^2}{r} \right) \psi_E(r, \theta, \varphi) = \\ E\psi_E(r, \theta, \varphi) \end{aligned} \quad (9.15)$$

## 9.4 The angular solution

We already know (see eq. 7.72) functions (spherical harmonics)  $Y_{l,m}(\theta, \varphi)$  such that

$$- \left( \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\varphi^2 \right) Y_{l,m}(\theta, \varphi) = l(l+1)Y_{l,m}(\theta, \varphi) \quad (9.16)$$

where

$$Y_{l,m}(\theta, \varphi) \propto P_{l,m}(\theta) e^{im\varphi} \quad (9.17)$$

$P_{l,m}(\theta)$  being an associated Legendre polynomial with  $l \in \{0, \mathbb{N}\}$ ,  $m = \{-l, -l+1, \dots, l-1, l\}$ , so that we can just assume

$$\psi_E(r, \theta, \varphi) = R_E(r) Y_{l,m}(\theta, \varphi) \quad (9.18)$$

and obtain an equation for  $R_E(r)$ . It is nevertheless useful to use algebraic techniques common in Quantum Mechanics to study the relation between the spherical harmonics and angular momentum, and explicitly show the properties of spherical harmonics. This is done in appendix 9.C (you may need to read first appendix 9.A).

## 9.5 The radial equation

Substituting eq. (9.18) in eq. (9.15) we obtain

$$-\frac{\hbar^2}{2m} \frac{1}{r^2} \partial_r (r^2 \partial_r) R_E(r) + \frac{\hbar^2 l(l+1)}{2mr^2} R_E(r) - \frac{e^2}{r} R_E(r) = E R_E(r) \quad (9.19)$$

where  $l$  is an integer (possibly also 0, see appendix 9.C).

### 9.5.1 First change of variable

Let us now define

$$\xi(r) = r R_E(r) \Rightarrow R(r) = \frac{\xi(r)}{r} \quad (9.20)$$

(we dropped the suffix  $E$  in  $\xi$  but should be clear that the solution is found for given values of  $E$  (and  $l$ )).

Since the radial equation contains the term  $\partial_r (r^2 \partial_r R_E(r)) / r^2$  let us first compute

$$r^2 \left( \partial_r \left( \frac{\xi(r)}{r} \right) \right) = r^2 \left( \frac{1}{r} \partial_r \xi(r) - \frac{1}{r^2} \xi(r) \right) = r \partial_r \xi(r) - \xi(r) \quad (9.21)$$

and then

$$\frac{1}{r^2} \partial_r \left( r^2 \left( \partial_r \left( \frac{\xi(r)}{r} \right) \right) \right) = \frac{1}{r^2} \partial_r (r \partial_r \xi(r) - \xi(r)) = \frac{1}{r^2} \partial_r \xi(r) + \frac{1}{r} \partial_r^2 \xi(r) - \frac{1}{r^2} \partial_r \xi(r) = \frac{1}{r} \partial_r^2 \xi(r) \quad (9.22)$$

The equation becomes then

$$-\frac{\hbar^2}{2m} \frac{1}{r} \partial_r^2 \xi(r) + \frac{\hbar^2 l(l+1)}{2mr^3} \xi(r) - \frac{e^2}{r^2} \xi(r) = E \frac{\xi(r)}{r} \quad (9.23)$$

Multiplying by  $r$  we get

$$\frac{\hbar^2}{2m} \partial_r^2 \xi(r) - \frac{\hbar^2 l(l+1)}{2mr^2} \xi(r) + \frac{e^2}{r} \xi(r) + E \xi(r) = 0 \quad (9.24)$$

or

$$\partial_r^2 \xi(r) + \frac{2m}{\hbar^2} \left( -\frac{\hbar^2 l(l+1)}{2mr^2} + \frac{e^2}{r} + E \right) \xi(r) = 0 \quad (9.25)$$

## 9.5 The radial equation

---

Our change of variables led thus to an equation in which only a second derivative and a multiplicative term dependent on  $r$  appear. We may hope this equation to be solved more easily.

For a reason that will be clear soon, we re-write again the equation as

$$\partial_r^2 \xi(r) + \frac{2me^2}{\hbar^2} \left( -\frac{\hbar^2 l(l+1)}{2mr^2 e^2} + \frac{1}{r} + \frac{E}{e^2} \right) \xi(r) = 0 \quad (9.26)$$

### 9.5.2 Dimensional analysis (characteristic length)

We wrote our potential as

$$\frac{e^2}{r} \quad (9.27)$$

You may be more familiar with

$$\frac{e^2}{4\pi \epsilon_0 r} \quad (9.28)$$

Our choice corresponds to a definition of the charge that makes dimensional analysis easier (since we do not introduce a new “charge” dimensionality, but define charge based on mass, length and time). The magnitude of the electrostatic force is

$$F = \frac{e^2}{r^2} \quad (9.29)$$

so that

$$\frac{[M][L]}{[T]^2} = [F] = \frac{[e]^2}{[L]^2} \quad (9.30)$$

or

$$[e]^2 = \frac{[M][L]^3}{[T]^2} \left( \Rightarrow [e] = \frac{[M]^{\frac{1}{2}}[L]^{\frac{3}{2}}}{[T]} \right) \quad (9.31)$$

Recalling that

$$[\hbar] = [E][T] = \frac{[M][L]^2}{[T]} \quad (9.32)$$

we verify that the following quantity is a length

$$\left[ \frac{\hbar}{e^2 m} \right] = \frac{[M]^2 [L]^4}{[T]^2} [M]^{-1} \frac{[T]^2}{[M][L]^3} = [L] \quad (9.33)$$

We may thus define the *Bohr radius*<sup>6</sup>

$$r_0 = \frac{\hbar}{e^2 m} \approx 5.310^{-11} \text{m} \quad (9.34)$$

Using this quantity, the equation becomes

$$\partial_r^2 \xi(r) + \frac{2}{r_0} \left( -\frac{l(l+1)}{2r^2} r_0 + \frac{1}{r} + \frac{E}{e^2} \right) \xi(r) = 0 \quad (9.35)$$

---

<sup>6</sup>Introduced in the first quantum theory of the atom by the Danish Nobel prize winner Niels Bohr.

### 9.5.3 Second change of variables

Since the equation itself is suggesting us a natural unit (a quantity with the dimension of length that appears in the equation) let us use it to measure lengths, and perform the change of variables

$$u = \frac{r}{r_0}, \Rightarrow r = r_0 u \quad (9.36)$$

$$\partial_r = \partial_u \frac{\partial u}{\partial r} = \frac{1}{r_0} \partial_u \Rightarrow \partial_r^2 = \frac{1}{r_0^2} \partial_u^2 \quad (9.37)$$

Under this change of variable, the differential equation becomes

$$\frac{1}{r_0^2} \partial_u^2 \xi(u) + \frac{2}{r_0} \left( -\frac{l(l+1)}{2r_0^2 u^2} r_0 + \frac{1}{r_0 u} + \frac{E}{e^2} \right) \xi(u) = 0 \quad (9.38)$$

or, multiplying by  $r_0^2$

$$\partial_u^2 \xi(u) + \left( -\frac{l(l+1)}{u^2} + \frac{2}{u} + \frac{r_0}{2e^2} E \right) \xi(u) = 0 \quad (9.39)$$

### 9.5.4 A further dimensional analysis (characteristic energy)

We may now note that

$$\left[ \frac{e^2}{r_0} \right] = \frac{[M][L]^3}{[T]^2[L]} = \frac{[M][L]^2}{[T]^2} = [E] \quad (9.40)$$

Let us call the characteristic energy

$$E_0 \equiv \frac{e^2}{2r_0} \quad (9.41)$$

and our differential equation becomes

$$\partial_u^2 \xi(u) + \left( -\frac{l(l+1)}{u^2} + \frac{2}{u} + \frac{E}{E_0} \right) \xi(u) = 0 \quad (9.42)$$

We are interested in studying electrons *in the atom*, meaning those that are bound to it. “Bound” means that the electron cannot reach, unless external energy is provided to it, an arbitrary distance from the proton. At this arbitrary (infinity) distance, the potential due to the proton is zero, so that the (classical) energy would be

$$\frac{p^2}{2m} \geq 0 \quad (9.43)$$

Since the electron (again, from a classical viewpoint) can go everywhere the conservation of energy law allows it to go, if it had  $E \geq 0$  it could go at

## 9.5 The radial equation

---

arbitrary distance. We thus assume, for bound electrons,  $E < 0$ . Since  $E_0 > 0$ , we write

$$n \equiv \sqrt{-\frac{E_0}{E}} > 0, \Rightarrow \frac{E}{E_0} = -\frac{1}{n^2}, \Rightarrow E = -\frac{E_0}{n^2} \quad (9.44)$$

At the moment,  $n$  is just a new name for a quantity that we know to be positive (but later we will show that  $n$  needs to be an integer).

The differential equation is now

$$\partial_u^2 \xi(u) + \left( -\frac{l(l+1)}{u^2} + \frac{2}{u} - \frac{1}{n^2} \right) \xi(u) = 0 \quad (9.45)$$

### 9.5.5 Asymptotic behaviour

The equation has simplified a lot, but in order to solve it we should better understand how it behaves for large and small values of  $r$  (far or close to the proton) or, equivalently, for large and small values of  $u$

#### Large $u$

For  $u \rightarrow \infty$ , we have

$$\frac{1}{u^2} \rightarrow 0, \quad \frac{1}{u} \rightarrow 0, \quad \frac{1}{n^2} = \text{const} \quad (9.46)$$

so that we may approximate the equation for the behaviour far from the proton as

$$\partial_u^2 \xi_{\text{far}}(u) = \frac{1}{n^2} \xi_{\text{far}}(u) \quad (9.47)$$

This is easily solved as

$$\xi_{\text{far}}(u) = e^{\pm \frac{u}{n}} \quad (9.48)$$

Now, since we have  $n > 0$  and  $u > 0$ , the only non-diverging solution is

$$\xi_{\text{far}}(u) = e^{-\frac{u}{n}} \quad (9.49)$$

#### Small $u$

For  $u \rightarrow 0$ , the term  $u^{-2}$  diverges faster than the others

$$\frac{1}{u^2} + \frac{1}{u} + c = \frac{1}{u^2} (1 + u + cu^2) \rightarrow \frac{1}{u^2} \quad (9.50)$$

so that we may approximate the equation for the behaviour far from the proton as

$$\partial_u^2 \xi_{\text{near}}(u) = \frac{l(l+1)}{u^2} \xi_{\text{near}}(u) \quad (9.51)$$



We try a solution as

$$\xi_{\text{near}}(u) = u^k \Rightarrow \partial_u \xi_{\text{near}}(u) = k u^{k-1} \Rightarrow \partial_u^2 \xi_{\text{near}}(u) = k(k-1) u^{k-2} \quad (9.52)$$

so that eq. (9.51) becomes

$$k(k-1) u^{k-2} = l(l+1) u^{k-2} \Rightarrow k(k-1) = l(l+1) \quad (9.53)$$

This equation has two solutions

$$k = l + 1, \quad k = -l \quad (9.54)$$

but, again, since  $l \geq 0$  in order for  $u^k$  not to diverge for  $u \rightarrow 0$ , we have<sup>7</sup>

$$\xi_{\text{near}}(u) = u^{l+1} \quad (9.55)$$

### The ansatz for the solution

The behaviour for small and large  $u$  may suggest us the following *ansatz* (guess) for  $\xi(u)$

$$\xi(u) = \xi_{\text{far}}(u) f(u) \xi_{\text{near}}(u) = u^{l+1} e^{-\frac{u}{n}} f(u) \quad (9.56)$$

To see if this assumption is reasonable, we have to insert it in the equation and see if we find a solution for  $f(u)$ .

#### 9.5.6 The equation for the function $f(u)$

Let us compute

$$\begin{aligned} \partial_u \xi(u) &= (l+1) u^l e^{-\frac{u}{n}} f(u) + u^{l+1} \left( -\frac{1}{n} \right) e^{-\frac{u}{n}} f(u) + u^{l+1} e^{-\frac{u}{n}} \partial_u f(u) = \\ u^{l+1} e^{-\frac{u}{n}} f(u) &\left( \frac{l+1}{u} - \frac{1}{n} + \frac{\partial_u f(u)}{f(u)} \right) = \xi(u) \left( \frac{l+1}{u} - \frac{1}{n} + \frac{\partial_u f(u)}{f(u)} \right) \end{aligned} \quad (9.57)$$

and

$$\begin{aligned} \partial_u^2 \xi(u) &= (\partial_u \xi(u)) \left( \frac{l+1}{u} - \frac{1}{n} + \frac{\partial_u f(u)}{f(u)} \right) + \xi(u) \partial_u \left( \frac{l+1}{u} - \frac{1}{n} + \frac{\partial_u f(u)}{f(u)} \right) = \\ \xi(u) &\left( \frac{l+1}{u} - \frac{1}{n} + \frac{\partial_u f(u)}{f(u)} \right)^2 + \xi(u) \left( -\frac{l+1}{u^2} + \frac{\partial_u^2 f(u)}{f(u)} - \frac{(\partial_u f(u))^2}{f(u)^2} \right) = \\ \xi(u) &\left( \frac{(l+1)^2}{u^2} + \frac{1}{n^2} + \frac{(\partial_u f(u))^2}{f(u)^2} - \frac{2(l+1)}{nu} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} - \frac{l+1}{u^2} + \frac{\partial_u^2 f(u)}{f(u)} - \frac{(\partial_u f(u))^2}{f(u)^2} \right) \\ &= \xi(u) \left( \frac{(l+1)^2}{u^2} + \frac{1}{n^2} - \frac{2(l+1)}{nu} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} - \frac{l+1}{u^2} + \frac{\partial_u^2 f(u)}{f(u)} \right) \end{aligned} \quad (9.58)$$

---

<sup>7</sup>Strictly speaking, for  $l = 0$  also  $k = -l$  is not diverging. Anyway, this process is more *guiding* us to the ansatz (9.56) than a strict derivation.

## 9.5 The radial equation

---

Now we note that

$$(l+1)^2 = (l+1)(l+1) = (l+1) + l(l+1) \quad (9.59)$$

so that

$$\begin{aligned} \partial_u^2 \xi(u) &= \xi(u) \left( \frac{l+1}{u^2} + \frac{l(l+1)}{u^2} - \frac{l+1}{u^2} - \frac{2(l+1)}{nu} + \frac{1}{n^2} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} \right. \\ &\quad \left. + \frac{\partial_u^2 f(u)}{f(u)} \right) = \xi(u) \left( \frac{l(l+1)}{u^2} - \frac{2(l+1)}{nu} + \frac{1}{n^2} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} + \frac{\partial_u^2 f(u)}{f(u)} \right) \end{aligned} \quad (9.60)$$

Substituting this in eq. (9.45) we get

$$\begin{aligned} &\xi(u) \left( \frac{l(l+1)}{u^2} - \frac{2(l+1)}{nu} + \frac{1}{n^2} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} \right. \\ &\quad \left. + \frac{\partial_u^2 f(u)}{f(u)} - \frac{l(l+1)}{u^2} + \frac{2}{u} - \frac{1}{n^2} \right) = 0 \Rightarrow \\ &\xi(u) \left( -\frac{2(l+1)}{nu} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} + \frac{\partial_u^2 f(u)}{f(u)} + \frac{2}{u} \right) = 0 \Rightarrow \\ &\left( -\frac{2(l+1)}{nu} + \frac{2(l+1)\partial_u f(u)}{uf(u)} - \frac{2\partial_u f(u)}{nf(u)} + \frac{\partial_u^2 f(u)}{f(u)} + \frac{2}{u} \right) = 0 \Rightarrow \\ &\frac{1}{f(u)} \partial_u^2 f(u) + \frac{2(l+1)n-2u}{nuf(u)} \partial_u f(u) + \frac{2n-2(l+1)}{nu} = 0 \end{aligned} \quad (9.61)$$

or, multiplying by  $f(u)$ ,

$$\partial_u^2 f(u) + \frac{2((l+1)n-u)}{nu} \partial_u f(u) + \frac{2(n-l-1)}{nu} f(u) = 0 \quad (9.62)$$

### 9.5.7 A third change of variables

This equation becomes simpler if we perform a further change of variables

$$k + \frac{2u}{n}, \quad \Rightarrow u = \frac{nk}{2}, \quad \Rightarrow \partial_u = \partial_k \frac{\partial k}{\partial u} = \frac{2}{n} \partial_u \quad (9.63)$$

so that

$$\begin{aligned} &\frac{4}{n^2} \partial_k^2 f(k) + \frac{2}{n} 2 \frac{(l+1)n - \frac{nk}{2}}{\frac{n^2 k}{2}} \partial_k f(k) + 2 \frac{n-l-1}{\frac{n^2 k}{2}} f(k) = 0 \\ &\Rightarrow \frac{4}{n^2} \left( \partial_k^2 f(k) + \frac{2(l+1)-k}{k} \partial_k f(k) + \frac{n-l-1}{k} f(k) \right) = 0 \end{aligned} \quad (9.64)$$

Multiplying by  $4k/n^2$  we get

$$k \partial_k^2 f(k) + (2(l+1) - k) \partial_k f(k) + (n-l-1) f(k) = 0 \quad (9.65)$$

### 9.5.8 Power series

Let us define

$$\alpha = 2(l + 1), \quad \beta = n - l - 1 \quad (9.66)$$

so that

$$k\partial_k^2 f(k) + (\alpha - k)\partial_k f(k) + \beta f(k) = 0 \quad (9.67)$$

We look for a power series as a solution

$$f(k) = \sum_{j=0}^{\infty} c_j k^j \quad (9.68)$$

This means that

$$k\partial_k^2 f(k) = k\partial_k \sum_{j=0}^{\infty} j c_j k^{j-1} = k \sum_{j=0}^{\infty} j(j-1) c_j k^{j-2} = \sum_{j=0}^{\infty} j(j-1) c_j k^{j-1} = \sum_{j=1}^{\infty} j(j-1) c_j k^{j-1} \quad (9.69)$$

the last expression being due to the fact that the  $j = 0$  term is equal to  $0^8$ .

Also

$$(\alpha - k)\partial_k f(k) = \sum_{j=0}^{\infty} \alpha j c_j k^{j-1} - \sum_{j=0}^{\infty} j c_j k^j = \sum_{j=1}^{\infty} \alpha j c_j k^{j-1} - \sum_{j=0}^{\infty} j c_j k^j \quad (9.70)$$

where again we got rid of one term that is anyway equal to zero *and that we do not need*. Finally, since

$$\beta f(k) = \sum_{j=0}^{\infty} \beta c_j k^j \quad (9.71)$$

we get

$$\begin{aligned} \sum_{j=1}^{\infty} (c_j(j(j-1))k^{j-1} + \alpha j c_j k^{j-1}) + \sum_{j=0}^{\infty} (\beta - j) c_j k^j &= 0 \\ \Rightarrow \sum_{j=1}^{\infty} j(c_j(j-1) + \alpha) k^{j-1} + \sum_{j=0}^{\infty} (\beta - j) c_j k^j &= 0 \end{aligned} \quad (9.72)$$

Let us use in the first sum

$$j' = j - 1 \quad \Rightarrow \quad j = j' + 1 \quad (9.73)$$

---

<sup>8</sup>Also the  $j = 1$  is equal to zero, but we perform only the manipulations that we are going to need.

## 9.5 The radial equation

---

so that

$$\sum_{j'=0}^{\infty} c_{j'+1}(j'+1)(j'+\alpha)k^{j'} + \sum_{j=0}^{\infty} (\beta-j)c_j k^j = 0 \quad (9.74)$$

and, re-naming  $j' = j$  again in the first sum

$$\begin{aligned} \sum_{j=0}^{\infty} c_{j+1}(j+1)(j+\alpha)k^j + \sum_{j=0}^{\infty} (\beta-j)c_j k^j &= 0 \\ \Rightarrow \sum_{j=0}^{\infty} k^j (c_{j+1}((j+1)(j+\alpha)) - c_j(j-\beta)) &= 0 \end{aligned} \quad (9.75)$$

This will be true for any  $k$  provided that beinequation

$$c_{j+1}((j+1)(j+\alpha)) - c_j(j-\beta) = 0 \quad (9.76)$$

or

$$c_{j+1} = c_j \frac{j-\beta}{(j+1)(j+\alpha)} \quad \forall j \quad (9.77)$$

### 9.5.9 Conditions imposed by the power series solution

If we perform the  $j \rightarrow \infty$  limit in the previous equation we get

$$c_{j+1} \approx \frac{c_j}{j} \quad (9.78)$$

but this asymptotic is the behaviour is like the one of the exponential series

$$\exp(k) = \sum_{j=0}^{\infty} c_j k^j = \sum_{j=0}^{\infty} \frac{1}{j!} k^j \Rightarrow c_j = \frac{1}{j!} \Rightarrow \frac{c_{j+1}}{c_j} = \frac{j!}{(j+1)!} = \frac{1}{j+1} \quad (9.79)$$

But this series diverges for  $k \rightarrow \infty$  (i.e. for large  $u$  and large  $r$ ). We want to avoid such a diverging solution<sup>9</sup>, and thus we need some

$$c_{\tilde{j}+1} = 0 \quad (9.80)$$

for a given integer  $\tilde{j}$ . But this means that the numerator in eq. (9.77) is zero

$$\tilde{j} = \beta = n - l - 1 \quad \Rightarrow n = l + 1 + \tilde{j} \quad (9.81)$$

Since  $\tilde{j} \in (0, \mathbb{N})$ ,  $l \in (0, \mathbb{N})$  we have

$$n \in \mathbb{N}, \quad n > l \quad (9.82)$$

---

<sup>9</sup>Wave solutions have to be normalised to be possible to identify as probability distribution functions; furthermore since we are looking for a bound state of the electron, we don't want the electron to be extremely (divergingly!) probable to be found far from the proton.

The function  $f$  (called a *confluent hypergeometric function* assumes form

$$f^j(k) \propto 1 + \frac{\beta}{\alpha}k + \frac{\beta(\beta+1)}{\alpha(\alpha+1)2!}k^2 + \frac{\beta(\beta+1)(\beta+2)}{\alpha(\alpha+1)(\alpha+2)3!}k^3 + \dots \quad (9.83)$$

with the sum stopping at

$$j = n - l - 1 \quad (9.84)$$

## 9.6 The solution

Summarising what we have found, physical solutions corresponding to bounded states of given energy for the Hydrogen atom may then be identified through the 3 numbers  $n$ ,  $l$  and  $m$

$$|n, m, l\rangle \quad (9.85)$$

All these numbers are  $\in \mathbb{Z}$ .  $n$  is a positive integer while  $l$  is a non-negative integer. We have  $l < n$ , so that for a given value of  $n$  we have  $l = 0, \dots, n-1$ , while for each value of  $l$  the integer  $m$  may assume values  $-l, -l+1, \dots, l-1, l$ . We know, as shown in detail in the appendices, that  $\hbar^2 l(l+1)$  is the value assumed by the magnitude of the angular momentum in the physical state, while  $\hbar m$  is the value assumed by  $L_z$ . The energy of the physical state is (eq. 9.44)

$$E = -\frac{E_0}{n^2} \quad (9.86)$$

depending on the only  $n$  (the energy is independent of the angular momentum). The minimum of this energy is assumed for  $n = 1$ .

The energy, or better the Hamiltonian operator  $\hat{H}$ , the total angular momentum  $\hat{L}^2$  and the  $z$  component  $\hat{L}_z$  are a set of commuting operators<sup>10</sup> that completely specify the state of the electron<sup>11</sup>.

Remembering

$$k = \frac{2u}{n} = \frac{2r}{nr_0} \quad (9.87)$$

we get that to the state  $|n, l, m\rangle$  corresponds the function (recall all the change of variables that we performed!)

$$\psi_{n,l,m}(r, \theta, \varphi) \equiv \frac{1}{r} \left(\frac{r}{r_0}\right)^{l+1} e^{-\frac{r}{nr_0}} f^{n-l-1} \left(\frac{2r}{nr_0}\right) Y_{l,m}(\theta, \varphi) \quad (9.88)$$

---

<sup>10</sup> $\hat{H}$  commutes with the angular momentum and its components since its expression is given by  $\hat{L}^2$  itself plus  $r$  dependent terms that are commuting with  $\hat{L}^2$ , since the latter depends only on angles.

<sup>11</sup>Actually the electron has also an internal angular momentum, its spin, but the description of spin goes beyond the purpose of this course.

## 9.6 The solution

---

The lowest energy state has  $n = 1$ , so necessarily  $l = m = 0$ , and since

$$f^0(k) \propto 1 \quad (9.89)$$

we have

$$\psi_{1,0,0}(r, \theta, \varphi) \propto e^{-\frac{r}{r_0}} \quad (9.90)$$

or, using

$$\int_0^\infty e^{-\frac{r}{r_0}} = \left[ -r_0 e^{-\frac{r}{r_0}} \right]_0^\infty = r_0 \quad (9.91)$$

we get

$$\psi_{1,0,0}(r, \theta, \varphi) = \frac{1}{r_0} e^{-\frac{r}{r_0}} \quad (9.92)$$

The general time dependent solution is

$$\Psi(t, r, \theta, \varphi) = \sum_{n=1}^{\infty} \sum_{l=0}^{n-1} \sum_{m=-l}^l e^{-i\frac{E_0}{\hbar n^2}t} \psi_{n,l,m}(r, \theta, \varphi) \quad (9.93)$$

# Appendix

## 9.A A very shorth introduction to analytical mechanics

There is a very useful function in Classical Mechanics, the Lagrangian function <sup>12</sup> that allows us to easily write down the energy and the equations of motions in different coordinate systems.

### 9.A.1 Cartesian coordinates

#### Single particle

In order for the Lagrangian formulation of mechanics to be valid, it has to be equal to the Newtonian one in Cartesian coordinates. We write, for a single particle moving under a potential  $U$

$$L(\mathbf{x}, \mathbf{v}) = \frac{1}{2}mv^2 - U(\mathbf{x}) \quad (9.94)$$

where we have obviously

$$v^2 = v_x^2 + v_y^2 + v_z^2 = \left(\frac{d}{dt}x\right)^2 + \left(\frac{d}{dt}y\right)^2 + \left(\frac{d}{dt}z\right)^2 \quad (9.95)$$

The first term

$$T = \frac{1}{2}mv^2 \quad (9.96)$$

---

<sup>12</sup>The great French mathematician Joseph-Louis Lagrange, was actually Italian, born in Turin (Kingdom of Sardinia) in 1736 by the name Giuseppe Lodovico Lagrangia. His mathematical treatment of Newtonian Mechanics is, along with the work by Irish William Rowan Hamilton (Dublin, 1805-1865) that extended and completed it, the basis of modern theoretical physics. Lagrange moved to France only when he was 51 (before that he had spent 20 years in Berlin). Lagrange, whose great-grand father was French, became a Senator of the French Republic and was buried, at his death in Paris in 1813, in the Panthéon along with other great contributors to glory of France.

## 9.A A very shorth introduction to analytical mechanics

---

is called *kinetic energy*, so that

$$L = T - U \quad (9.97)$$

is a shorth hand for the Lagrangian.

Let us now define

$$p_i = \frac{\partial L}{\partial v_i} \quad (9.98)$$

e.g.,

$$p_x = \frac{\partial L}{\partial v_x} = mv_x \quad (9.99)$$

These  $p$  are clearly equal to the momenta of newtonian Mechanics. The *Lagrange equations* are now

$$\frac{d}{dt}p_i = \frac{\partial L}{\partial x_i} \quad (9.100)$$

that, in Cartesian coordinantes are equal to the Newton equations of motion

$$m \frac{d^2}{dt^2}x_i = -\frac{\partial U}{\partial x_i} \quad (9.101)$$

### Many particles

We write the Lagrangian as

$$L(\mathbf{x}_i, \mathbf{v}_i) = \sum_i \frac{1}{2}m_i v_i^2 - U(\mathbf{x}_i) \quad (9.102)$$

where

$$v_i^2 = v_{i,x}^2 + v_{i,y}^2 + v_{i,z}^2 \quad (9.103)$$

We define again

$$p_{i,j} = \frac{\partial L}{\partial v_{i,j}} \quad (9.104)$$

and the Lagrange equations

$$\frac{d}{dt}p_{i,j} = \frac{\partial L}{\partial x_{i,j}} \quad (9.105)$$

are again equivalent to the Newtonian ones



## Energy

The energy (or Hamiltonian function) is defined by

$$H(p_{i,j}, x_{i,j}) = \sum_{i,j} p_{i,j} v_{i,j} - L(x_{i,j}, v_{i,j}(p_{i,j})) \quad (9.106)$$

Let us compute this for a single particle. We need to obtain the  $v$ s as functions of the  $p$ s. From

$$v_i = \frac{p_i}{m} \quad (9.107)$$

we get

$$H = \sum_{i=x,y,z} \frac{p_i^2}{m} - \sum_{i=x,y,z} \frac{p_i^2}{2m} + U(\mathbf{x}) = \sum_i \frac{p_i^2}{2m} + U(\mathbf{x}) = T + U \quad (9.108)$$

which is clearly the energy of the Newtonian system. While  $H$  is a function of momenta and positions, it may be shown that for a large class of systems it is equal to a constant  $E$  (the energy).

Indeed, if  $L$ , as we assumed above, is a function of the  $x$  and  $v$ , but not an explicit function of time, we have

$$\frac{d}{dt}H = \sum_{i,j} \dot{p}_{i,j} v_{i,j} + \sum_{i,j} p_{i,j} \dot{v}_{i,j} - \sum_{i,j} \frac{\partial L}{\partial x_{i,j}} \dot{x}_{i,j} - \sum_{i,j} \frac{\partial L}{\partial v_{i,j}} \dot{v}_{i,j} \quad (9.109)$$

We have use Newton's notation for time derivatives

$$\frac{df}{dt} \equiv \dot{f} \quad (9.110)$$

Now, since

$$x_{i,j} \dot{=} v_{i,j}, \quad \frac{\partial L}{\partial v_{i,j}} = p_{i,j}, \quad \frac{\partial L}{\partial x_{i,j}} = p_{i,j} \quad (9.111)$$

we get

$$\frac{d}{dt}H = \sum_{i,j} \dot{p}_{i,j} v_{i,j} + \sum_{i,j} p_{i,j} \dot{v}_{i,j} - \sum_{i,j} \dot{p}_{i,j} v_{i,j} - \sum_{i,j} p_{i,j} \dot{v}_{i,j} = 0 \quad (9.112)$$

### 9.A.2 Arbitrary coordinates

The Lagrangian formulation becomes extremely useful when we are using non-Cartesian coordinates. Let us assume that we may write the Cartesian coordinates  $x_{i,j}$  as functions of non-Cartesian coordinates  $q_k$

$$x_{i,j} = x_{i,j}(q_k) \quad (9.113)$$

## 9.A A very shorth introduction to analytical mechanics

---

For example, in case of a single particle, we may write  $x$ ,  $y$  and  $z$  as functions of the spherical coordinates

$$x = x(r, \theta, \varphi), \quad y = y(r, \theta, \varphi), \quad z = z(r, \theta, \varphi) \quad (9.114)$$

so that we have, for example  $q_1 = r$ ,  $q_2 = \theta$  and  $q_3 = \varphi$ . In this case, we have exactly 3 non-Cartesian coordinates expressing 3 Cartesian ones, but the situation may be different, and we may have less  $qs$  than  $xs$ . For example, if a particle is constrained to move on a sphere of radius  $r = R$ , we may use the only  $\theta$  and  $\varphi$  coordinates,

$$x = x(\theta, \varphi), \quad y = y(\theta, \varphi), \quad z = z(\theta, \varphi) \quad (9.115)$$

It may be shown that everything we need to do to compute the equations of motion and the energy in the new coordinate systems is to express the Lagrangian in them. Regarding the potential energy  $U$  we just need to re-express it as a function of the  $q$  using the relations 9.113 or the kinetic energy we write

$$T = \frac{1}{2} \sum_{i,j} m_i \dot{x}_{i,j}^2 = \frac{1}{2} \sum_{i,j} m_i (\dot{x}_{i,j})(\dot{x}_{i,j}) \quad (9.116)$$

Now we use

$$\dot{x}_{i,j} = \sum_k \frac{\partial x_{i,j}}{\partial q_k} \dot{q}_k \quad (9.117)$$

and obtain

$$T = \frac{1}{2} \sum_{i,j} m_i \sum_{k,l} \frac{\partial x_{i,j}}{\partial q_k} \frac{\partial x_{i,j}}{\partial q_l} \dot{q}_k \dot{q}_l \equiv \sum_{k,l} T_{k,l} \dot{q}_k \dot{q}_l \quad (9.118)$$

where we defined

$$\frac{1}{2} \sum_{i,j} m_i \frac{\partial x_{i,j}}{\partial q_k} \frac{\partial x_{i,j}}{\partial q_l} \equiv T_{k,l} \quad (9.119)$$

After we have obtained  $L(q_i, \dot{q}_i)$  we use the same technique that we applied in the Cartesian case,

$$p_i = \frac{\partial L}{\partial \dot{q}_i} \quad (9.120)$$

$$\dot{p}_i = \frac{\partial L}{\partial q_i} \quad (9.121)$$

$$H(q_i, p_i) = \sum_i p_i \dot{q}_i - L(q_i, \dot{q}_i(p_i)) \quad (9.122)$$

## 9.B Two body problem

### 9.B.1 Introduction

As a first application of the Lagrangian method we may show how to deal with a system composed of two particles (electron and proton) or two “bodies” that may be approximated as particles by ignoring their internal structure (Sun and Earth, or the proton itself), by reducing it to a problem dealing with a single particle (actually most of the following treatment may be understood just using Newtonian dynamics).

### 9.B.2 Newtonian treatment

#### Definitions

Let us call the proton “body 1”, and identify it by its mass and position and momentum in an arbitrary Cartesian frame

$$m_1, \quad \mathbf{r}_1, \quad \mathbf{p}_1 = m_1 \dot{\mathbf{r}}_1 \quad (9.123)$$

and the electron “body 2”

$$m_2, \quad \mathbf{r}_2, \quad \mathbf{p}_2 = m_2 \dot{\mathbf{r}}_2 \quad (9.124)$$

Although for the proton-electron problem (as for the Sun-Earth one) we have  $m_1 \gg m_2$  the following analysis does not depend on the  $m_1/m_2$  ratio.

Let us first define the relative distance as

$$\mathbf{r} \equiv \mathbf{r}_2 - \mathbf{r}_1 \quad (9.125)$$

and the relative velocity as

$$\mathbf{v} \equiv \dot{\mathbf{r}} = \mathbf{v}_2 - \mathbf{v}_1 = \frac{\mathbf{p}_2}{m_2} - \frac{\mathbf{p}_1}{m_1} \quad (9.126)$$

The centre of mass of the system is defined as usual by

$$\mathbf{R} \equiv \frac{m_2 \mathbf{r}_2 + m_1 \mathbf{r}_1}{m_1 + m_2} \quad (9.127)$$

The denominator is the total mass

$$M \equiv m_1 + m_2 \quad (9.128)$$

so that

$$\mathbf{R} \equiv \frac{m_2 \mathbf{r}_2 + m_1 \mathbf{r}_1}{M} \quad (9.129)$$

## 9.B Two body problem

---

The velocity of the centre of mass is

$$\mathbf{V} \equiv \dot{\mathbf{R}} \equiv \frac{m_2 \mathbf{v}_2 + m_1 \mathbf{v}_1}{M} \quad (9.130)$$

If we multiply this velocity by  $M$  we get the total momentum (that we may call  $\mathbf{P}$ )

$$M\mathbf{V} = m_2 \mathbf{v}_2 + m_1 \mathbf{v}_1 = \mathbf{p}_2 + \mathbf{p}_1 = \mathbf{P} \quad (9.131)$$

All this definitions were quite intuitive. We now define the *reduced mass*

$$m = \frac{m_1 m_2}{M} \quad (9.132)$$

and the *relative momentum*, the momentum of a particle with reduced mass and relative velocity

$$\mathbf{p} = m\mathbf{v} = m \left( \frac{\mathbf{p}_2}{m_2} - \frac{\mathbf{p}_1}{m_1} \right) \quad (9.133)$$

### Energy

The energy of the system is

$$E = \frac{p_1^2}{2m_1} + \frac{p_2^2}{2m_2} + U(|r_2 - r_1|) \quad (9.134)$$

Here we have done an important assumption about the interaction potential, namely that it depends only on the distance between the two particles, as it is true for gravitational (Sun-Earth) and electrical (electron-proton) forces<sup>13</sup>.

We may show that, when expressed using centre of mass variables, the energy becomes

$$E = \frac{P^2}{2M} + \frac{p^2}{2m} + U(r) \quad (9.135)$$

---

<sup>13</sup>It may be easily shown that a potential in the form  $U(\mathbf{r}_2 - \mathbf{r}_1)$  implies conservation of momentum, or the “weak” form of Newton’s third law, i.e. that the forces between two particles have to be equal and opposite, while a potential in the form  $U(|r_2 - r_1|)$  implies the “strong” form of Newton’s third law, namely that the forces have to be directed along the direction  $\mathbf{r}_2 - \mathbf{r}_1$ . It is thus natural to expect that the fundamental forces of nature, at least in their classical form, may be expressed through a  $U(|r_2 - r_1|)$  potential (we are not considering here the possibility of velocity dependent forces, such as magnetic forces).

It is obvious that  $U(|r_2 - r_1|) = U(r)$ . For the kinetic energy we have

$$\begin{aligned}
 2T &= \frac{P^2}{M} + \frac{p}{m} = \frac{p_2^2 + p_1^2 + 2\mathbf{p}_1 \cdot \mathbf{p}_2}{M} + \frac{1}{m} \frac{m^2 (p_2^2 m_1^2 + p_1^2 m_2^2 - 2m_1 m_2 \mathbf{p}_1 \cdot \mathbf{p}_2)}{m_1^2 m_2^2} \\
 &= \frac{p_2^2 + p_1^2 + 2\mathbf{p}_1 \cdot \mathbf{p}_2}{m_1 + m_2} + \frac{m_1 m_2}{m_1 + m_2} \frac{p_2^2 m_1^2 + p_1^2 m_2^2 - 2m_1 m_2 \mathbf{p}_1 \cdot \mathbf{p}_2}{(m_1 m_2)^2} \\
 &= \frac{p_2^2 + p_1^2 + 2\mathbf{p}_1 \cdot \mathbf{p}_2}{m_1 + m_2} + \frac{1}{m_1 + m_2} \frac{p_2^2 m_1^2 + p_1^2 m_2^2 - 2m_1 m_2 \mathbf{p}_1 \cdot \mathbf{p}_2}{m_1 m_2} \\
 &= \frac{p_2^2 + p_1^2 + 2\mathbf{p}_1 \cdot \mathbf{p}_2}{m_1 + m_2} + \frac{1}{m_1 + m_2} \left( \frac{p_2^2 m_1}{m_2} + \frac{p_1^2 m_2}{m_1} - 2\mathbf{p}_1 \cdot \mathbf{p}_2 \right) \\
 &= p_2^2 \left[ \frac{1}{m_1 + m_2} \left( 1 + \frac{m_1}{m_2} \right) \right] + p_1^2 \left[ \frac{1}{m_1 + m_2} \left( 1 + \frac{m_1}{m_2} \right) \right] \\
 &= p_2^2 \left[ \frac{1}{m_1 + m_2} \frac{m_2 + m_1}{m_2} \right] + p_1^2 \left[ \frac{1}{m_1 + m_2} \frac{m_1 + m_2}{m_2} \right] = \frac{p_1^2}{m_1} + \frac{p_2^2}{m_2}
 \end{aligned} \tag{9.136}$$

From the expression of the energy, we see that it splits into the energy of the centre of mass

$$E^{CM} = \frac{P^2}{2M} \tag{9.137}$$

i.e. of a particle with mass  $M$  subject to no potential, and thus moving with constant velocity, and the energy of relative motion

$$E^{Rel} = \frac{p}{2m} + U(r) \tag{9.138}$$

We may thus study the two body problem just like a single body one. The distance  $r$  is the distance between the two particles, but the position of the proton is not the centre of an inertial frame. An actual inertial frame is given by the position of the centre of mass  $\mathbf{R}$ . Anyway, if  $m_1 \gg m_2$ , we have

$$\mathbf{R} = \frac{m_1 \mathbf{r}_1 + m_2 \mathbf{r}_2}{m_1 + m_2} = \frac{m_1 \mathbf{r}_1}{m_1 \left( 1 + \frac{m_2}{m_1} \right)} + \frac{m_2 \mathbf{r}_2}{m_1 \left( 1 + \frac{m_2}{m_1} \right)} \approx \mathbf{r}_1 \tag{9.139}$$

Again, although the mass entering in the relative energy is not the electron mass but the reduced mass, if  $m_1 \gg m_2$  we have

$$m = \frac{m_1 m_2}{m_1 + m_2} = \frac{m_1 m_2}{m_1 \left( 1 + \frac{m_2}{m_1} \right)} \approx m_2 \tag{9.140}$$

### 9.B.3 Lagrangian treatment

In the Newtonian treatment, the reduced mass and relative momentum were introduced as a guess. The Lagrangian treatment just starts from the defi-

## 9.B Two body problem

---

inition of the centre of mass and relative distance (the  $q$  variables)

$$\mathbf{r} \equiv \mathbf{r}_2 - \mathbf{r}_1, \quad \mathbf{R} \equiv \frac{m_2 \mathbf{r}_2 + m_1 \mathbf{r}_1}{m_1 + m_2} \quad (9.141)$$

and of the corresponding derivatives ( $\dot{q}$  variables)

$$\mathbf{v} \equiv \dot{\mathbf{r}} = \mathbf{v}_2 - \mathbf{v}_1, \quad \mathbf{V} \equiv \dot{\mathbf{R}} \equiv \frac{m_2 \mathbf{v}_2 + m_1 \mathbf{v}_1}{M} \quad (9.142)$$

To write the Lagrangian in the new variables we need an expression of the  $x$  variables as functions of the  $q$  ones. We use

$$\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1 \Rightarrow \mathbf{r}_2 = \mathbf{r} + \mathbf{r}_1 \quad (9.143)$$

so that

$$\mathbf{R} = \frac{m_1 \mathbf{r}_1 + m_2 (\mathbf{r} + \mathbf{r}_1)}{m_1 + m_2} = \mathbf{r}_1 + \mathbf{r} \frac{m_2}{m_1 + m_2} \Rightarrow \mathbf{r}_1 = \mathbf{R} - \mathbf{r} \frac{m_2}{m_1 + m_2} \quad (9.144)$$

Similarly

$$\mathbf{r} = \mathbf{r}_2 - \mathbf{r}_1 \Rightarrow \mathbf{r}_1 = \mathbf{r}_2 - \mathbf{r} \quad (9.145)$$

so that

$$\mathbf{R} = \frac{m_2 \mathbf{r}_2 + m_1 (\mathbf{r}_2 - \mathbf{r})}{m_1 + m_2} = \mathbf{r}_2 - \mathbf{r} \frac{m_1}{m_1 + m_2} \Rightarrow \mathbf{r}_2 = \mathbf{R} + \mathbf{r} \frac{m_1}{m_1 + m_2} \quad (9.146)$$

These latter relations show very clearly that if  $m_1 \gg m_2$  we have

$$\mathbf{r}_2 \approx \mathbf{R} + \mathbf{r} \quad \mathbf{r}_1 \approx \mathbf{R} \quad (9.147)$$

Trivially, by derivation, we have

$$\mathbf{v}_1 = \mathbf{V} - \mathbf{v} \frac{m_2}{m_1 + m_2}, \quad \mathbf{v}_2 = \mathbf{V} + \mathbf{v} \frac{m_1}{m_1 + m_2} \quad (9.148)$$

We have now

$$m_1 v_1^2 = m_1 V^2 + \frac{m_1 m_2^2}{(m_1 + m_2)^2} v^2 - 2\mathbf{V} \cdot \mathbf{v} \frac{m_2 m_1}{m_1 + m_2} \quad (9.149)$$

and

$$m_2 v_2^2 = m_2 V^2 + \frac{m_2 m_1^2}{(m_1 + m_2)^2} v^2 + 2\mathbf{V} \cdot \mathbf{v} \frac{m_2 m_1}{m_1 + m_2} \quad (9.150)$$

so that

$$\begin{aligned} m_1 v_1^2 + m_2 v_2^2 &= (m_1 + m_2) V^2 + \frac{m_1 m_2^2 + m_2 m_1^2}{(m_1 + m_2)^2} v^2 = M V^2 + \frac{(m_1 m_2)(m_1 + m_2)}{(m_1 + m_2)^2} v^2 \\ &= M V^2 + \frac{m_1 m_2}{m_1 + m_2} v^2 = M V^2 + m v^2 \end{aligned} \quad (9.151)$$

We see that now  $M$  and  $m$  are naturally defined by the Lagrangian formalism. The Lagrangian function is

$$L = \frac{1}{2} (m_1 v_1^2 + m_2 v_2^2) - U(|r_2 - r_1|) = \frac{1}{2} (MV^2 + mv^2) - U(r) \quad (9.152)$$

Again from the Lagrange formalism

$$p_i = \frac{\partial L}{\partial \dot{q}_i} \quad (9.153)$$

we get

$$\mathbf{P} = M\mathbf{V}, \quad \mathbf{p} = m\mathbf{v} \quad (9.154)$$

The energy is

$$E = \mathbf{P} \cdot \mathbf{V} + \mathbf{p} \cdot \mathbf{v} - L \quad (9.155)$$

using

$$\mathbf{V} = \frac{\mathbf{P}}{M}, \quad \mathbf{v} = \frac{\mathbf{p}}{m} \quad (9.156)$$

and substituting, we get

$$E = \frac{P^2}{2M} + \frac{p^2}{2m} + U(r) \quad (9.157)$$

The equation of motion are

$$\dot{P}_i = \frac{\partial L}{\partial R_i} = 0 \quad (9.158)$$

so that the centre of mass moves with constant momentum (velocity), and

$$\dot{p}_i = \frac{\partial L}{\partial R_i} = -\frac{\partial U}{\partial r_i} \quad (9.159)$$

justifying the treatment of the relative motion as a single particle under the effect of the potential  $U$ .

## 9.C Angular momentum in Classical and Quantum Mechanics

### 9.C.1 Definitions

The angular momentum is defined as the vector product

$$\mathbf{L} = \mathbf{x} \wedge \mathbf{p} \quad (9.160)$$

## 9.C Angular momentum in Classical and Quantum Mechanics

---

This expression may be written also using the Levi-Civita symbol  $\varepsilon$

$$L_i = \sum_{j,k} \varepsilon_{i,j,k} x_j p_k \quad (9.161)$$

Here the indexes are assuming values between 1 and 3, and the Levi-Civita symbol is defined as having

$$\varepsilon_{1,2,3} = 1 \quad (9.162)$$

and being completely asymmetric, i.e. it changes sign if we switch any index. As a result terms with repeated indexes are zero, e.g.

$$\varepsilon_{1,2,2} = -\varepsilon_{1,2,2} \Rightarrow \varepsilon_{1,2,2} = 0 \quad (9.163)$$

terms in which the indexes are “in order” are 1, e.g.

$$\varepsilon_{3,1,2} = -\varepsilon_{2,1,3} = -\varepsilon_{1,2,3} = 1 \quad (9.164)$$

while terms in which the indexes are “not in order” are -1, e.g.

$$\varepsilon_{3,2,1} = -\varepsilon_{3,1,2} = -1 \quad (9.165)$$

So we have, for example,

$$L_z = L_3 = \sum_{j,k} \varepsilon_{3,j,k} x_j p_k \quad (9.166)$$

$j$  and  $k$  may not assume value 3 (since the corresponding  $\varepsilon$  would be zero) so that

$$L_z = L_3 = \varepsilon_{3,1,2} x_1 p_2 + \varepsilon_{3,2,1} x_2 p_1 = x p_y - y p_x \quad (9.167)$$

### 9.C.2 Lagrangian for a central potential

We have seen that when dealing with a two body problem in which the potential is in the form  $U(|r_2 - r_1|)$ , the Lagrangian for the relative motion may be written as

$$L = \frac{1}{2} m v^2 - U(r) \quad (9.168)$$

Since the potential depends only on  $r$ , we may want to pass to spherical coordinates. The lagrangian formalism asks us to re-write

$$v^2 = \dot{x}^2 + \dot{y}^2 + \dot{z}^2 \quad (9.169)$$

as a function of  $\dot{r}$ ,  $\dot{\theta}$  and  $\dot{\varphi}$ . We may obviously use

$$\dot{x} = \partial_r \dot{r} + \partial_\theta \dot{\theta} + \partial_\varphi \dot{\varphi} \quad (9.170)$$



etc., but an easier way is to remember, from chapter 1, that

$$v = \frac{d\mathbf{s}}{dt} \Rightarrow v^2 = \left(\frac{ds}{dt}\right)^2, \quad ds^2 = dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\varphi^2 \Rightarrow v^2 = \dot{r}^2 + r^2 \dot{\theta}^2 + r^2 \sin^2 \theta \dot{\varphi}^2 \quad (9.171)$$

and thus

$$L = \frac{1}{2}m \left( \dot{r}^2 + r^2 \dot{\theta}^2 + r^2 \sin^2 \theta \dot{\varphi}^2 \right) - U(r) \quad (9.172)$$

We compute the momenta

$$p_r = \frac{\partial L}{\partial \dot{r}} = m\dot{r} \quad (9.173)$$

$$p_\theta = \frac{\partial L}{\partial \dot{\theta}} = mr^2 \dot{\theta} \quad (9.174)$$

$$p_\varphi = \frac{\partial L}{\partial \dot{\varphi}} = mr^2 \sin^2 \theta \dot{\varphi} \quad (9.175)$$

Since  $U = U(r)$  we have

$$\dot{p}_\theta = \frac{\partial L}{\partial \theta} = 0 \Rightarrow p_\theta = \text{const} \quad (9.176)$$

$$\dot{p}_\varphi = \frac{\partial L}{\partial \varphi} = 0 \Rightarrow p_\varphi = \text{const} \quad (9.177)$$

At time  $t = 0$ ,  $\mathbf{r}$  and  $\mathbf{v}$  assume given values, and their linear combinations form a plane. Let us call this plane the  $x, y$  plane. On this plane (eq. 1.27) we have  $\sin \theta = 1$ , and, since the velocity is directed along the  $\theta = \pi/2$  plane,  $\dot{\theta} = 0$ . As a result the initial value of the  $\theta$  momentum is

$$p_\theta = mr^2 \dot{\theta} = 0 \quad (9.178)$$

But since this is a constant, we have

$$\dot{\theta} = 0 \quad \forall t, \Rightarrow \theta = \frac{\pi}{2}, \sin \theta = 1 \quad \forall t \quad (9.179)$$

The Lagrangian becomes

$$L = \frac{1}{2}m \left( \dot{r}^2 + r^2 \dot{\varphi}^2 \right) - U(r) \quad (9.180)$$

and the  $\varphi$  momentum is

$$p_\varphi = mr^2 \dot{\varphi} \quad (9.181)$$

This is related to the conservation of angular momentum. Indeed, if the motion is limited to the  $z = 0$  plane,

$$L_x = \varepsilon_{1,2,3} x_2 p_3 + \varepsilon_{1,3,2} x_3 p_2 = y p_z - z p_y = 0; \quad L_y = \varepsilon_{2,1,3} x_1 p_3 + \varepsilon_{2,3,1} x_3 p_1 = -x p_z + z p_x = 0 \quad (9.182)$$

## 9.C Angular momentum in Classical and Quantum Mechanics

---

and thus

$$\mathbf{L} = L_z \mathbf{e}_z \quad (9.183)$$

This  $L_z$  is exactly  $p_\varphi$ . Indeed,

$$L_z = |\mathbf{L}| = \mathbf{r} \wedge \mathbf{p} = mrv \sin \alpha \quad (9.184)$$

where  $\alpha$  is the angle between  $\mathbf{r}$  and  $\mathbf{v}$ . Now  $v \sin \alpha$  is the projection of  $\mathbf{v}$  in the direction orthogonal to  $\mathbf{r}$ , which is given, at first order, by  $r\dot{\varphi}$  (Fig. 9.C.1) so that




Figure 9.C.1: Geometrical interpretation of  $p_\varphi$ .

$$L_z = mr^2\dot{\varphi} \quad (9.185)$$

Another way to see it is to use an (instantaneous) frame with  $x$  and  $r$  aligned, so that

$$\varphi = 0 \Rightarrow x = r, \quad y = 0, \quad v_x = \dot{r} \cos \varphi + r \sin \varphi \dot{\varphi} = \dot{r}, \quad v_y = \dot{r} \sin \varphi + r \cos \varphi \dot{\varphi} = r\dot{\varphi} \quad (9.186)$$

and finally

$$L_z = m(xv_y - yv_x) = mr^2\dot{\varphi} \quad (9.187)$$

Let us thus have

$$p_\varphi = L_z = mr^2\dot{\varphi} \Rightarrow \dot{\varphi} = \frac{L_z}{mr^2} \quad (9.188)$$

and

$$L = \frac{1}{2} \left( mr\dot{r}^2 + \frac{L_z^2}{mr} \right) - U(r) \quad (9.189)$$

Now, using

$$\dot{r} = \frac{p_r}{m} \quad (9.190)$$

we have

$$H = p_r \dot{r} + p_\varphi \dot{\varphi} - \frac{1}{2} \left( m\dot{r}^2 + \frac{L_z^2}{mr^2} \right) + U(r) = \frac{p_r^2}{m} + \frac{p_\varphi^2}{mr^2} - \frac{p_r^2}{2m} - \frac{p_\varphi^2}{2mr^2} + U(r)$$

$$\frac{p_r^2}{2m} + \frac{p_\varphi^2}{2mr^2} + U(r)$$

(9.191)

or, remembering that  $p_\varphi^2$  gives the magnitude of the angular momentum  $\mathbf{L}$ ,

$$H = \frac{p_r^2}{2m} + \frac{L^2}{2mr^2} + U(r)$$

(9.192)

This is equivalent to the motion of a particle in a one dimensional potential *effective potential*

$$\frac{L^2}{2mr^2} + U(r)$$

(9.193)

### 9.C.3 Angular momentum in Quantum Mechanics

#### A guess for the expression of $L^2$

For the solution of the classical problem, you may check any good book in Classical Mechanics. We are now interested in comparing

$$H = \frac{p_r^2}{2m} + \frac{L^2}{2mr^2} + U(r)$$

(9.194)

with its Quantum equivalent, the time Schrödinger equation (refer to 9.15)

$$H\Psi(t, r, \theta, \varphi) = \left( \frac{\hbar^2}{2m} \left[ \frac{1}{r^2} \partial_r (r^2 \partial_r) + \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \right] + U(r) \right) \Psi_E(t, r, \theta, \varphi)$$

(9.195)

We are left to identify the quantum operators

$$\hat{p}_r^2 = \hbar^2 \frac{1}{r^2} \partial_r (r^2 \partial_r)$$

(9.196)

$$\hat{L}^2 = \hbar^2 \left[ \frac{1}{r^2 \sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{r^2 \sin^2 \theta} \partial_\varphi^2 \right]$$

(9.197)

We will thus assume that this is the form assumed by the squared angular momentum operator in Quantum Mechanics.

The angular momentum is obviously given by

$$\hat{L}_i = \sum_{j,k} \varepsilon_{ijk} \hat{x}_j \hat{y}_k$$

(9.198)

## 9.C Angular momentum in Classical and Quantum Mechanics

---

with the position operator being a multiplicative operator (since when we find a potential that is a function of a position we get a multiplicative operator from it)

$$\hat{x}_j \Psi(t, \mathbf{x}) = x_j \Psi(t, \mathbf{x}) \quad (9.199)$$

and as usual

$$\hat{p}_k \Psi(t, \mathbf{x}) = -i\hbar \partial_k \Psi(t, \mathbf{x}) \quad (9.200)$$

The computation showing that eq. (9.197) actually derives from eq. (9.198) is quite lengthy and left for appendix 9.D.

### Physical meaning of self-adjoint operators in Quantum Mechanics

In Quantum Mechanics, normalised vectors in a space provided with a scalar product represent physical states, and self-adjoint operators represent physical observables. Given a state  $|\Psi\rangle$  and an operator  $A$ ,

$$\langle \Psi | A | \Psi \rangle \equiv \langle A \rangle_\Psi \quad (9.201)$$

gives the *expectation value* of  $A$  in the state  $\Psi$ , i.e. the average value assumed by the observable in that state (average over a large number of experiments). If  $\Psi$  is an eigenvector of  $A$ , e.g., if we consider the Hamiltonian observable  $\hat{H}$  and one of its eigenvectors  $\psi_E$ , we have

$$\langle H \rangle_{\psi_E} = \langle \psi_E | \hat{H} | \psi_E \rangle = E \langle \psi_E | \psi_E \rangle = E \quad (9.202)$$

Eigenvectors are states in which the observable assumes a fixed value, i.e. repeating the experiment we get always the same result and (using again  $\hat{H}$  as an example)

$$(\Delta H)^2 = \langle \psi_E | \hat{H}^2 | \psi_E \rangle - \left( \langle \psi_E | \hat{H} | \psi_E \rangle \right)^2 = E^2 - E^2 = 0 \quad (9.203)$$

In general, since the eigenvectors of a self-adjoint operator form a basis, we have (using again  $\hat{H}$  as an example)

$$|\Psi\rangle = \sum_i c_i |\psi_{E_i}\rangle \quad (9.204)$$

where the  $c_i$  are complex numbers. The postulates of Quantum Mechanics tell us that single experiments will find one of the  $E_i$  values with probability  $|c_i|^2$ , and the variation will be non zero

$$(\Delta H)^2 > 0 \quad (9.205)$$

For the operator  $L^2$  we know (eq. 7.72 and eq. 9.197) that

$$\hat{L}^2 Y_{l,m} = l(l+1)\hbar^2 \quad (9.206)$$

This means that the spherical harmonics are the eigenvectors of  $\hat{L}^2$  (states with a fixed angular momentum magnitude), and that angular momentum may assume only quantised values

$$L^2 = l(l + 1)\hbar^2 \quad (9.207)$$

### 9.C.4 Commutators in Quantum Mechanics

#### Physical meaning

In section 6.3.6 we have defined the commutator

$$[A, B] = AB - BA \quad (9.208)$$

and we have discussed how self-adjoint operators admit a common basis of eigenvectors only if they commute

$$[A, B] = 0 \quad (9.209)$$

It is indeed clear that if  $\Psi$  is an eigenvector of  $A$  and  $B$ , let us say with eigenvalues  $a$  and  $b$ , at least we have

$$AB|\Psi\rangle = bA|\Psi\rangle = ba|\Psi\rangle = bA|\Psi\rangle = BA|\Psi\rangle \quad (9.210)$$

This means that if

$$\langle\Psi|[A, B]|\Psi\rangle \neq 0 \quad (9.211)$$

Then  $\Psi$  may not be an eigenvector of both operators. Physically, this means that it is impossible to have a determined value of both observables in the physical state corresponding to  $\Psi$ .

#### The $[\hat{x}, \hat{p}]$ commutator

The most important commuting relation in Quantum Physics is the one between the position operator  $\hat{x}$ , related to the physical observable of measuring the position of a particle, and the momentum operator  $\hat{p}$ , related to the physical observable of measuring the momentum of a particle.

Let us compute

$$[\hat{x}, \hat{p}_x] \quad (9.212)$$

by having the resulting operator act on an arbitrary function of  $x$ ,

$$\begin{aligned} [\hat{x}, \hat{p}_x]\psi(x) &= -i\hbar(x\partial_x\psi - \partial_x(x\psi(x))) = \\ &= -i\hbar(x\partial_x\psi(x) - x\partial_x\psi(x) - \psi(x)) = i\hbar\psi(x) \end{aligned} \quad (9.213)$$

## 9.C Angular momentum in Classical and Quantum Mechanics

---

so that

$$[\hat{x}, \hat{p}_x] = i\hbar \mathbf{1} \quad (9.214)$$

This tells us that is *impossible to find a physical state in which both position and velocity are exactly defined (eigenvectors), or, equivalently that it is impossible to measure at the same time the position and momentum of a particle*. This is, fundamentally, the content of the celebrated *Uncertainty principle* due to Heisenberg (which is studied in more detail in appendix 9.E).

Obviously we have

$$[\hat{x}, \hat{y}] = xy - yx = xy - xy = 0 \quad (9.215)$$

and

$$[\hat{x}, \hat{p}_y] = -i\hbar(x\partial_y - \partial_y x) = (x\partial_y - x\partial_y) = 0 \quad (9.216)$$

$$[\hat{p}_x, \hat{p}_y] = -i\hbar(\partial_x\partial_y - \partial_y\partial_x) = 0 \quad (9.217)$$

since derivatives commute, so that

$$[\hat{x}_k, \hat{p}_j] = i\hbar\delta_{k,j}, \quad [\hat{x}_k, \hat{x}_j] = [\hat{p}_k, \hat{p}_j] = 0 \quad (9.218)$$

From a physical point of view, this means that different component of position, momentum, or component of position and momentum in different directions may be measured with arbitrary precision at the same (i.e. we may find common eigenstates of these observables). The uncertainty principle only applies to position and momentum in the same direction.

### Are the operators that we are considering self-adjoint?

Using the commuting properties of  $\hat{x}$  and  $\hat{p}$  we may now show that if these two operators are self-adjoint, then also all the components of  $\hat{L}$  are self-adjoint. Indeed,

$$\begin{aligned} \hat{L}_i^\dagger &= \left( \sum_{j,k} \varepsilon_{i,j,k} \hat{x}_j \hat{p}_k \right)^\dagger = \sum_{j,k} \varepsilon_{i,j,k} (\hat{x}_j \hat{p}_k)^\dagger = \\ &= \sum_{j,k} \varepsilon_{i,j,k} \hat{p}_k^\dagger \hat{x}_j^\dagger = \sum_{j,k} \varepsilon_{i,j,k} \hat{p}_k \hat{x}_j = \sum_{j,k} \varepsilon_{i,j,k} \hat{x}_j \hat{p}_k = \hat{L}_i \end{aligned} \quad (9.219)$$

where we have used the fact that for  $\varepsilon_{i,j,k}$  to be non-zero we need  $j \neq k$  and thus  $[\hat{x}_j, \hat{p}_k] = 0$ .

We still have to show that the position and momentum operators are self-adjoint. Establishing if an operator is self-adjoint is often a non-trivial

problem in functional analysis, one of the main issues being to show that the domain of the adjoint operator is the same as the original one. We ignore here the subtle problems and show that

$$\langle \psi, \hat{x}\phi \rangle = \int x\psi(x)^*\phi(x)dx = \langle \hat{x}\psi, \phi \rangle \quad (9.220)$$

and

$$\begin{aligned} \langle \psi, \hat{p}_x\phi \rangle &= \hbar \int \psi(x)^*i\partial_x\phi(x) = \hbar \int (-i\psi(x))^* \partial_x\phi(x) = \\ &= -\hbar \int \partial_x(-i\psi(x)\phi(x)) dx + \hbar \int (i\partial_x\psi(x))^*\phi(x)dx = \\ &= -\hbar [-i\psi(x)\phi(x)]_{\text{boundary}} + \hbar \int (i\partial_x\psi(x))^*\phi(x)dx = \hbar \int (i\partial_x\psi(x))^*\phi(x)dx = \langle \hat{p}_x\psi, \phi \rangle \end{aligned} \quad (9.221)$$

where we have assumed that the boundary contributions are equal to zero.

### Properties of commutators

What are the commuting properties of the angular momentum  $\hat{L}$ ? In order to obtain them, we will need to compute expressions like

$$[\hat{L}_x, \hat{L}_y] = [\hat{y}\hat{p}_z - \hat{z}\hat{p}_y, \hat{z}\hat{p}_x - \hat{x}\hat{p}_z] \quad (9.222)$$

It is thus useful to learn some formulae that may facilitate these computations, as for example the formula for a commutator with a sum

$$[A + B, C] = [A, C] + [B, C] \quad (9.223)$$

since

$$(A + B)C - C(A + B) = AC - CA + BC - CB \quad (9.224)$$

Obviously we have

$$[A, B] = -[B, A] \quad (9.225)$$

so that

$$[C, A + B] = -([A, C] + [B, C]) = [C, A] + [C, B] \quad (9.226)$$

These equations show that if  $B$  commutes with  $C$ , it may be ignored

$$[A + B, C] = [A, C] \quad \text{if } [B, C] = 0 \quad (9.227)$$

We also have that if  $\alpha$  is a number,

$$\alpha[A, B] = [\alpha A, B] = [A, \alpha B] \quad (9.228)$$

## 9.C Angular momentum in Classical and Quantum Mechanics

---

since

$$[\alpha A, B] = \alpha AB - \alpha BA \quad (9.229)$$

Finally we have

$$[AB, C] = A[B, C] + [A, C]B \quad (9.230)$$

since

$$ABC - CAB = ABC - ACB + ACB - CAB \quad (9.231)$$

From this we obtain

$$[C, AB] = -[AB, C] = A[C, B] + [C, A]B \quad (9.232)$$

We also see that we have

$$[AB, C] = [A, C]B \quad \text{if } [B, C] = 0 \quad (9.233)$$

$$[AB, C] = A[B, C] \quad \text{if } [A, C] = 0 \quad (9.234)$$

### Commutators between components of $\hat{L}$

Let us use these properties to compute

$$\begin{aligned} [\hat{L}_x, \hat{L}_y] &= [\hat{y}\hat{p}_z - \hat{z}\hat{p}_y, \hat{z}\hat{p}_x - \hat{x}\hat{p}_z] = [\hat{y}\hat{p}_z - \hat{z}\hat{p}_y, \hat{z}\hat{p}_x] - [\hat{y}\hat{p}_z - \hat{z}\hat{p}_y, \hat{x}\hat{p}_z] \\ &= [\hat{y}\hat{p}_z, \hat{z}\hat{p}_x] - [\hat{z}\hat{p}_y, \hat{z}\hat{p}_x] - [\hat{y}\hat{p}_z, \hat{x}\hat{p}_z] + [\hat{z}\hat{p}_y, \hat{x}\hat{p}_z] \end{aligned} \quad (9.235)$$

Further applying the rules, we see that

$$[\hat{y}\hat{p}_z, \hat{z}\hat{p}_x] = \hat{y}[\hat{p}_z, \hat{z}\hat{p}_x] = \hat{y}[\hat{p}_z, \hat{z}]\hat{p}_x = -i\hbar\hat{y}\hat{p}_x \quad (9.236)$$

since  $\hat{y}$  commutes with  $\hat{z}\hat{p}_x$  by commuting with both operators

$$[\hat{y}, \hat{z}\hat{p}_x] = [\hat{y}, \hat{z}]\hat{p}_x = 0 \quad (9.237)$$

and so on. Similarly, you may see that

$$[\hat{z}\hat{p}_y, \hat{x}\hat{p}_z] = 0 = [\hat{z}\hat{p}_y, \hat{x}\hat{p}_z], \quad [\hat{z}\hat{p}_y, \hat{x}\hat{p}_z] = \hat{x}\hat{p}_y[\hat{z}, \hat{p}_z] = i\hbar\hat{x}\hat{p}_y \quad (9.238)$$

and finally

$$[\hat{L}_x, \hat{L}_y] = i\hbar(\hat{x}\hat{p}_y + \hat{y}\hat{p}_x) \quad (9.239)$$

Working case by case, we may show

$$[\hat{L}_j, \hat{L}_k] = i\varepsilon_{l,j,k}\hat{L}_l \quad (9.240)$$



With some formal manipulation, anyway, we may prove the above formula directly. We need the expression

$$\sum_i \varepsilon_{i,j,k} \varepsilon_{i,l,m} = \delta_{j,l} \delta_{k,m} - \delta_{j,m} \delta_{k,l} \quad (9.241)$$

Indeed assume  $j = l$  and  $k = m$ , with  $j \neq k$ . On the right side we get 1, and on the left we get (since we cannot have repetitions in the  $\varepsilon$  indexes)

$$\varepsilon_{i,j,k} \varepsilon_{i,j,k} \text{ with } i \neq j, i \neq k \quad (9.242)$$

which is either  $1^2$  or  $(-1)^2$ . If  $j = m$  and  $k = l$ , again with  $j \neq k$ , on the right side we have -1 and on the left

$$\varepsilon_{i,j,k} \varepsilon_{i,k,j} \text{ with } i \neq j, i \neq k \quad (9.243)$$

which is necessarily -1 due to the  $k, j$  inversion. If we have  $j = k$  we have

$$\sum_i \varepsilon_{i,j,j} \varepsilon_{i,l,m} = 0 = \delta_{j,l} \delta_{j,m} - \delta_{j,m} \delta_{j,l} \quad (9.244)$$

Finally, if  $j \neq l \neq k$ , (and  $j \neq k$ , since for  $j = k$  we already shown that we get a zero)

$$\delta_{j,l} \delta_{j,m} - \delta_{j,m} \delta_{k,l} = 0 \quad (9.245)$$

but also

$$\sum_i \varepsilon_{i,j,k} \varepsilon_{i,l,m} = 0 \quad (9.246)$$

since it is impossible to find  $i$  such that  $i \neq j, i \neq k, i \neq l$  in order to avoid repetitions in the indexes. The same applies if we replace  $l$  with  $m$ . In this way we show that eq. (9.241) applies.

Once we got this useful expression, we may show, applying multiple times

## 9.C Angular momentum in Classical and Quantum Mechanics

the commutator properties, that

$$\begin{aligned}
 [\hat{L}_j, \hat{L}_k] &= \left[ \sum_{l,m} \varepsilon_{j,l,m} \hat{x}_l \hat{p}_m, \sum_{n,q} \varepsilon_{k,n,q} \hat{x}_n \hat{p}_q \right] = \sum_{l,m,n,q} \varepsilon_{j,l,m} \varepsilon_{k,n,q} [\hat{x}_l \hat{p}_m, \hat{x}_n \hat{p}_q] = \\
 &\sum_{l,m,n,q} \varepsilon_{j,l,m} \varepsilon_{k,n,q} (\hat{x}_l [\hat{p}_m, \hat{x}_n \hat{p}_q] + [\hat{x}_l, \hat{x}_n \hat{p}_q] \hat{p}_m) = i\hbar \sum_{l,m,n,q} \varepsilon_{j,l,m} \varepsilon_{k,n,q} (-\hat{x}_l \delta_{n,m} \hat{p}_q + \hat{x}_n \delta_{l,q} \hat{p}_m) = \\
 &i\hbar \left( - \sum_{l,m,q} \varepsilon_{j,l,m} \varepsilon_{k,m,q} \hat{x}_l \hat{p}_q + \sum_{l,m,n} \varepsilon_{j,l,m} \varepsilon_{k,n,l} \hat{x}_n \hat{p}_m \right) = \\
 &i\hbar \left( - \sum_{l,m,q} \varepsilon_{m,j,l} \varepsilon_{m,q,k} \hat{x}_l \hat{p}_q + \sum_{l,m,n} \varepsilon_{l,m,j} \varepsilon_{l,k,n} \hat{x}_n \hat{p}_m \right) = \\
 &i\hbar \left( - \sum_{l,q} \delta_{j,q} \delta_{l,k} \hat{x}_l \hat{p}_q + \sum_{l,q} \delta_{j,k} \delta_{l,q} \hat{x}_l \hat{p}_q + \sum_{m,n} \delta_{m,k} \delta_{j,n} \hat{x}_n \hat{p}_m - \sum_{m,n} \delta_{m,n} \delta_{j,k} \hat{x}_n \hat{p}_m \right)
 \end{aligned} \tag{9.247}$$

Now, the sum of the second and fourth term is zero, as can be seen just by changing the names of the dummy indexes in the latter ( $m \rightarrow q$ ,  $n \rightarrow l$ )

$$\begin{aligned}
 \sum_{l,q} \delta_{j,k} \delta_{l,q} \hat{x}_l \hat{p}_q - \sum_{m,n} \delta_{m,n} \delta_{j,k} \hat{x}_n \hat{p}_m &= \sum_{l,q} \delta_{j,k} \delta_{l,q} \hat{x}_l \hat{p}_q - \sum_{q,l} \delta_{q,l} \delta_{j,k} \hat{x}_l \hat{p}_q = \\
 \sum_{l,q} \delta_{j,k} \delta_{l,q} \hat{x}_l \hat{p}_q - \sum_{l,q} \delta_{j,k} \delta_{l,q} \hat{x}_l \hat{p}_q &= 0
 \end{aligned} \tag{9.248}$$

Now we obtain, from the remaining terms, changing again  $m \rightarrow q$ ,  $n \rightarrow l$  in the latter, and recalling that sums and deltas do not depend on the index order, and using once again eq. (9.241) in the right-left direction

$$\begin{aligned}
 [\hat{L}_j, \hat{L}_k] &= i\hbar \left( - \sum_{l,q} \delta_{j,q} \delta_{l,k} \hat{x}_l \hat{p}_q + \sum_{m,n} \delta_{m,k} \delta_{j,n} \hat{x}_n \hat{p}_m \right) = i\hbar \left( - \sum_{l,q} \delta_{j,q} \delta_{l,k} \hat{x}_l \hat{p}_q + \sum_{q,l} \delta_{q,k} \delta_{j,l} \hat{x}_l \hat{p}_q \right) \\
 &i\hbar \left( - \sum_{l,q} \delta_{j,q} \delta_{l,k} + \sum_{q,l} \delta_{q,k} \delta_{j,l} \right) \hat{x}_l \hat{p}_q = i\hbar \sum_{q,l} (\delta_{q,k} \delta_{l,j} - \delta_{q,j} \delta_{l,k}) \hat{x}_l \hat{p}_q = \\
 &i\hbar \sum_{r,q,l} \varepsilon_{r,q,l} \varepsilon_{r,k,j} \hat{x}_l \hat{p}_q = i\hbar \sum_r \varepsilon_{r,k,j} \left( \sum_{q,l} \varepsilon_{r,q,l} \hat{x}_l \hat{p}_q \right) = i\hbar \sum_r \varepsilon_{r,k,j} L_r
 \end{aligned} \tag{9.249}$$

i.e. eq. (9.240).

We can also obtain, much more easily

$$\begin{aligned}
 [\hat{L}_j, \hat{x}_k] &= \left[ \sum_{l,m} \varepsilon_{j,l,m} \hat{x}_l \hat{p}_m, \hat{x}_k \right] = \sum_{l,m} \varepsilon_{j,l,m} \hat{x}_l [\hat{p}_m, \hat{x}_k] = \\
 &- i\hbar \sum_{l,m} \varepsilon_{j,l,m} \hat{x}_l \delta_{m,k} = -i\hbar \sum_l \varepsilon_{j,l,k} \hat{x}_l = i\hbar \sum_l \varepsilon_{j,k,l} \hat{x}_l
 \end{aligned} \tag{9.250}$$

and

$$\begin{aligned}
 [\hat{L}_j, \hat{p}_k] &= \left[ \sum_{l,m} \varepsilon_{j,l,m} \hat{x}_l \hat{p}_m, \hat{p}_k \right] = \sum_{l,m} \varepsilon_{j,l,m} [\hat{x}_l, \hat{p}_k] \hat{p}_m = \\
 &i\hbar \sum_{l,m} \varepsilon_{j,l,m} \hat{p}_m \delta_{l,k} = i\hbar \sum_m \varepsilon_{j,k,m} \hat{p}_m
 \end{aligned} \tag{9.251}$$

We may see that the commutators of  $\hat{L}$  with the same  $\hat{L}$ ,  $\hat{p}$  and  $\hat{x}$  all satisfy the rule

$$[\hat{L}_j, \hat{A}_k] = i\hbar \sum_l \varepsilon_{j,k,l} \hat{A}_l \tag{9.252}$$

Using *Group Theory*, it may be shown that this is due to the fact that all these operators are vectors, and rotate as vectors under the  $L$ , which are the generators of rotations. We may also notice that different components of  $L$  do not commute between themselves, since rotations do not commute. These properties of Quantum Mechanics go anyway beyond the space that we have in this course.

It is nevertheless important to remember that a scalar, such as for example the (square of) the magnitude of  $\mathbf{L}$ , is left unchanged under rotations, and thus we expect  $\hat{L}^2$  to commute with all  $\hat{L}_j$ . Indeed,

$$\begin{aligned}
 [\hat{L}_j, \hat{L}^2] &= [\hat{L}_j, \sum_k \hat{L}_k^2] = \sum_k [\hat{L}_j, \hat{L}_k \hat{L}_k] = \sum_k \left( [\hat{L}_j, \hat{L}_k] \hat{L}_k + \hat{L}_k [\hat{L}_j, \hat{L}_k] \right) = \\
 &i\hbar \left( \sum_{k,l} \varepsilon_{j,k,l} \hat{L}_l \hat{L}_k + \sum_{k,l} \varepsilon_{j,k,l} \hat{L}_k \hat{L}_l \right)
 \end{aligned} \tag{9.253}$$

Let us now switch the names of the  $k$  and  $l$  indexes in the last sum, and then use the asymmetry of  $\varepsilon$

$$[\hat{L}_j, \hat{L}^2] = \dots = i\hbar \left( \sum_{k,l} \varepsilon_{j,k,l} \hat{L}_l \hat{L}_k + \sum_{k,l} \varepsilon_{j,l,k} \hat{L}_l \hat{L}_k \right) = i\hbar \left( \sum_{k,l} \varepsilon_{j,k,l} \hat{L}_l \hat{L}_k - \sum_{k,l} \varepsilon_{j,k,l} \hat{L}_l \hat{L}_k \right) = 0 \tag{9.254}$$

## 9.C Angular momentum in Classical and Quantum Mechanics

---

If you are baffled by formal computations, you may always deal with explicit ones such as

$$[\hat{L}_z, \hat{L}_x^2 + \hat{L}_y^2 + \hat{L}_z^2] = [\hat{L}_z, \hat{L}_x^2] + [\hat{L}_z, \hat{L}_y^2] = \hat{L}_x[\hat{L}_z, \hat{L}_x] + [\hat{L}_z, \hat{L}_x]\hat{L}_x + \hat{L}_y[\hat{L}_z, \hat{L}_y] + [\hat{L}_z, \hat{L}_y]\hat{L}_y = i\hbar(L_xL_y + L_yL_x - L_yL_x - L_xL_y) = 0 \quad (9.255)$$

The operator  $\hat{L}^2$  is self-adjoint, since

$$(\hat{L}_i^2)^\dagger = (\hat{L}_i\hat{L}_i)^\dagger = \hat{L}_i^\dagger\hat{L}_i^\dagger = \hat{L}_i\hat{L}_i = \hat{L}_i^2 \quad (9.256)$$

and

$$(\hat{L}^2)^\dagger = (\hat{L}_x^2 + \hat{L}_y^2 + \hat{L}_z^2)^\dagger = \hat{L}_x^2 + \hat{L}_y^2 + \hat{L}_z^2 = \hat{L}^2 \quad (9.257)$$

### 9.C.5 The operators $\hat{L}_+$ , $\hat{L}_-$ and their properties

Let us now define

$$\hat{L}_+ = \hat{L}_x + i\hat{L}_y, \quad \hat{L}_- = \hat{L}_x - i\hat{L}_y \quad (9.258)$$

We have

$$[\hat{L}_z, \hat{L}_+] = [\hat{L}_z, \hat{L}_x] + i[\hat{L}_z, \hat{L}_y] = i\hbar(\varepsilon_{3,1,2}\hat{L}_y + i\varepsilon_{3,2,1}\hat{L}_z) = i\hbar\hat{L}_y + \hbar\hat{L}_z = \hbar\hat{L}_+ \quad (9.259)$$

and

$$[\hat{L}_z, \hat{L}_-] = [\hat{L}_z, \hat{L}_x] - i[\hat{L}_z, \hat{L}_y] = i\hbar(\varepsilon_{3,1,2}\hat{L}_y - i\varepsilon_{3,2,1}\hat{L}_z) = i\hbar\hat{L}_y - \hbar\hat{L}_z = -\hbar\hat{L}_- \quad (9.260)$$

and

$$[\hat{L}_+, \hat{L}_-] = [\hat{L}_x + i\hat{L}_y, \hat{L}_x - i\hat{L}_y] = i[\hat{L}_y, \hat{L}_x] - i[\hat{L}_x, \hat{L}_y] = -2i(i)\hat{L}_z = 2\hat{L}_z \quad (9.261)$$

These operators are clearly *not self-adjoint*

$$(\hat{L}_+)^\dagger = (\hat{L}_x + i\hat{L}_y)^\dagger = \hat{L}_x - i\hat{L}_y = \hat{L}_- \quad (9.262)$$

$$(\hat{L}_-)^\dagger = (\hat{L}_x - i\hat{L}_y)^\dagger = \hat{L}_x + i\hat{L}_y = \hat{L}_+ \quad (9.263)$$

A useful relation is the following

$$\hat{L}_+\hat{L}_- + \hat{L}_-\hat{L}_+ = 2\hat{L}_x^2 + 2\hat{L}_y^2 + i\hat{L}_y\hat{L}_x - i\hat{L}_x\hat{L}_y - i\hat{L}_y\hat{L}_x + i\hat{L}_x\hat{L}_y = 2(\hat{L}_x^2 + \hat{L}_y^2) \quad (9.264)$$

from which we obtain

$$\hat{L}^2 = \frac{1}{2}(\hat{L}_+\hat{L}_- + \hat{L}_-\hat{L}_+) + \hat{L}_z^2 \quad (9.265)$$

Using the commuting properties of  $\hat{L}_\pm$  we can re-write this as

$$\begin{aligned}\hat{L}^2 &= \frac{1}{2} \left( \hat{L}_+ \hat{L}_- + \hat{L}_+ \hat{L}_- - \hat{L}_+ \hat{L}_- + \hat{L}_- \hat{L}_+ \right) + \hat{L}_z^2 = \\ &\frac{1}{2} \left( 2\hat{L}_+ \hat{L}_- + [\hat{L}_-, \hat{L}_+] \right) + \hat{L}_z^2 = \hat{L}_+ \hat{L}_- - \hbar \hat{L}_z + \hat{L}_z^2\end{aligned}\quad (9.266)$$

or

$$\begin{aligned}\hat{L}^2 &= \frac{1}{2} \left( \hat{L}_+ \hat{L}_- + \hat{L}_- \hat{L}_+ - \hat{L}_- \hat{L}_+ + \hat{L}_- \hat{L}_+ \right) + \hat{L}_z^2 = \\ &\frac{1}{2} \left( 2\hat{L}_- \hat{L}_+ + [\hat{L}_+, \hat{L}_-] \right) + \hat{L}_z^2 = \hat{L}_- \hat{L}_+ + \hbar \hat{L}_z + \hat{L}_z^2\end{aligned}\quad (9.267)$$

### 9.C.6 Eigenvectors and eigenvalues of $\hat{L}$

In appendix 9.D we are going to show that

$$\hat{L}_z = -i\hbar\partial_\varphi \quad (9.268)$$

Considering that for each function we have  $f(\varphi) = f(\varphi + 2\pi)$ , the eigenvectors of this operator are

$$|m\rangle \equiv \frac{1}{\sqrt{2\pi}} e^{im\phi}, \quad \hat{L}_z |m\rangle = \hbar |m\rangle, \quad m \in (0, \mathbb{N}) \quad (9.269)$$

We choose the normalization so that (section 2.8.3)

$$\langle n|m\rangle = \delta_{n,m} \quad (9.270)$$

Since

$$[\hat{L}_z, \hat{L}_x] \neq 0 \neq [\hat{L}_z, \hat{L}_y] \quad (9.271)$$

the eigenvectors  $|m\rangle$  are not in general eigenvectors of  $\hat{L}_x$  and  $\hat{L}_y$ , so that a physical state with a well defined value of  $\mathbf{L}$  in the  $z$  direction has not a well defined value of  $\hat{L}_x$  and  $\hat{L}_y$ , i.e. any attempt to measure these observables will give a result with a finite variance, different repetitions of the experiment under the same conditions giving a spread of values<sup>14</sup>.

Let us now remember that  $\hat{L}^2$  and  $\hat{L}_z$  are commuting self adjoint operators (eq. 9.254), so that they admit a common basis of orthonormal eigenvectors (eq. 6.106). Let us call these vectors  $|a, m\rangle$

$$\hat{L}^2 |a, m\rangle = a |a, m\rangle \quad \hat{L}_z |a, m\rangle = m |a, m\rangle \quad (9.272)$$

---

<sup>14</sup>The state  $|0\rangle$  is an exception, being an eigenvector of all  $\hat{L}$  components. The  $\hat{L}$  components may commute when acting on some vector, a situation different from the one we have for the  $[\hat{x}, \hat{p}] = i\hbar\mathbf{1}$  commutator. See also the discussion in appendix 9.E.

### 9.C Angular momentum in Classical and Quantum Mechanics

---

We already know (eq. 9.269) that  $m \in (0, \mathbb{N})$ . It is trivial to show that  $a \geq 0$

$$a = \langle a, m | \hat{L}^2 | a, m \rangle = \sum_i \langle a, m | \hat{L}_i^2 | a, m \rangle = \sum_i \langle a, m | \hat{L}_i | \hat{L}_i | a, m \rangle = \sum_i \| \hat{L}_i | a, m \rangle \|^2 \geq 0 \quad (9.273)$$

We can even show something more, since

$$a = \langle a, m | \hat{L}^2 | a, m \rangle = \| \hat{L}_x | a, m \rangle \|^2 + \| \hat{L}_y | a, m \rangle \|^2 + \| \hat{L}_z | a, m \rangle \|^2 \quad (9.274)$$

but since  $\hat{L}_z | a, m \rangle = \hbar | a, m \rangle$

$$\langle a, m | \hat{L}_z | \hat{L}_z | a, m \rangle = \hbar^2 m^2 \langle a, m | a, m \rangle = \hbar^2 m^2 \quad (9.275)$$

and thus

$$a = \| \hat{L}_x | a, m \rangle \|^2 + \| \hat{L}_y | a, m \rangle \|^2 + \hbar^2 m^2 \geq \hbar^2 m^2 \quad (9.276)$$

or

$$\sqrt{a} \geq \hbar |m| \quad (9.277)$$

As shown above, we have also

$$[\hat{L}_z, \hat{L}_\pm] \neq 0 \quad (9.278)$$

but the action of these operators on the  $|a, m\rangle$  eigenvectors is extremely simple (and that is the reason we introduced them). Indeed we have

$$\begin{aligned} \hat{L}_z \hat{L}_+ | a, m \rangle &= (\hat{L}_z \hat{L}_+ - \hat{L}_+ \hat{L}_z + \hat{L}_+ \hat{L}_z) | a, m \rangle = ([\hat{L}_z, \hat{L}_+] + \hat{L}_+ \hat{L}_z) | a, m \rangle \\ &= (\hbar \hat{L}_+ + m \hbar \hat{L}_+) | a, m \rangle = \hbar(m+1) (\hat{L}_+ | a, m \rangle) \end{aligned} \quad (9.279)$$

This means that when acting on  $\hat{L}_+ | a, m \rangle$ ,  $\hat{L}_z$  multiplies it by  $\hbar(m+1)$ , i.e.  $\hat{L}_+ | a, m \rangle$  is an eigenvector of  $\hat{L}_z$

$$\hat{L}_+ | a, m \rangle \propto |m+1\rangle \quad (9.280)$$

But it has also to be an eigenvector of  $\hat{L}^2$  with eigenvalue  $a$  since  $[\hat{L}^2, \hat{L}_+] = 0$

$$\hat{L}^2 \hat{L}_+ | a, m \rangle = \hat{L}_+ a | a, m \rangle = a \hat{L}_+ | a, m \rangle \quad (9.281)$$

and thus

$$\hat{L}_+ | a, m \rangle \propto |a, m+1\rangle \quad (9.282)$$

Similarly

$$\begin{aligned} \hat{L}_z \hat{L}_- |a, m\rangle &= (\hat{L}_z \hat{L}_- - \hat{L}_- \hat{L}_z + \hat{L}_- \hat{L}_z) |a, m\rangle = ([\hat{L}_z, \hat{L}_-] + \hat{L}_- \hat{L}_z) |a, m\rangle \\ &= (-\hbar \hat{L}_+ + m\hbar \hat{L}_+) |a, m\rangle = \hbar(m-1) (\hat{L}_+ |a, m\rangle) \end{aligned} \quad (9.283)$$

so that

$$\hat{L}_- |m\rangle \propto |a, m-1\rangle \quad (9.284)$$

If we now consider  $m$  such that  $\hbar m \geq -\sqrt{a}$  and  $n \in \mathbb{N}$ , with  $\hbar n > \sqrt{a}$  we get

$$\hat{L}_+^n |a, m\rangle = |a, m+n\rangle \quad (9.285)$$

But since

$$\hbar^2(m+n)^2 > a \quad (9.286)$$

we have built an eigenvector whose eigenvalue is larger than  $\hbar\sqrt{a}$  in contradiction with what stated above. How is this possible?

We know that

$$\hat{L}_z(\hat{L}_+ |a, m\rangle) = \hbar(m+1)\hat{L}_+ |a, m\rangle \quad (9.287)$$

This actually implies that

$$\hat{L}_+ |a, m\rangle \propto |a, m+1\rangle \quad (9.288)$$

unless

$$\hat{L}_+ |a, m\rangle = 0 \quad (9.289)$$

Thus, to avoid the contradiction, there must be a  $m_{\text{MAX}}$  such that

$$\hat{L}_+ |a, m_{\text{MAX}}\rangle = 0 \quad (9.290)$$

We may choose instead  $m \leq \sqrt{a}$  as our starting vector and  $n \in \mathbb{N}$ , with  $\hbar n > \sqrt{a}$  to obtain

$$\hat{L}_-^n |a, m\rangle = |a, m-n\rangle \quad (9.291)$$

Again, since

$$\hbar^2(m-n)^2 > a \quad (9.292)$$

there must be also a  $m_{\text{MIN}}$  such that

$$\hat{L}_- |a, m_{\text{MIN}}\rangle = 0 \quad (9.293)$$

## 9.C Angular momentum in Classical and Quantum Mechanics

---

We now see that (refer to eq. 9.267)

$$a = \langle a, m_{\text{MAX}} | \hat{L}^2 | a, m_{\text{MAX}} \rangle = \langle a, m_{\text{MAX}} | \hat{L}_- \hat{L}_+ + \hbar \hat{L}_z + \hat{L}_z^2 | a, m_{\text{MAX}} \rangle = 0 + \hbar^2 m_{\text{MAX}} + \hbar^2 m_{\text{MAX}}^2 \quad (9.294)$$

or

$$a = m_{\text{MAX}}(m_{\text{MAX}} + 1) \quad (9.295)$$

Similarly (refer to eq. 9.266)

$$a = \langle a, m_{\text{MIN}} | \hat{L}^2 | a, m_{\text{MIN}} \rangle = \langle a, m_{\text{MIN}} | \hat{L}_+ \hat{L}_- - \hbar \hat{L}_z + \hat{L}_z^2 | a, m_{\text{MIN}} \rangle = 0 - \hbar^2 m_{\text{MIN}} + \hbar^2 m_{\text{MIN}}^2 \quad (9.296)$$

or

$$a = \hbar^2 m_{\text{MIN}}(m_{\text{MIN}} - 1) \quad (9.297)$$

To find the relation between  $m_{\text{MAX}}$  and  $m_{\text{MIN}}$  we solve

$$x^2 - x = y^2 + y \Rightarrow x^2 - y^2 = x + y \Rightarrow (x + y)(x - y) = x + y \quad (9.298)$$

The solution

$$x - y = 1 \Rightarrow x = y + 1 \quad (9.299)$$

cannot be accepted (the minimum would be higher than the maximum) thus we have

$$x + y = 0 \Rightarrow m_{\text{MIN}} = -m_{\text{MAX}} \quad (9.300)$$

Let us call

$$m_{\text{MAX}} = l \in (0, \mathbb{N}) \quad (9.301)$$

since all the  $m$  are integers. We now have

$$a = \hbar^2 l(l + 1), m = -l, -l + 1, \dots, l - 1, l \quad (9.302)$$

But these are exactly the eigenvalues admitted by the spherical harmonics, a result that we had previously presented without proof.

We may thus identify

$$|\hbar^2 l(l + 1), m\rangle = Y_{l,m}(\theta, \varphi) \quad (9.303)$$

In general, these eigenvectors are identified simply as

$$|l, m\rangle = Y_{l,m}(\theta, \varphi) \quad (9.304)$$



A spherical harmonic corresponds thus to a physical state in which the angular momentum assumes magnitude  $L^2 = \hbar^2 l(l+1)$  while its  $z$  component has value  $L_z = \hbar m$

We know that

$$|l, 0\rangle \propto P_l(\cos \theta) \quad (9.305)$$

we may then use the operators  $\hat{L}_\pm$  (see appendix 9.D for their explicit form) to build

$$\hat{L}_+^n |l, 0\rangle \propto |l, n\rangle, \quad \hat{L}_-^n |l, 0\rangle \propto |l, -n\rangle \quad (9.306)$$

Nevertheless, if we want orthonormality

$$\langle l, m | l', m' \rangle = \delta_{l,l'} \delta_{m,m'} \quad (9.307)$$

we need also to compute the proportionality constant. We may do it as follows

$$\begin{aligned} \hbar^2 l(l+1) &= \langle l, m | \hat{L}^2 | l, m \rangle = \langle l, m | \hat{L}_- \hat{L}_+ | l, m \rangle + \hbar^2 (m + m^2) = \\ \Rightarrow \hbar^2 (l(l+1) - m(m+1)) &= \sum_{l'} \sum_{m=-l'}^{l'} \langle l, m | \hat{L}_- | l', m' \rangle \langle l', m' | \hat{L}_+ | l, m \rangle \end{aligned} \quad (9.308)$$

In the last step we used the fact that the common eigenvectors of  $\hat{L}^2$  and  $\hat{L}_z$  are a basis<sup>15</sup>, so that (eq. 6.140)

$$\sum_{l'} \sum_{m=-l'}^{l'} |l', m'\rangle \langle l', m'| = \mathbb{1} \quad (9.309)$$

a very common technique in Quantum Mechanics. But since

$$\hat{L}_+ |l, m\rangle \propto |l, m+1\rangle, \quad \hat{L}_- |l, m\rangle \propto |l, m-1\rangle \quad (9.310)$$

and

$$\langle l, m | l', m' \rangle = \delta_{l,l'} \delta_{m,m'} \quad (9.311)$$

the only surviving term of the form

$$\langle l, m | \hat{L}_- | l', m' \rangle \quad (9.312)$$

is

$$\langle l, m | \hat{L}_- | l, m+1 \rangle \quad (9.313)$$

---

<sup>15</sup>For functions defined on a sphere of given radius, the spherical harmonics basis.

## 9.C Angular momentum in Classical and Quantum Mechanics

---

and the only surviving term of the form

$$\langle l', m' | \hat{L}_+ | l, m \rangle \quad (9.314)$$

is

$$\langle l, m + 1 | \hat{L}_+ | l, m \rangle \quad (9.315)$$

We have thus

$$\begin{aligned} \hbar^2 (l(l+1) - m(m+1)) &= \langle l, m | \hat{L}_- | l, m + 1 \rangle \langle l, m + 1 | \hat{L}_+ | l, m \rangle = \\ &\langle l, m | \hat{L}_+ | l, m + 1 \rangle \langle l, m + 1 | \hat{L}_+ | l, m \rangle = \|\langle l, m + 1 | \hat{L}_+ | l, m \rangle\|^2 \end{aligned} \quad (9.316)$$

where we used  $\hat{L}_-^\dagger = \hat{L}_+$  and  $\langle a | b \rangle = (\langle b | a \rangle)^*$ . Now we know that we have a proportionality law

$$\hat{L}_+ | l, m \rangle = \alpha_m^+ | l, m + 1 \rangle \quad (9.317)$$

so that

$$\|\langle l, m + 1 | \hat{L}_+ | l, m \rangle\|^2 = (\alpha_m^+)^2 \quad (9.318)$$

and we finally find that

$$\alpha_m^+ = e^{i\beta} \hbar \sqrt{l(l+1) - m(m+1)} \quad (9.319)$$

The complex phase  $e^{i\beta}$  is an arbitrary convention and may be fixed to 1. We see that since

$$\hat{L}_+ | l, m \rangle = \sqrt{l(l+1) - m(m+1)} | l, m + 1 \rangle \quad (9.320)$$

we have as wanted

$$\hat{L}_+ | l, l \rangle = 0 \quad (9.321)$$

In a similar way, we have

$$\begin{aligned} \hbar^2 l(l+1) &= \langle l, m | \hat{L}^2 | l, m \rangle = \langle l, m | \hat{L}_+ \hat{L}_- | l, m \rangle + \hbar^2 (-m + m^2) = \\ &\Rightarrow \hbar^2 (l(l+1) - m(m-1)) = \sum_{l'} \sum_{m=-l'}^{l'} \langle l, m | \hat{L}_+ | l', m' \rangle \langle l', m' | \hat{L}_- | l, m \rangle = \\ &\langle l, m | \hat{L}_+ | l, m - 1 \rangle \langle l, m - 1 | \hat{L}_- | l, m \rangle = \langle l, m | \hat{L}_- | l, m + 1 \rangle \langle l, m + 1 | \hat{L}_+ | l, m \rangle = \\ &\|\langle l, m - 1 | \hat{L}_+ | l, m \rangle\|^2 \end{aligned} \quad (9.322)$$

From

$$\hat{L}_- | l, m \rangle = \alpha_m^- | l, m - 1 \rangle \quad (9.323)$$

we get

$$\alpha_m^- = e^{i\beta} \hbar \sqrt{l(l+1) - m(m-1)} \quad (9.324)$$

Fixing again the phase to one we obtain

$$\hat{L}_-|l, m\rangle = \sqrt{l(l+1) - m(m-1)}|l, m-1\rangle \quad (9.325)$$

If we act on  $|l, -l\rangle$  we have

$$\hat{L}_+|l, -l\rangle = \sqrt{l(l+1) - (-l)(-l-1)}|l, -l-1\rangle = 0 \quad (9.326)$$

## 9.D Explicit form of $\hat{L}^2$ in spherical coordinates

Deriving eq. (9.197) is not conceptually difficult but a little lengthy.

### 9.D.1 Change of coordinate

Let us start by writing down the change of coordinates laws

#### From spherical to Cartesian

$$z = r \cos \theta \quad (9.327)$$

$$x = r \sin \theta \cos \varphi \quad (9.328)$$

$$y = r \sin \theta \sin \varphi \quad (9.329)$$

#### From Cartesian to spherical

These ones are a little bit more annoying

$$r = \sqrt{x^2 + y^2 + z^2} \quad (9.330)$$

$$\cos \theta = \frac{z}{r} \Rightarrow \theta = \arccos \left( \frac{z}{\sqrt{x^2 + y^2 + z^2}} \right) \quad (9.331)$$

$$\tan \varphi = \frac{y}{x} \Rightarrow \varphi = \operatorname{atan} \frac{y}{x} \quad (9.332)$$

### 9.D.2 Partial derivatives

#### Useful formulae

Let us first recall the derivatives of  $\arccos$

$$\frac{d}{dx} \arccos w = \frac{-1}{\sqrt{1-w^2}} \quad (9.333)$$

and  $\operatorname{atan}$

$$\frac{d}{dx} \operatorname{atan} w = \frac{1}{1+w^2} \quad (9.334)$$

## 9.D Explicit form of $\hat{L}^2$ in spherical coordinates

---

### Derivatives of $\theta$

We have

$$\begin{aligned} \frac{\partial \theta}{\partial x} &= \frac{-1}{\sqrt{1 - \frac{z^2}{r^2}}} z \frac{\partial}{\partial x} \left( \frac{1}{r} \right) = \frac{-1}{\sqrt{1 - \frac{z^2}{r^2}}} z \left( \frac{-1}{r^2} \right) \frac{\partial}{\partial x} \left( \sqrt{x^2 + y^2 + z^2} \right) = \\ &= \frac{1}{\sqrt{1 - \frac{z^2}{r^2}}} z \left( \frac{1}{r^2} \right) \frac{1}{2} \frac{2x}{r} = \frac{zx}{\sqrt{1 - \cos^2 \theta} r^3} = \frac{r^2 \cos \theta \sin \theta \cos \varphi}{\sqrt{1 - \cos^2 \theta} r^3} = \frac{\cos \theta \sin \theta \cos \varphi}{\sin \theta r} \end{aligned} \quad (9.335)$$

So that

$$\partial_x \theta = \frac{\cos \theta \cos \varphi}{r} \quad (9.336)$$

Proceeding similarly, although not writing passages very similar to those performed above, we have

$$\frac{\partial \theta}{\partial y} = \frac{zy}{\sin \theta r^3} = \frac{r^2 \cos \theta \sin \theta \sin \varphi}{\sin \theta r^3} \quad (9.337)$$

So that

$$\partial_y \theta = \frac{\cos \theta \sin \varphi}{r} \quad (9.338)$$

Finally

$$\begin{aligned} \frac{\partial \theta}{\partial z} &= \frac{-1}{\sqrt{1 - \frac{z^2}{r^2}}} \left( \frac{1}{r} - \frac{z}{r^2} \frac{\partial r}{\partial z} \right) = \frac{-1}{\sqrt{1 - \frac{z^2}{r^2}}} \left( \frac{1}{r} - \frac{z^2}{r^3} \right) = \\ &= \frac{-1}{\sin \theta} \frac{r^2 - z^2}{r^3} = \frac{-1}{r \sin \theta} \left( 1 - \frac{z^2}{r^2} \right) = \frac{-1}{r \sin \theta} (1 - \cos^2 \theta) = \frac{-1}{r \sin \theta} \sin^2 \theta \end{aligned} \quad (9.339)$$

so that

$$\partial_z \theta = \frac{-\sin \theta}{r} \quad (9.340)$$

### Derivatives of $r$

These are the easiest ones

$$\frac{\partial r}{\partial x} = \frac{1}{2} \frac{2x}{\sqrt{x^2 + y^2 + z^2}} = \frac{x}{r} = \frac{r \sin \theta \cos \varphi}{r} \quad (9.341)$$

so that

$$\partial_x r = \sin \theta \cos \varphi \quad (9.342)$$

Similarly

$$\frac{\partial r}{\partial y} = \frac{1}{2} \frac{2y}{\sqrt{x^2 + y^2 + z^2}} = \frac{y}{r} = \frac{r \sin \theta \sin \varphi}{r} \quad (9.343)$$

so that

$$\partial_y r = \sin \theta \sin \varphi \quad (9.344)$$

and

$$\frac{\partial r}{\partial z} = \frac{1}{2} \frac{2z}{\sqrt{x^2 + y^2 + z^2}} = \frac{z}{r} = \frac{r \cos \theta}{r} \quad (9.345)$$

so that

$$\partial_z r = \cos \theta \quad (9.346)$$

### Derivatives of $\varphi$

Trivially,

$$\partial_z \varphi = 0 \quad (9.347)$$

Then

$$\frac{\partial \varphi}{\partial x} = \frac{1}{1 + \frac{y^2}{x^2}} \frac{-y}{x^2} = \frac{x^2}{x^2 + y^2} \frac{-y}{x^2} = \frac{-y}{x^2 + y^2} = -\frac{r \sin \theta \sin \varphi}{r^2 \sin^2 \theta} \quad (9.348)$$

so that

$$\partial_x \varphi = \frac{-\sin \varphi}{r \sin \theta} \quad (9.349)$$

and

$$\frac{\partial \varphi}{\partial y} = \frac{1}{1 + \frac{y^2}{x^2}} \frac{1}{x} = \frac{x^2}{x^2 + y^2} \frac{1}{x} = \frac{x}{x^2 + y^2} = \frac{r \sin \theta \cos \varphi}{r^2 \sin^2 \theta} \quad (9.350)$$

so that

$$\partial_y \varphi = \frac{\cos \varphi}{r \sin \theta} \quad (9.351)$$

### 9.D.3 Components of $\hat{L}$

#### Explicit form of $\hat{L}_z$

We have

$$\begin{aligned} \hat{L}_z &= -i\hbar (x\partial_y - y\partial_x) = \\ &= -i\hbar \left( r \sin \theta \cos \varphi \left( \frac{\partial \theta}{\partial y} \partial_\theta + \frac{\partial \varphi}{\partial y} \partial_\varphi + \frac{\partial r}{\partial y} \partial_r \right) - r \sin \theta \sin \varphi \left( \frac{\partial \theta}{\partial x} \partial_\theta + \frac{\partial \varphi}{\partial x} \partial_\varphi + \frac{\partial r}{\partial x} \partial_r \right) \right) = \\ &= -i\hbar \left( r \sin \theta \cos \varphi \frac{\cos \theta \sin \varphi}{r} \partial_\theta + r \sin \theta \cos \varphi \frac{\cos \varphi}{r \sin \theta} \partial_\varphi + r \sin \theta \cos \varphi \sin \theta \sin \varphi \partial_r + \right. \\ &\quad \left. - r \sin \theta \sin \varphi \frac{\cos \theta \cos \varphi}{r} \partial_\theta + r \sin \theta \sin \varphi \frac{\sin \varphi}{r \sin \theta} \partial_\varphi - r \sin \theta \sin \varphi \sin \theta \cos \varphi \partial_r \right) = \\ &= -i\hbar \left( \sin \theta \cos \varphi \cos \theta \sin \varphi \partial_\theta + \cos^2 \varphi \partial_\varphi + r \sin \theta \cos \varphi \sin \theta \sin \varphi \partial_r + \right. \\ &\quad \left. - \sin \theta \sin \varphi \cos \theta \cos \varphi \partial_\theta + \sin^2 \varphi \partial_\varphi - r \sin \theta \sin \varphi \sin \theta \cos \varphi \partial_r \right) = -i\hbar (\cos^2 \varphi + \sin^2 \varphi) \partial_\varphi \end{aligned} \quad (9.352)$$

## 9.D Explicit form of $\hat{L}^2$ in spherical coordinates

---

so that

$$\hat{L}_z = -i\hbar\partial_\varphi \quad (9.353)$$

### Explicit form of $\hat{L}_x$

We have

$$\begin{aligned} \hat{L}_x &= -i\hbar(y\partial_z - z\partial_y) = \\ &= -i\hbar\left(r\sin\theta\sin\varphi\left(\frac{\partial\theta}{\partial z}\partial_\theta + \frac{\partial\varphi}{\partial z}\partial_\varphi + \frac{\partial r}{\partial z}\partial_r\right) - r\cos\theta\left(\frac{\partial\theta}{\partial y}\partial_\theta + \frac{\partial\varphi}{\partial y}\partial_\varphi + \frac{\partial r}{\partial y}\partial_r\right)\right) = \\ &= -i\hbar\left(-r\sin\theta\sin\varphi\frac{\sin\theta}{r}\partial_\theta + r\sin\theta\sin\varphi\cos\theta\partial_r + \right. \\ &\quad \left. - r\cos\theta\frac{\cos\theta\sin\varphi}{r}\partial_\theta - r\cos\theta\frac{\cos\varphi}{r\sin\theta}\partial_\varphi - r\cos\theta\sin\theta\sin\varphi\partial_r\right) = \\ &= -i\hbar\left(-\sin^2\theta\sin\varphi\partial_\theta + r\sin\theta\sin\varphi\cos\theta\partial_r - \right. \\ &\quad \left. - \cos^2\theta\sin\varphi\partial_\theta - \frac{\cos\theta\cos\varphi}{\sin\theta}\partial_\varphi - r\cos\theta\sin\theta\sin\varphi\partial_r\right) = \\ &= -i\hbar(-\sin\varphi\partial_\theta - \cotan\theta\cos\varphi\partial_\varphi) \end{aligned} \quad (9.354)$$

so that

$$\hat{L}_x = i\hbar(\sin\varphi\partial_\theta + \cotan\theta\cos\varphi\partial_\varphi) \quad (9.355)$$

### Explicit form of $\hat{L}_y$

We have

$$\begin{aligned} \hat{L}_y &= -i\hbar(z\partial_x - x\partial_z) = \\ &= -i\hbar\left(r\cos\theta\left(\frac{\partial\theta}{\partial x}\partial_\theta + \frac{\partial\varphi}{\partial x}\partial_\varphi + \frac{\partial r}{\partial x}\partial_r\right) - r\sin\theta\cos\varphi\left(\frac{\partial\theta}{\partial z}\partial_\theta + \frac{\partial\varphi}{\partial z}\partial_\varphi + \frac{\partial r}{\partial z}\partial_r\right)\right) = \\ &= -i\hbar\left(r\cos\theta\frac{\cos\theta\cos\varphi}{r}\partial_\theta + r\cos\theta\frac{-\sin\varphi}{\sin\theta r}\partial_\theta + r\cos\theta\sin\theta\cos\varphi\partial_r + \right. \\ &\quad \left. + r\sin\theta\cos\varphi\frac{\sin\theta}{r}\partial_\theta - r\sin\theta\cos\varphi\cos\theta\partial_r\right) = \\ &= -i\hbar\left(\cos^2\theta\cos\varphi\partial_\theta - \frac{\cos\theta\sin\varphi}{\sin\theta}\partial_\varphi + r\cos\theta\sin\theta\cos\varphi\partial_r - \right. \\ &\quad \left. + \sin^2\theta\cos\varphi\partial_\theta - r\cos\theta\sin\theta\cos\varphi\partial_r\right) = \\ &= -i\hbar(\cos\varphi\partial_\theta - \cot\theta\sin\varphi\partial_\varphi) \end{aligned} \quad (9.356)$$

so that

$$\hat{L}_y = i\hbar(\cot\theta\sin\varphi\partial_\varphi - \cos\varphi\partial_\theta) \quad (9.357)$$

### Explicit form of $\hat{L}^2$

We first write down

$$\hat{L}_x^2 = -\hbar^2 (\sin^2 \varphi \partial_\theta^2 + \cot^2 \theta \cos^2 \varphi \partial_\varphi^2 + \sin \varphi \cos \varphi \partial_\theta (\cot \theta \partial_\varphi) + \cot \theta \cos \varphi \partial_\varphi (\sin \varphi \partial_\theta)) \quad (9.358)$$

and

$$\hat{L}_y^2 = -\hbar^2 (\cos^2 \varphi \partial_\theta^2 + \cot^2 \theta \sin^2 \varphi \partial_\varphi^2 - \sin \varphi \cos \varphi \partial_\theta (\cot \theta \partial_\varphi) - \cot \theta \sin \varphi \partial_\varphi (\cos \varphi \partial_\theta)) \quad (9.359)$$

Their sum gives

$$\begin{aligned} \hat{L}_x^2 + \hat{L}_y^2 &= -\hbar^2 (\partial_\theta^2 + \cot^2 \theta \partial_\varphi^2 + \cot \theta \cos \varphi \partial_\varphi (\sin \varphi \partial_\theta) - \cot \theta \sin \varphi \partial_\varphi (\cos \varphi \partial_\theta)) = \\ &= -\hbar^2 (\partial_\theta^2 + \cot^2 \theta \partial_\varphi^2 + \cot \theta \cos^2 \varphi \partial_\theta + \cot \theta \sin^2 \varphi \partial_\theta + \cot \theta \cos \varphi \sin \varphi \partial_\varphi \partial_\theta - \cot \theta \cos \varphi \sin \varphi \partial_\theta \partial_\varphi) = \\ &= -\hbar^2 (\partial_\theta^2 + \cot^2 \theta \partial_\varphi^2 + \cot \theta \partial_\theta) \end{aligned} \quad (9.360)$$

And summing  $\hat{L}_z^2$  we get

$$\hat{L}^2 = -\hbar^2 (\partial_\theta^2 + (\cot^2 \theta + 1) \partial_\varphi^2 + \cot \theta \partial_\theta) \quad (9.361)$$

now, since

$$\cot^2 \theta + 1 = \frac{\cos^2 \theta}{\sin^2 \theta} + 1 = \frac{1}{\sin^2 \theta} \quad (9.362)$$

we have

$$\hat{L}^2 = -\hbar^2 \left( \partial_\theta^2 + \cot \theta \partial_\theta + \frac{1}{\sin^2 \theta} \partial_\varphi^2 \right) \quad (9.363)$$

Finally, we see that

$$\partial_\theta (\sin \theta \partial_\theta) = \sin \theta \partial_\theta^2 + \cos \theta \partial_\theta \Rightarrow \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) = \partial_\theta^2 + \cot \theta \partial_\theta \quad (9.364)$$

and thus

$$\hat{L}^2 = -\hbar^2 \left( \frac{1}{\sin \theta} \partial_\theta (\sin \theta \partial_\theta) + \frac{1}{\sin^2 \theta} \partial_\varphi^2 \right) \quad (9.365)$$

## 9.E Mathematical formulation of the uncertainty principle

### 9.E.1 Expectation values

Let us consider a physical state

$$|\Psi\rangle = \sum_i c_i |a_i\rangle \quad (9.366)$$

## 9.E Mathematical formulation of the uncertainty principle

---

The  $c_i$  are complex numbers

$$c_i \in \mathbb{C}, \quad \sum_i |c_i|^2 = 1 \quad (9.367)$$

and the  $|a_i\rangle$  are orthonormal

$$\langle a_j | a_i \rangle = \delta_{j,i} \quad (9.368)$$

so that

$$\langle \Psi | \Psi \rangle = \sum_{i,j} c_i^* c_j \langle a_j | a_i \rangle = \sum_i |c_i|^2 = 1 \quad (9.369)$$

this state is the superposition of states  $|a_i\rangle$  on which the observable  $A$  (for example, a component of the angular momentum) assumes value  $a_i \in \mathbb{R}$ . The  $a_i$  all the possible values assumed by  $A$ <sup>16</sup>. This means that when we perform an experiment to measure  $A$ , the result will always be one of the  $a_i$ . If the physical state is  $|a_i\rangle$ , each experiment will always give  $a_i$  as a result, and this is indeed the physical definition of  $|a_i\rangle$ . In the case of the state  $|\Psi\rangle$ , an experiment is going to give the result  $a_i$  with probability  $p_i = |c_i|^2$ <sup>17</sup>. This means that if we perform  $N$  experiments, we expect (for large  $N$ ) to have the result  $a_i$  for  $n_i$  times, with

$$n_i = |c_i|^2 N \Rightarrow |c_i|^2 = \frac{n_i}{N} = p_i \quad (9.370)$$

In the general case then the value of  $A$  is not fixed. What is uniquely determined is its average value, or *expectation value*

$$\langle A \rangle_\Psi = \sum_i |c_i|^2 a_i = \frac{\sum_i n_i a_i}{N} = \sum_i p_i a_i \in \mathbb{R} \quad (9.371)$$

i.e. a weighted average in which each result is weighted with its probability  $p_i = |c_i|^2$ . The expectation value is clearly linear, i.e. if on the same state (and thus with the same probability  $p_i$ ) the observable  $B$  assumes value  $b_i$ , we have

$$\langle A + \beta B \rangle_\Psi = \sum_i p_i (a_i + \beta b_i) = \sum_i p_i a_i + \beta \sum_i p_i b_i = \langle A \rangle_\Psi + \beta \langle B \rangle_\Psi \quad (9.372)$$

Clearly, the expectation value of a constant is the constant itself

$$\langle \alpha \rangle_\Psi = \sum_i p_i \alpha = \alpha \quad (9.373)$$

---

<sup>16</sup>They may be continuous, although we will assume for simplicity them to be discrete. And indeed, the existence of discrete values for physical observables is one of the features of Quantum Mechanics, as we have seen for the angular momentum and energy.

<sup>17</sup>A large number of experimental results has led physicists to build this formulation of the laws of nature.



It is also important to know what is the spread of  $A$  around its average value  $\langle A \rangle$ . This may be done by measuring

$$\Delta A_{\Psi}^2 \equiv \langle A^2 \rangle_{\Psi} - \langle A \rangle_{\Psi}^2 \quad (9.374)$$

or, equivalently (using the properties of expectation values)

$$\langle (A - \langle A \rangle_{\Psi})^2 \rangle_{\Psi} = \langle A^2 + \langle A \rangle_{\Psi}^2 - 2A \langle A \rangle_{\Psi} \rangle_{\Psi} = \langle A^2 \rangle_{\Psi} + \langle A \rangle_{\Psi}^2 - 2 \langle A \rangle_{\Psi} \langle A \rangle_{\Psi} = \Delta A_{\Psi}^2 \quad (9.375)$$

Since  $(A - \langle A \rangle_{\Psi})^2 \geq 0$ , and obviously all probabilities are  $p_i \geq 0$  we have  $\Delta A_{\Psi}^2 \geq 0$ , with  $\Delta A_{\Psi}^2 = 0$  only if  $A = \langle A \rangle_{\Psi}$ , i.e. if  $|\Psi\rangle = |a_i\rangle$ .

Until now, we have talked about  $A$  as a physical observable, i.e. without any reference to self-adjoint operators. But we assumed that we may write any state as a combination of states in which  $A$  assumes a fixed value. We may thus assume the  $|a_i\rangle$  to be the eigenvectors of the self adjoint operator  $\hat{A}$ , that is the mathematical representation of the physical observable  $A$ <sup>18</sup>. We have thus

$$\hat{A}|a_i\rangle = a_i|a_i\rangle \quad (9.376)$$

or, (eq. 6.151)

$$\hat{A} = \sum_i a_i |a_i\rangle \langle a_i| \quad (9.377)$$

We can see that

$$\begin{aligned} \langle \Psi | \hat{A} | \Psi \rangle &= \left\langle \sum_i c_i a_i |a_i\rangle \left| \sum_j c_j a_j \right. \right\rangle = \sum_i c_i^* \langle a_i | \left| \sum_j c_j a_j \right. \rangle = \sum_{i,j} c_i^* c_j a_j \langle a_i | a_j \rangle = \\ &= \sum_{i,j} c_i^* c_j a_j \delta_{i,j} = \sum_i |c_i|^2 a_i = \langle A \rangle_{\Psi} \end{aligned} \quad (9.378)$$

explaining why we have considered these expressions as equivalent of average values of observables.

We may define the following operator (clearly self-adjoint, since  $\langle A \rangle_{\Psi}$  is just a real number)

$$\hat{\Delta A}_{\Psi} \equiv \hat{A} - \langle A \rangle_{\Psi} \quad (9.379)$$

We have

$$\begin{aligned} \|\hat{\Delta A}_{\Psi} |\Psi\rangle\|^2 &= \langle \Psi | \hat{\Delta A}_{\Psi} | \hat{\Delta A}_{\Psi} | \Psi \rangle = \langle \Psi | (\hat{\Delta A}_{\Psi})^2 | \Psi \rangle = \\ &= \langle \Psi | \hat{A}^2 + \langle A \rangle_{\Psi}^2 - 2\hat{A} \langle A \rangle_{\Psi} | \Psi \rangle = \Delta A_{\Psi}^2 \end{aligned} \quad (9.380)$$

---

<sup>18</sup>We use the expression  $\hat{A}$  just here to make a distinction between observable and operator and clarify some aspects.

## 9.E Mathematical formulation of the uncertainty principle

---

### Uncertainty principle

Now that we have clarified the physical and mathematical meaning of the spread  $\Delta A_\Psi^2$ , we may show the relation (uncertainty principle)

$$\Delta A_\Psi^2 \Delta B_\Psi^2 \geq \frac{1}{4} |\langle [\hat{A}, \hat{B}] \rangle_\Psi|^2 \quad (9.381)$$

that relates the spreads of operators  $A$  and  $B$  on the state  $|\Psi\rangle$  to the expectation value on  $\Psi$  of their commutator.

To show it we need the Schwartz inequality, which is explicitly demonstrated below, but that states the quite intuitive result that the magnitude of a scalar product is smaller than the product of the magnitude of the two vectors

$$|\langle \alpha | \beta \rangle|^2 \leq \langle \alpha | \alpha \rangle \langle \beta | \beta \rangle \quad (9.382)$$

Assuming this to be true we have

$$\begin{aligned} \Delta A_\Psi^2 \Delta B_\Psi^2 &= \langle \Psi | \hat{\Delta} A_\Psi | \hat{\Delta} A_\Psi | \Psi \rangle \langle \Psi | \hat{\Delta} B_\Psi | \hat{\Delta} B_\Psi | \Psi \rangle \geq \\ &|\langle \Psi | \hat{\Delta} A_\Psi | \hat{\Delta} B_\Psi | \Psi \rangle|^2 = |\langle \Psi | \hat{\Delta} A_\Psi \hat{\Delta} B_\Psi | \Psi \rangle|^2 \end{aligned} \quad (9.383)$$

where we used  $(\hat{\Delta} A_\Psi)^\dagger = \hat{\Delta} A_\Psi$ . Now we notice that for each two operators  $\hat{C}$  and  $\hat{D}$  we have

$$\hat{C}\hat{D} = \frac{1}{2} (\hat{C}\hat{D} - \hat{D}\hat{C} + \hat{C}\hat{D} + \hat{D}\hat{C}) = \frac{1}{2} ([\hat{C}, \hat{D}] + \{\hat{C}, \hat{D}\}) \quad (9.384)$$

where we have defined the *anticommutator*

$$\{\hat{C}, \hat{D}\} = \hat{C}\hat{D} + \hat{D}\hat{C} \quad (9.385)$$

Substituting in eq. (9.383) (replacing the operators  $\hat{C}$  and  $\hat{D}$  with  $\hat{\Delta} A_\Psi$  and  $\hat{\Delta} B_\Psi$ ) we get

$$\begin{aligned} \Delta A_\Psi^2 \Delta B_\Psi^2 &\geq \frac{1}{4} |\langle \Psi | [\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi] + \{\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi\} | \Psi \rangle|^2 = \\ &\frac{1}{4} |\langle \Psi | [\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi] | \Psi \rangle + \langle \Psi | \{\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi\} | \Psi \rangle|^2 \end{aligned} \quad (9.386)$$

Now, since

$$\{\hat{C}, \hat{D}\}^\dagger = (\hat{C}\hat{D} + \hat{D}\hat{C})^\dagger = \hat{D}\hat{C} + \hat{C}\hat{D} = \{\hat{C}, \hat{D}\} \quad (9.387)$$

we have

$$\begin{aligned} \langle \Psi | \{\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi\} | \Psi \rangle &= \langle \Psi | \{\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi\} | \Psi \rangle = (\langle \Psi | \{\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi\} | \Psi \rangle)^* \\ &\Rightarrow \langle \Psi | \{\hat{\Delta} A_\Psi, \hat{\Delta} B_\Psi\} | \Psi \rangle \in \mathbb{R} \end{aligned} \quad (9.388)$$

On the other hand

$$[\hat{C}, \hat{D}]^\dagger = (\hat{C}\hat{D} - \hat{D}\hat{C})^\dagger = \hat{D}\hat{C} - \hat{C}\hat{D} = -[\hat{C}, \hat{D}] \quad (9.389)$$

we have

$$\begin{aligned} \langle \Psi | [\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] | \Psi \rangle &= -\langle \Psi | [\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] | \Psi \rangle = -(\langle \Psi | [\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] | \Psi \rangle)^* \\ &\Rightarrow \langle \Psi | [\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] | \Psi \rangle \in \mathbb{I} \end{aligned} \quad (9.390)$$

But if  $\alpha \in \mathbb{R}$  and  $\beta \in \mathbb{I}$  we have

$$|\alpha + \beta|^2 = (\alpha - \beta)(\alpha + \beta) = \alpha^2 - \beta^2 = \alpha^2 + |\beta|^2 \quad (9.391)$$

so that eq. (9.386) becomes

$$\Delta A_\Psi^2 \Delta B_\Psi^2 \geq \frac{1}{4} |\langle \Psi | [\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] | \Psi \rangle|^2 + \frac{1}{4} |\langle \Psi | \{\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi\} | \Psi \rangle|^2 \geq \frac{1}{4} |\langle \Psi | [\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] | \Psi \rangle|^2 \quad (9.392)$$

But

$$[\hat{\Delta}A_\Psi, \hat{\Delta}B_\Psi] = (\hat{A} - \langle A \rangle_\Psi)(\hat{B} - \langle B \rangle_\Psi) - (\hat{B} - \langle B \rangle_\Psi)(\hat{A} - \langle A \rangle_\Psi) = \hat{A}\hat{B} - \hat{B}\hat{A} \quad (9.393)$$

since  $\langle A \rangle_\Psi$  and  $\langle B \rangle_\Psi$  are just numbers and commute with everything, so that

$$\Delta A_\Psi^2 \Delta B_\Psi^2 \geq \frac{1}{4} |\langle \Psi | [\hat{A}, \hat{B}] | \Psi \rangle|^2 \quad (9.394)$$

### 9.E.2 Schwartz inequality

This inequality is very easy to show. We obviously know that

$$\langle \gamma | \gamma \rangle, \quad \forall |\gamma\rangle \quad (9.395)$$

let us pick

$$|\gamma\rangle = |\alpha\rangle - \frac{\langle \beta | \alpha \rangle}{\langle \beta | \beta \rangle} |\beta\rangle \quad (9.396)$$

so that (using  $\langle \beta | \alpha \rangle^* = \langle \alpha | \beta \rangle$ )

$$\begin{aligned} 0 \leq \langle \gamma | \gamma \rangle &= \left( \langle \alpha | - \frac{\langle \alpha | \beta \rangle}{\langle \beta | \beta \rangle} \langle \beta | \right) |\alpha\rangle - \frac{\langle \beta | \alpha \rangle}{\langle \beta | \beta \rangle} |\beta\rangle = \langle \alpha | \alpha \rangle - \frac{\langle \alpha | \beta \rangle}{\langle \beta | \beta \rangle} \langle \beta | \alpha \rangle - \frac{\langle \beta | \alpha \rangle}{\langle \beta | \beta \rangle} \langle \alpha | \beta \rangle + \frac{|\langle \beta | \alpha \rangle|^2}{\langle \beta | \beta \rangle^2} \langle \beta | \beta \rangle = \\ &= \langle \alpha | \alpha \rangle - \frac{|\langle \beta | \alpha \rangle|^2}{\langle \beta | \beta \rangle^2} \langle \beta | \beta \rangle - \frac{|\langle \beta | \alpha \rangle|^2}{\langle \beta | \beta \rangle^2} \langle \beta | \beta \rangle + \frac{|\langle \beta | \alpha \rangle|^2}{\langle \beta | \beta \rangle^2} \langle \beta | \beta \rangle = \langle \alpha | \alpha \rangle - \frac{|\langle \beta | \alpha \rangle|^2}{\langle \beta | \beta \rangle} \\ &\Rightarrow \langle \alpha | \alpha \rangle \geq \frac{|\langle \beta | \alpha \rangle|^2}{\langle \beta | \beta \rangle} \Rightarrow \langle \alpha | \alpha \rangle \langle \beta | \beta \rangle \geq |\langle \beta | \alpha \rangle|^2 \end{aligned} \quad (9.397)$$

which proves the inequality.

---

## 9.E Mathematical formulation of the uncertainty principle

### 9.E.3 Physical meaning

#### Position and momentum

For the  $\hat{x}$  and  $\hat{p}$  operators we have, for example

$$\Delta x_{\Psi}^2 \Delta p_{x\Psi}^2 \geq \frac{1}{4} |\langle \Psi | [\hat{x}, \hat{p}_x] | \Psi \rangle|^2 = \frac{\hbar^2}{4} \quad (9.398)$$

on all states, since the commutator is a multiple of the identity. What happens if we have an eigenstate of  $p$  with  $\Delta p_{x\Psi}^2 = 0$ ? How can the relation be satisfied? We now that such an eigenstate is in the form

$$\Psi(x) = Ae^{ipx} \quad (9.399)$$

so that the probability (refer to eq. 3.62) of finding the particle in  $x$  is

$$|\Psi(x)|^2 = |A|^2 = \text{const} \quad (9.400)$$

The particle is *everywhere with the same probability* and thus its spread in position is infinite! Actually the physical sense of this mathematical result is that it is impossible to build a physical state with a perfectly defined  $p$  or  $x$ . All physical states will have a spread in position and moment, their product satisfying eq. (9.398). The more we try to localise the particle in space, the less we will know its momentum, and viceversa!

#### Angular momentum

States such as  $|m\rangle$  in which one component of the angular momentum assumes a defined value *do exist*. If then on these states we have  $\Delta L_{z|m}^2 = 0$  how can we satisfy

$$\Delta L_{z|m}^2 \Delta L_{x|m}^2 \geq \frac{1}{4} |\langle m | [\hat{L}_z, \hat{L}_y] | m \rangle|^2 = \frac{\hbar^2}{4} |\langle m | \hat{L}_y | m \rangle|^2 \quad (9.401)$$

The solution is easy. We know that

$$\hat{L}_y = \frac{-i}{2} (\hat{L}_+ - \hat{L}_-) = \frac{-i}{2} (\hat{L}_x + i\hat{L}_y - \hat{L}_x + i\hat{L}_y) \quad (9.402)$$

so that the action of  $\hat{L}_y$  on  $|m\rangle$  creates a combination of  $|m+1\rangle$  and  $|m-1\rangle$  (but no  $|m\rangle$ ) that have 0 scalar product with  $\langle m|$ , so that

$$\langle m | \hat{L}_y | m \rangle = 0 \quad (9.403)$$

# Bibliography